



US008835445B2

(12) **United States Patent**
Chen et al.(10) **Patent No.:** US 8,835,445 B2(45) **Date of Patent:** Sep. 16, 2014(54) **DIHYDROFOLATE REDUCTASE
INHIBITORS**

FOREIGN PATENT DOCUMENTS

(75) Inventors: **Zhiyong Chen**, San Diego, CA (US);
Christopher J. Creighton, San Diego,
CA (US); **Mark Cunningham**, San
Diego, CA (US); **John Finn**, San Diego,
CA (US); **Mark Hilgers**, San Diego, CA
(US); **Michael Jung**, Los Angeles, CA
(US); **Lucy Aguirre Kohnen**,
Cambridge, MA (US); **Thanh Lam**, San
Diego, CA (US); **Xiaoming Li**, San
Diego, CA (US); **Mark Stidham**, San
Diego, CA (US); **Les Tari**, San Diego,
CA (US); **Michael Trzoss**, San Diego,
CA (US); **Junhu Zhang**, San Diego, CA
(US)WO WO 94/18980 9/1994
WO WO 2006/050843 5/2006

OTHER PUBLICATIONS

(73) Assignee: **Trius Therapeutics, Inc.**, San Diego,
CA (US)(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 73 days.(21) Appl. No.: **13/151,683**(22) Filed: **Jun. 2, 2011**(65) **Prior Publication Data**

US 2012/0136014 A1 May 31, 2012

Related U.S. Application Data(60) Provisional application No. 61/350,886, filed on Jun.
2, 2010.(51) **Int. Cl.****A01N 43/54** (2006.01)
A61K 31/517 (2006.01)
C07D 239/72 (2006.01)
C07D 401/00 (2006.01)(52) **U.S. Cl.**USPC **514/266.1**; 514/266.2; 544/283;
544/284(58) **Field of Classification Search**USPC 544/283, 284; 540/484; 514/266.1,
514/266.2

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,534,518 A 7/1996 Henrie et al.
7,678,803 B2* 3/2010 Huang et al. 514/266.2
2004/0229890 A1 11/2004 Berthel et al.
2006/0211715 A1 9/2006 Berthel et al.
2008/0070935 A1* 3/2008 Huang et al. 514/264.11
2009/0118311 A1 5/2009 Davidson
2009/0312305 A1 12/2009 Beard et al.


Vippagunta et al.*

Finn et al. Structure-Based Design of New DHFR-based Antibacte-
rial Agents (Part 2): (Benzimidazol-1-yl)-2,4-diaminoquinazolines
SAR, retrieved from [http://www.triusrx.com/pdfs/Structure-Based-Design-of-New-DHFR-Antibacterial-Agents\(Part%202\).pdf](http://www.triusrx.com/pdfs/Structure-Based-Design-of-New-DHFR-Antibacterial-Agents(Part%202).pdf) (on
Apr. 5, 2012).Rosowsky and Papatanasopoulos "Quinazolines. XIII. Synthesis of
Polycyclic 2,4-Diaminopyrimidines from Aromatic Amine Hydro-
chlorides and Sodium Dicyanamide 1,2," J. Org. Chem., 39(22):
3293-5 (1974).Cikotiene and Morkunas, "The First and Efficient Synthesis of
7-Aryl-6-methoxycarbonylquinazolines via Unexpected Reaction of
6-Arylethynylpyrimidine-5-carbaldehydes and Methyl
Mercaptoacetate," Synlett (2): 284-286 (2009).Chemical Abstracts Service, Columbus Ohio, US: CAS Registry No.
1027917-45-8 (Jun. 1, 2011); CAS Registry No. 1137439-84-9 (Jun.
1, 2011); CAS Registry No. 159018-81-2 (Jun. 1, 2011); CAS Reg-
istry No. 180268-99-9 (Jun. 1, 2011); CAS Registry No. 52306-21-5
(Jun. 1, 2011); CAS Registry No. 887232-80-6 (Jun. 1, 2011); CAS
Registry No. 887232-82-8 (Jun. 1, 2011); CAS Registry No. 887232-
91-9 (Jun. 1, 2011); CAS Registry No. 887232-96-4 (Jun. 1, 2011);
CAS Registry No. 887233-15-0 (Jun. 1, 2011); CAS Registry No.
887233-17-2 (Jun. 1, 2011); CAS Registry No. 887233-19-4 (Jun. 1,
2011); CAS Registry No. 887233-21-8 (Jun. 1, 2011); CAS Registry
No. 887233-26-3 (Jun. 1, 2011); CAS Registry No. 887233-28-5
(Jun. 1, 2011).

* cited by examiner

Primary Examiner — Paul V. Ward(74) *Attorney, Agent, or Firm* — DLA Piper LLP (US)(57) **ABSTRACT**

The present disclosure provides compounds of Formula I:

or a pharmaceutically acceptable salt thereof, wherein R⁵, R⁶
and Z are as described herein. The disclosure also provides
pharmaceutical compositions thereof; and methods for inhib-
iting DHFR activity; and methods for treating cell prolifera-
tive diseases, autoimmune disease, inflammatory disease or
bacterial, fungal or parasitic infection by administering a
compound of Formula I.**15 Claims, No Drawings**

1

**DIHYDROFOLATE REDUCTASE
INHIBITORS****CROSS REFERENCE TO RELATED
APPLICATIONS**

This application claims the benefit of priority under 35 U.S.C. §119(e) to U.S. Provisional Application No. 61/350,886 filed on Jun. 2, 2010, the disclosure of which is hereby incorporated by reference in their entirety for all purposes.

FIELD OF THE INVENTION

The disclosure relates to compounds and compositions that inhibit the dihydrofolate reductase (DHFR) family of enzymes; methods of making and using these compounds and compositions for the treatment and prophylaxis of diseases in mammals including the treatment of cell proliferative diseases such as cancers and inflammation, and to their use as anti-infective agents for bacterial, fungal or parasitic infection.

BACKGROUND OF THE INVENTION

Bacterial infections pose a continuing medical problem because anti-bacterial drugs eventually engender resistance in the bacteria on which they are used. Consequently, a need exists for new drugs with efficacy against pathogenic bacteria for use in the therapy and prophylaxis of bacterial infections.

One target for development of anti-bacterial drugs has been dihydrofolate reductase (DHFR), an enzyme on the synthetic pathway to purine and pyrimidine nucleotides. DHFR inhibitors have been disclosed in US 20090118311, which is hereby incorporated by reference in its entirety.

Folate (pteroylglutamate) is a vitamin which is a key component in the biosynthesis of purine and pyrimidine nucleotides. Following absorption, dietary folate is reduced to dihydrofolate and then further reduced to tetrahydrofolate by the enzyme dihydrofolate reductase (DHFR). Inhibition of DHFR leads to a reduction in nucleotide biosynthesis resulting in inhibition of DNA biosynthesis and reduced cell division. DHFR inhibitors are widely used in the treatment of cancer, cell proliferative diseases such as rheumatoid arthritis, psoriasis and transplant rejection. DHFR inhibitors have also found use as anti-infective and anti-parasitic agents. Many types of DHFR inhibitor compounds have been suggested, and several such compounds are used as anti-cancer, anti-inflammatory, anti-infective and anti-parasitic agents.

Methotrexate is the most widely used DHFR inhibitor, which contains a glutamate functionality that enables it to be actively transported into and retained inside of cells. However, cancer cells can become resistant to methotrexate by modifying this active transport mechanism. Furthermore, non-mammalian cells lack the active transport system and methotrexate has limited utility as an anti-infective agent. Lipophilic DHFR inhibitors which can be taken up by passive diffusion into cells have therefore been developed both to circumvent cancer cell resistance and for use as anti-infective agents. However, agents that passively diffuse into cells also exit the cell readily. Thus, there remains a need in the art for DHFR inhibitors that accumulate in cells in a way that does not depend on the active transport mechanism of methotrexate or on the lipophilicities.

SUMMARY OF THE INVENTION


The disclosure addresses these needs by providing new compounds and pharmaceutical compositions thereof, for

2

inhibiting DHFR activity and for treating cell proliferative diseases, autoimmune diseases, inflammatory diseases or bacterial, fungal or parasitic infections. Thus, in one embodiment the disclosure provides compounds of Formula I

5

Formula I



10

15

wherein:

Z is a five to 10-membered aryl or heteroaryl ring compound comprising 0-4 heteroatoms independently selected from the group consisting of N, O, and S;

20

R represents one or more moieties independently selected from the group consisting of H, halo, lower alkyl, lower alkoxy, CF₃, amino, CN, and optionally-substituted aryl, and optionally-substituted heterocyclyl;

25

R⁵ is selected from the group consisting of H, halo, lower alkyl, and lower alkoxy;

30

R⁶ is selected from the group consisting of H, halo, lower alkyl, lower alkoxy, CF₃, 5-membered heteroarylmethylene, and amino; and

35

pharmaceutically acceptable salts, solvates, and prodrugs thereof.

40

The disclosure also provides pharmaceutical compositions of the compound of Formula I, and methods for inhibiting DHFR activity and for treating cell proliferative diseases, autoimmune diseases, inflammatory diseases or bacterial, fungal or parasitic infections using the compound of Formula I.

45

DETAILED DESCRIPTION OF THE INVENTION

Unless otherwise specified, technical terms take the meanings specified in the McGraw-Hill Dictionary of Scientific and Technical Terms, 6th edition.

50

Definitions

“Alkyl” refers to substituents derived from optionally-substituted saturated hydrocarbons, whether cyclic or acyclic, by removal of one or more hydrogen atoms.

55

“Lower alkyl” refers to optionally-substituted acyclic alkyl groups having one to six carbon atoms, and to cyclic alkyl groups having three to six carbon atoms.

Similarly, “lower alkoxy” refers to —OR groups where R comprises a chain of one to six carbon atoms.

60

“Aryl” refers to optionally-substituted monocyclic and fused bicyclic carbocyclic groups having from five to 12 carbon atoms and having at least one aromatic ring. Examples of particular aryl groups include phenyl and naphthyl.

65

“Heterocyclic,” “heterocycle,” and “heterocyclyl” refer to optionally-substituted monocyclic and fused bicyclic groups, saturated or unsaturated, aromatic or non-aromatic, having the specified number of members and including 1-4 heteroatoms selected from N, O and S. Examples include tetrahydrofuran, dihydropyran, tetrahydropyran, pyran, oxetane, thietane, 1,4-dioxane, 1,3-dioxane, 1,3-dioxalane, piperidine, piperazine, tetrahydropyrimidine, pyrrolidine, morpholine, thiomorpholine, thiazolidine, oxazolidine, tetrahydrothiopyran, and tetrahydrothiophene.

ment of the patients, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

Preferably these compositions are in unit dosage forms from such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories; for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition

is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

In one aspect, the disclosure provides a compound of Formula I, wherein the compound is selected from Table 1.

TABLE 1




#	Structure	Name
1		7-(2-methoxyphenyl)quinazoline-2,4-diamine
2		7-(2-aminophenyl)quinazoline-2,4-diamine
3		7-(2,5-dimethoxyphenyl)quinazoline-2,4-diamine

TABLE 1-continued







#	Structure	Name
4		7-(4-aminophenyl)quinazoline-2,4-diamine
5		7-(4-aminophenyl)-6-methylquinazoline-2,4-diamine
6		1-(3-(2,4-diaminoquinazolin-7-yl)-phenyl)ethanone
7		1-(3-(2,4-diamino-6-methylquinazolin-7-yl)-phenyl)ethanone
8		7-(2,5-dimethoxyphenyl)-6-methylquinazoline-2,4-diamine
9		7-(1H-indol-4-yl)-6-methylquinazoline-2,4-diamine

TABLE 1-continued

#	Structure	Name
10		6-((1H-imidazol-1-yl)methyl)-7-(3,4-dimethoxyphenyl)quinazoline-2,4-diamine
11		3-(2,4-diamino-6-methylquinazolin-7-yl)benzotrile
12		3-amino-5-(2,4-diamino-6-methylquinazolin-7-yl)benzotrile
13		2-(3-(2,4-diamino-6-methylquinazolin-7-yl)phenyl)acetotrile
14		7-(1-(2-methoxyethyl)-5,6-dimethyl-1H-benzo[d]imidazol-2-yl)quinazoline-2,4-diamine

TABLE 1-continued





#	Structure	Name
15		7-(2-cyclopropyl-5,6-dimethyl-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
16		7-(5,6-dimethyl-2-(pyridin-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
17		7-(2-(furan-2-yl)-5,6-dimethoxy-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
18		7-(5,6-dimethyl-2-(thiophen-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine

TABLE 1-continued

#	Structure	Name
19		7-(5,6-dimethyl-2-morpholino-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
20		7-(2-(3,3-difluorobutylthio)-6-methoxy-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
21		7-(5,6-dimethyl-2-(thiophen-3-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
22		7-[5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl]quinazoline-2,4-diamine
23		6-chloro-7-(5,6-dimethoxy-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine

TABLE 1-continued

#	Structure	Name
24		7-(5,6-dimethoxy-2-(thiazol-2-yl)-1H-benzo[d]-imidazol-1-yl)-6-methylquinazoline-2,4-diamine
25		6-chloro-7-(5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
26		7-(5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]-imidazol-1-yl)-6-methylquinazoline-2,4-diamine
27		6-methyl-7-(1H-pyrrolo[2,3-b]pyridin-5-yl)quinazoline-2,4-diamine
28		7-(5-chloro-2-methoxypyridin-4-yl)-6-methylquinazoline-2,4-diamine

TABLE 1-continued









#	Structure	Name
29		7-(2,4-dimethoxypyrimidin-5-yl)-6-methylquinazoline-2,4-diamine
30		N-(3'-(2,4-diaminoquinazolin-7-yl)-4'-ethoxybiphenyl-3-yl)methanesulfonamide
31		7-(2-ethoxynaphthalen-1-yl)-6-methylquinazoline-2,4-diamine
32		methyl 1-(2,4-diaminoquinazolin-7-yl)-2-(thiazol-2-yl)-1H-benzo[d]imidazole-5-carboxylate
33		6-methyl-7-(3,4,5-trimethoxyphenyl)quinazoline-2,4-diamine

TABLE 1-continued

#	Structure	Name
34		6-chloro-7-(5,6-dimethyl-2-(thiazol-2-yl))-1H-benzo[d]imidazol-1-ylquinazoline-2,4-diamine
35		7-(5,6-dimethoxy-2-(thiazol-2-yl))-1H-benzo[d]imidazol-1-ylquinazoline-2,4-diamine
36		7-(5-methyl-2-(thiophen-2-yl))-1H-benzo[d]imidazol-1-ylquinazoline-2,4-diamine
37		7-(2-furan-2-yl)-5-methoxy-3H-imidazo[4,5-b]pyridine-3-ylquinazoline-2,4-diamine
38		7-(2-chloro-5-methoxy-4-pyridin-4-yl)-6-methylquinazoline-2,4-diamine

TABLE 1-continued

#	Structure	Name
39		tert-butyl 1-(2,4-diaminoquinazolin-7-yl)-5-methyl-2-thiophen-2-yl)-1H-imidazole-4-carboxylate
40		3-(2,4-diamino-6-methylquinazolin-7-yl)-4-ethoxy-N,N-diethylbenzamide
41		7-(1H-indol-3-yl)-6-methylquinazolin-2,4-diamine

In another aspect, the disclosure provides a pharmaceutical composition of a compound of Formula I together with a pharmaceutically acceptable carrier.

In another aspect, the disclosure provides methods for impeding growth of a bacterium by exposing the bacteria to a compound of Formula I.

In another aspect, the disclosure provides methods for treating a mammal prophylactically or therapeutically for a bacterial infection by administering to the mammal a compound of Formula I.

In another aspect, the disclosure provides methods for treating a mammal prophylactically or therapeutically for a bacterial infection comprising administering to the mammal a pharmaceutical composition of a compound of Formula I together with a pharmaceutically acceptable carrier.

In another aspect, the disclosure provides methods for inhibiting DHFR activity by contacting the DHFR enzyme with an amount of a compound of Formula I effective for such inhibition.

In another aspect, the disclosure provides methods for treating cell proliferative diseases, autoimmune disease, inflammatory disease or bacterial, fungal or parasitic infection, by administering to a subject in need thereof, an effective amount of a compound of Formula I.

In another aspect, the disclosure provides methods for treating cell proliferative diseases, autoimmune disease, inflammatory disease or bacterial, fungal or parasitic infection, by administering to a subject in need thereof, an effective


amount of a compound of Formula I, wherein the disease is cancer, rheumatoid arthritis, or fungal, bacterial or parasitic infection.

EXAMPLES

Some aspects of the disclosure can be further illustrated by the following non-limiting examples.


General Procedure for the Preparation of Diaminoquinazolines:

General procedures for the preparation of the diaminoquinazoline




23

-continued




24


-continued



Scheme 2




Scheme 4






Specific examples of diaminoquinazoline compounds prepared by these methods are provided in Schemes 5-7.

Scheme 3



Scheme 5





Example 1

Preparation of 3'-(cyanomethyl)-5-fluoro-2-methylbiphenyl-4-carbonitrile (3)

Tetrakis(triphenylphosphine)palladium (10 mg, 0.1 mmol) was added into the solution of 4-bromo-2-fluoro-5-methylbenzonitrile (110 mg, 0.5 mmol), K_2CO_3 70% aqueous solution (1 mL) and 3-(cyanomethyl)phenylboronic acid (96 mg, 0.6 mmol) in 10 ml N,N-dimethyl formamide, the solution was heated at 150° C. under microwave condition for 20 minutes, and the reaction mixture was partitioned with 20 mL water and 50 mL ethyl acetate, the aqueous layer was extracted with ethyl acetate (100 mL \times 2), and the combined organic layer was washed by saturated solution of sodium bicarbonate, sodium chloride, and dried by sodium sulfate, and concentrated by give quantitative products 3'-(cyanomethyl)-5-fluoro-2-methylbiphenyl-4-carbonitrile 3 (114 mg, 88% yield, confirmed by LCMS: 251.10) without purification for next steps.




Example 2

Preparation of 2-(3-(2-(4-(cyanomethyl)phenyl)-5-fluoro-2-methylquinazolin-7-yl)phenyl)acetonitrile (4)

To the above solution of 3'-(cyanomethyl)-5-fluoro-2-methylbiphenyl-4-carbonitrile 3 was added guanidine carbonate (180 mg, 0.5 mmol) and N,N-dimethyl formamide (5 mL). The reaction was heated at 150° C. under microwave condition for 20 minutes. The mixture was purified by high performance liquid chromatography to yield compound 4 as white solid trifluoroacetic acid salt (89 mg, Yield: 68%, confirmed by LCMS: 290.15 and HPLC).

26

Scheme 6



Example (3)

Preparation of 4-(5,6-dimethyl-1H-benzimidazol-2-yl)-2-fluorobenzonitrile (3)

To a solution of 2-fluoro-4-formylbenzonitrile 1 (100 mg, 0.7 mmol) and 4,5-dimethyl benzene-1,2-diamine 2 (109 mg, 0.8 mmol) in ethanol (10 ml) was added a catalytic amount of p-toluenesulfonic acid (0.5 mg, 0.5 mmol). The mixture was stirred reflux for 2 hours and concentrated by rotor evaporation. The crude material was purified through column chromatography with 40% ethyl acetate in hexane to yield compound 3 as white solid (68% yield, 121 mg, confirmed by LCMS: 266.20).

27

Example (4)

Preparation of 2-fluoro-4-(1-(2-methoxyethyl)-5,6-dimethyl-1H-benzo[d]imidazol-2-yl)benzonitrile (4)


4-(5,6-dimethyl-1H-benzo[d]imidazol-2-yl)-2-fluorobenzonitrile 3 (110 mg, 0.4 mmol) was combined with 1-bromo-2-methoxyethane (76 mg, 0.55 mmol), NaH (13 mg, 0.55 mmol), and N,N-dimethyl formamide (5 ml). The reaction was heated at 100° C. for 1.5 hours. The product solution (confirmed by LCMS: 324.22) was carried on to the next step without purification.

Example 5

Preparation of 2-fluoro-4-(1-(2-methoxyethyl)-5,6-dimethyl-1H-benzo[d]imidazol-2-yl)benzonitrile (5)


To the above solution of 2-fluoro-4-(1-(2-methoxyethyl)-5,6-dimethyl-1H-benzo[d]imidazol-2-yl)benzonitrile 4 was added guanidine carbonate (180 mg, 0.5 mmol) and N,N-dimethyl formamide (5 mL). The reaction was heated at 150° C. under microwave condition for 20 minutes. The mixture was purified by high performance liquid chromatography to yield compound 5 as white solid trifluoroacetic acid salt (96 mg, Yield: 64%, confirmed by LCMS: 363.15 and HPLC).

Scheme 7



28

-continued



Example 6

General Method for the Preparation of 7-(benzimidazol-1-yl)-2,4-diaminoquinazolines

The Preparation of 2-(5,6-dimethyl-1H-benzo[d]imidazol-2-yl)thiazole (3): To a well-dissolved solution of 4,5-dimethyl-1,2-phenylenediamine (1.382 g, 10 mmol) in methanol (75 ml) was added catalytic amount of acetic acid (121 mg, 2 mmol) followed by the drop-wise addition over 30 minutes of the solution of 2-thiazole carboxaldehyde (1.25 g, 11 mmol) in methanol (25 ml). The resulting mixture was stirred at room temperature while oxygen was gently bubbling through from one-layer balloon for two days. The solvent was then removed by rotor evaporation. The residue was purified through reverse phase chromatography (150 g Teledyne HP C18 column; 35% of CH₃CN (0.1% TFA) in water (0.1% TFA). Acetonitrile in the combined eluates was removed by rotor evaporation, and the resulting aqueous solution was neutralized by sodium bicarbonate. It was then extracted with dichloromethane (150 ml×3). The combined organic layers were dried over magnesium sulfate, and the solvent was taken off by rotor evaporation to afford the title compound as yellow solid (69.4% yield, 2.08 g, confirmed by LCMS: 230.09).

Example 7

Preparation of 4-(5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl)-2-fluorobenzonitrile (5)

2-(5,6-Dimethyl-1H-benzo[d]imidazol-2-yl)thiazole 3 (174 mg, 0.5 mmol) was combined with 2,4-difluorobenzonitrile (76.5 mg, 0.55 mmol), potassium carbonate (139 mg, 1 mmol), and anhydrous N,N-dimethyl formamide (5 mL). The reaction was heated at 100° C. under microwave condition for 3 hours. The crude product solution (confirmed by LCMS: 349.13) was carried on to the next step without purification.

Example 8

Preparation of 7-(5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine (6)

To the above solution of 4-(5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl)-2-fluorobenzonitrile 5 was added guanidine carbonate (360.4 mg, 2 mmol), potassium carbonate (552.8 mg, 4 mmol), and anhydrous N,N-dimethyl formamide (2 mL). The reaction was heated at 140° C. under microwave condition for 20 minutes. The mixture was puri-

29

fied by high performance liquid chromatography to yield compound 6 as white solid trifluoroacetic acid salt (80 mg, Yield: 60%, confirmed by LCMS: 388.43 and HPLC).


Example 9

General Method for the Preparation of 7-aryl-2,4-diaminoquinazolines: (Scheme 4)

Tetrakis(triphenylphosphine)palladium (10 mg, 0.1 mmol) was added into the solution of 7-bromo-6-methylquinazolin-2,4-diamine (126 mg, 0.5 mmol), K_2CO_3 70% aqueous solution (1 mL) and substituted arylboronic acid (0.6 mmol) in 10 ml N,N-dimethyl formamide, the solution was heated at 150° C. under microwave condition for 20 minutes, and the reaction mixture was partitioned with 20 mL water and 50 mL ethyl acetate, the aqueous layer was extracted with ethyl acetate (100 mL \times 2), and the combined organic layer was washed by saturated solution of sodium bicarbonate, sodium chloride, and dried by sodium sulfate, concentrated and the residue was subjected to pre-HPLC separation to give the desired compounds.



Example 10

Preparation of tert-butyl 1-(2,4-diaminoquinazolin-7-yl)-5-methyl-2-thiophen-2-yl)-1H-imidazole-4-carboxylate (compound 39)



30

-continued



tert-butyl 2-(hydroxyimino)-3-oxobutanoate 1 (1.85 g, 9.89 mmol) was mixed with thiophen-2-ylmethanamine 2 (1.07 mL, 10.44 mmol) in anhydrous acetonitrile (30 mL). The resulting mixture was heated to reflux for 3 hrs, upon cooling, the suspension was filtered, and the filtered material was washed with a small amount of acetonitrile to afford the desired solid product 3 (LC-MS: 265.10). Compound 3 was converted to tert-butyl 1-(2,4-diaminoquinazolin-7-yl)-5-methyl-2-thiophen-2-yl)-1H-imidazole-4-carboxylate (compound 39) by the method described in Examples 6-9.

Further examples of diaminoquinazoline compounds prepared by these methods are provided in

TABLE 2

#	Structure	Name
1		7-(2-methoxyphenyl)quinazoline-2,4-diamine

TABLE 2-continued








#	Structure	Name
2		7-(2-aminophenyl)quinazoline-2,4-diamine
3		7-(2,5-dimethoxyphenyl)quinazoline-2,4-diamine
4		7-(4-aminophenyl)quinazoline-2,4-diamine
5		7-(4-aminophenyl)-6-methylquinazoline-2,4-diamine
6		1-(3-(2,4-diaminoquinazolin-7-yl)-phenyl)ethanone
7		1-(3-(2,4-diamino-6-methylquinazolin-7-yl)-phenyl)ethanone
8		7-(2,5-dimethoxyphenyl)-6-methylquinazoline-2,4-diamine

TABLE 2-continued

#	Structure	Name
9		7-(1H-indol-4-yl)-6-methylquinazoline-2,4-diamine
10		6-((1H-imidazol-1-yl)methyl)-7-(3,4-dimethoxyphenyl)quinazoline-2,4-diamine
11		3-(2,4-diamino-6-methylquinazolin-7-yl)benzotrile
12		3-amino-5-(2,4-diamino-6-methylquinazolin-7-yl)benzotrile
13		2-(3-(2,4-diamino-6-methylquinazolin-7-yl)phenyl)acetone
14		7-(1-(2-methoxyethyl)-5,6-dimethyl-1H-benzo[d]imidazol-2-yl)quinazoline-2,4-diamine

TABLE 2-continued





#	Structure	Name
15		7-(2-cyclopropyl-5,6-dimethyl-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
16		7-(5,6-dimethyl-2-(pyridin-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
17		7-(2-(furan-2-yl)-5,6-dimethoxy-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
18		7-(5,6-dimethyl-2-(thiophen-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine

TABLE 2-continued

#	Structure	Name
19		7-(5,6-dimethyl-2-morpholino-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
20		7-(2-(3,3-difluorobutylthio)-6-methoxy-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
21		7-(5,6-dimethyl-2-(thiophen-3-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
22		7-[5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl]quinazoline-2,4-diamine
23		6-chloro-7-(5,6-dimethoxy-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine

TABLE 2-continued

#	Structure	Name
24		7-(5,6-dimethoxy-2-(thiazol-2-yl)-1H-benzo[d]-imidazol-1-yl)-6-methylquinazoline-2,4-diamine
25		6-chloro-7-(5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
26		7-(5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]-imidazol-1-yl)-6-methylquinazoline-2,4-diamine
27		6-methyl-7-(1H-pyrrolo[2,3-b]pyridin-5-yl)quinazoline-2,4-diamine
28		7-(5-chloro-2-methoxypyridin-4-yl)-6-methylquinazoline-2,4-diamine

TABLE 2-continued









#	Structure	Name
29		7-(2,4-dimethoxypyrimidin-5-yl)-6-methylquinazoline-2,4-diamine
30		N-(3'-(2,4-diaminoquinazolin-7-yl)-4'-ethoxybiphenyl-3-yl)methanesulfonamide
31		7-(2-ethoxynaphthalen-1-yl)-6-methylquinazoline-2,4-diamine
32		methyl 1-(2,4-diaminoquinazolin-7-yl)-2-(thiazol-2-yl)-1H-benzo[d]imidazole-5-carboxylate
33		6-methyl-7-(3,4,5-trimethoxyphenyl)quinazoline-2,4-diamine

TABLE 2-continued

#	Structure	Name
34		6-chloro-7-(5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
35		7-(5,6-dimethoxy-2-thiazol-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
36		7-(5-methyl-2-thiophen-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine
37		7-(2-furan-2-yl)-5-methoxy-3H-imidazol[4,5-b]pyridine-3-yl)quinazoline-2,4-diamine
38		7-(2-chloro-5-methoxy-3-pyridin-4-yl)-6-methylquinazoline-2,4-diamine

TABLE 2-continued

#	Structure	Name
39		tert-butyl 1-(2,4-diaminoquinazolin-7-yl)-5-methyl-2-thiophen-2-yl)-1H-imidazole-4-carboxylate
40		3-(2,4-diamino-6-methylquinazolin-7-yl)-4-ethoxy-N,N-diethylbenzamide
41		7-(1H-indol-3-yl)-6-methylquinazolin-2,4-diamine

Determination of Anti-Bacterial Efficacy

Antibacterial Activity Assays

Antibacterial activity as measured by the minimal inhibitory concentrations (MIC) and minimal bactericidal concentrations of compounds are well known (see., e.g., National Committee for Clinical Laboratory Standards 2000 Performance standards for antimicrobial disk susceptibility tests: approved standard, 7th ed. M2-A7, vol. 20, no. 1, Committee for Clinical Laboratory Standards, Wayne, Pa.)

In vitro testing for antibacterial activity may be accomplished through use of a whole-cell bacterial growth inhibition assay. For example, an agar dilution assay identifies a substance that inhibits bacterial growth. Microtiter plates are prepared with serial dilutions of the test compound, adding to the preparation a given amount of growth substrate, and providing a preparation of bacteria. Inhibition of bacterial growth is determined, for example, by observing changes in optical densities of the bacterial cultures. Inhibition of bacterial growth is determined, for example, by comparing (in the presence and absence of a test compound) the rate of growth or the absolute growth of bacterial cells. Inhibition includes a reduction of one of the above measurements by at least 20%.

The compounds of the present invention are active against a wide range of bacteria. In some embodiments, the bacteria are Gram-positive bacteria including methicillin-susceptible and methicillin-resistant Staphylococci (including *Staphylo-*

coccus aureus, *Staphylococcus epidermidis*, *Staphylococcus haemolyticus*, *Staphylococcus hominis*, *Staphylococcus saprophyticus*, and coagulase-negative Staphylococci), glycopeptide intermediary-susceptible *Staphylococcus aureus* (GISA), penicillin-susceptible and penicillin-resistant Streptococci (including *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus avium*, *Streptococcus bovis*, *Streptococcus lactis*, *Streptococcus sangius* and Streptococci Group C, Streptococci Group G and viridans Streptococci), enterococci (including vancomycin-susceptible and vancomycin-resistant strains such as *Enterococcus faecalis* and *Enterococcus faecium*), *Bacillus anthracis*, *Clostridium difficile*, *Clostridium clostridioforme*, *Clostridium innocuum*, *Clostridium perfringens*, *Clostridium ramosum*, *Haemophilus influenzae*, *Listeria monocytogenes*, *Corynebacterium jeikeium*, *Bifidobacterium* spp., *Eubacterium aerofaciens*, *Eubacterium lentum*, *Lactobacillus acidophilus*, *Lactobacillus casei*, *Lactobacillus plantarum*, *Lactococcus* spp., *Leuconostoc* spp., *Pediococcus*, *Peptostreptococcus anaerobius*, *Peptostreptococcus asaccharolyticus*, *Peptostreptococcus magnus*, *Peptostreptococcus micros*, *Peptostreptococcus prevotii*, *Peptostreptococcus productus*, *Propionibacterium acnes*, and *Actinomyces* spp. In some embodiments, the Gram-positive bacterium is *Staphylococcus aureus*.

47


TABLE 3

Enzymatic and Antibacterial Activity of Compounds 1-41				
Compound #	MIC in $\mu\text{g/ml}$		DHFR Ki (nM)	
	<i>S. aureus</i>	<i>S. aureus</i> in 20% serum	human DHFR	<i>S. aureus</i> DHFR
1	2	4	1138	2.7
2	8	8	3922	15.3
3	4	8	345	4
4	4	2	702	2.7
5	1	1	138	0.89
6	4	8	323	4.4
7	0.5	1	26.3	0.48
8	1	1	27.5	0.66
9	0.25	1	245	0.61
10	8	2	53.6	0.51
11	1	2	55	1.2
12	1	1	44	0.65
13	0.5	2	44	0.5
14	4	4	263	1.6
15	≤ 0.5	1	1164	0.14
16	≤ 0.125	0.25	572	0.022
17	≤ 0.125	0.25	943	0.007
18	≤ 0.125	0.5	1026	0.005
19	1	4	1086	0.6
20	0.5	4	586	0.12
21	≤ 0.125	0.25	663	0.012
22	≤ 0.125	≤ 0.125	93.5	0.002
23	≤ 0.125	0.5	320	0.011
24	≤ 0.5	≤ 0.5	170	0.019
25	≤ 0.125	1	7.6	0.011
26	≤ 0.125	≤ 0.125	2.9	0.03
27	4	4	235	2.6
28	0.5	1	56	0.72
29	0.5	0.5	54.7	0.4
30	≤ 0.5	2	5.8	0.02
31	≤ 0.125	0.5	4.5	0.026
32	0.25	2	53.1	0.021
33	≤ 0.5	1	47.1	0.91
34	≤ 0.125	2	8.8	0.016
35	0.25	0.25	854	0.16
36	1	1	314	0.005
37	0.25	0.5	797	0.039
38	1	1	52.4	1.4
39	1	2	1913	0.2
40	0.25	0.25	34	0.025
41	0.25	0.5	50.5	0.11

Although the disclosure has been described with reference to the above examples, it will be understood that modifications and variations are encompassed within the spirit and scope of the disclosure. Accordingly, the disclosure is limited only by the following claims.

The invention claimed is:

1. A compound having the structure



wherein

Z is a seven to 10-membered heteroaryl ring comprising 0-4 heteroatoms independently selected from the group consisting of N, O, and S;

when two rings comprise the seven to 10-membered heteroaryl ring, the two rings are heteroaromatic;

48

R represents one or more moieties independently selected from the group consisting of H, halo, lower alkyl, lower alkoxy, CF_3 , amino, CN, CO_2CH_3 , optionally-substituted aryl, and optionally-substituted heterocyclyl;

5 R^5 is selected from the group consisting of H, halo, lower alkyl, and lower alkoxy;

10 R^6 is selected from the group consisting of H, halo, lower alkyl, lower alkoxy, CF_3 , 5-membered heteroarylmethylene, and amino; and pharmaceutically acceptable salts and solvates thereof.

2. The compound of claim 1, wherein Z is selected from the group consisting of optionally-substituted isoquinoline, benzofuran, benzothiophene, indole, pyrrolopyridine, imidazopyridine, indazole benzimidazole, benzothiazole, quinoline, and benzotriazole.

3. The compound of claim 2, wherein Z is optionally-substituted benzimidazole.


4. The compound of claim 1, wherein R is selected from the group consisting of halo, optionally-substituted lower alkyl, lower alkoxy, and heterocyclyl.

5. The compound of claim 1, wherein R is selected from the group consisting of Me, Et, i-Pr, Cl, F, MeO, cyano, and CF_3 .

25 6. The compound of claim 1, wherein R is an optionally-substituted heteroaryl group.

7. The compound of claim 1, wherein R^5 and R^6 are independently selected from the group consisting of F, Cl, lower alkyl, OMe, CF_3 , NMe_2 , and imidazolylmethyl.


30 8. The compound of claim 1, where the compound is



35 7-(5,6-dimethyl-2-(pyridin-2-yl)-1H-benzo[d]-imidazole-1-yl)quinazoline-2,4-diamine.

9. The compound of claim 1, where the compound is


(Formula I)



55 7-[5,6-dimethyl-2-(thiazol-2-yl)-1H-benzo[d]-imidazole-1-yl]quinazoline-2,4-diamine.


49

10. The compound of claim 1, where the compound is



7-(5,6-dimethoxy-2-(thiazol-2-yl)-1H-benzo[d]imidazol-1-yl)quinazoline-2,4-diamine.

11. The compound of claim 1, where the compound is



50

7-(2-(furan-2-yl)-5,6-dimethoxy-1H-benzo[d]-imidazol-1-yl)quinazoline-2,4-diamine.

12. The compound of claim 1, where the compound is

5


10

15

20

25

30



7-(5,6-dimethyl-2-(thiophen-2-yl)-1H-benzo[d]-imidazol-1-yl)quinazoline-2,4-diamine.

13. The compound of claim 1, where the seven to 10-membered heteroaryl ring is an eight to 10-membered heteroaryl ring.

14. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable diluent or carrier.

15. A method of impeding growth of a bacterium by exposing the bacterium to a compound of claim 1.

* * * * *