

Chapter 10

Trace Metal Utilization in Chloroplasts

Sabeeha Merchant*

Department of Chemistry and Biochemistry, University of California, Los Angeles,
CA 90095, U.S.A.

Summary	200
I. Introduction	200
A. Transition Metals Function as Redox Catalysts	200
B. Trace Metal Deficiency Impacts the Chloroplast	201
C. Metalloprotein Assembly—Thermodynamics vs. Kinetics	201
D. Fe, Cu and Mn	201
II. Fe	202
A. Ferritin	202
B. Heme, FeS and Fe Cofactor Synthesis	203
1. Heme	203
2. FeS	204
a. Discovery and Function of Prototypical Bacterial, Nif, Isc and Other Components	204
b. Eukaryotic Homologs of Nif and Isc Components Function in Mitochondria	205
c. Plastid Components that are Nif/Isc-Related	205
d. The More Recently Discovered Suf System Functions in FeS Cluster Assembly in Bacteria and Plastids	206
e. Directly Discovered Plastid Components	206
3. Hydrogenase Fe Cluster	207
C. Transport of Iron into Chloroplasts	207
D. Fe-Deficiency Impacts the Photosynthetic Apparatus	208
E. Fe-Sensing	209
III. Cu	209
A. Copper-Protein Assembly	209
1. Transporters	209
2. Chaperones	210
B. Cu Deficiency and Regulation by Cu	210
1. <i>Chlamydomonas</i>	210
2. <i>Arabidopsis</i>	211
3. Redistribution of Copper	211
IV. Mn	211
A. Manganese Transport	211
B. Manganese Deficiency	212
V. Questions for Future Investigation	212
Acknowledgments	213
References	213

*Author for correspondence, email: merchant@chem.ucla.edu

Summary

Redox reactions, which are central to metabolism, depend on redox active functional groups on enzymes such as cysteinyl thiols, organic cofactors like pyridine or flavin nucleotides, or metal cofactors such as Fe, Cu, Mn, or Mo. Accordingly, certain metals are nutritionally essential for life. The green organs of plants display the most descriptive symptoms of metal-deficiency because of the importance of the chloroplast in various metal-dependent redox pathways. The impact of trace element deficiency on chloroplast function at the molecular level has been studied most extensively in microorganisms such as the alga *Chlamydomonas*, for whom the growth media are more readily manipulated. Studies of trace metal distribution and its regulation in cyanobacteria are also relevant to our understanding of chloroplast processes. Fe is the most limiting metal nutrient for all forms of life. In chloroplasts, Fe is found as a redox-active cofactor in FeS centers, heme, mononuclear and di-iron enzymes, and also in ferritin, which functions as an iron “store” and iron “buffer” to maintain intracellular iron homeostasis. The plastid is the key organelle for heme biosynthesis, but FeS cluster synthesis occurs in both plastids and mitochondria. The machinery for cluster synthesis is derived from bacteria, with the process in mitochondria derived from the Isc system and that in plastids containing components of both the Suf system and the Isc system. In iron-deficient chloroplasts, the abundance of iron-containing proteins and specific chlorophyll-proteins is reduced by hierarchical post-transcriptional regulatory mechanisms that may receive signals from iron-dependent enzymes in the tetrapyrrole biosynthetic pathway. The abundant copper enzymes in chloroplasts include plastocyanin and, in some plants, polyphenol oxidase in the thylakoid lumen, and CuZn-superoxide dismutase in the stroma of plants but not green algae. Distributive copper transporters and chaperones are responsible for delivery of the metal to specific sub-organellar compartments. Again, a hierarchical pattern of copper allocation is noted, with plastocyanin receiving copper with higher priority in *Arabidopsis* where plastocyanin is essential, but not in *Chlamydomonas* where a heme protein can substitute for plastocyanin function. A master regulator of copper nutrition called Crr1 regulates degradation of apoplastocyanin in *Chlamydomonas*. The mechanisms of manganese delivery and distribution have not been studied in eukaryotic photosynthetic organisms but, by analogy to metal uptake pathways required for loading Mn-enzymes in bacteria and mitochondria, could involve MntA and MntH/Nramp-like transporters. Mn-deficiency impacts the water oxidation machinery in the chloroplast and also mitochondrial superoxide dismutase.

I. Introduction

A. The Transition Metals Function as Redox Catalysts

Redox reactions are central to metabolism. Biosynthetic reactions involve the reduction of inorganic compounds—CO₂, nitrate, N₂ and sulfate—to more reduced compounds such as carbohydrates, fatty acids and functional groups such as amines, alcohols and thiols. Energy producing reactions involve the oxidation of reduced organic compounds at the expense of a terminal electron acceptor, which is oxygen in most aerobic organisms. The relevant pathways are replete with enzymes that carry redox active cofactors, commonly pyridine and flavin nucleotides or various metal centers. Several transition metals are useful biological catalysts because ionic species of different oxidation states form

stable chelates with functional groups found in proteins (Merchant and Dreyfuss, 1998). The use of particular metals in biology does not reflect their abundance in the earth's crust. Those elements that are bioavailable at neutral pH, either because of the high solubility of the aqua complexes of the low oxidation states (e.g. Cu²⁺ and Fe²⁺) or of the corresponding hydrated oxyanions for the higher oxidation states (e.g. molybdate and vanadate), are used preferentially (Kaim and Schwederski, 1994; Raven *et al.*, 1999).

The use of these metals for catalysis in particular pathways makes organisms that use those pathways dependent on the availability of such metals, leading to the concept of essential or beneficial nutrients (Frieden, 1985; Marschner, 1995). Accordingly, organisms have evolved mechanisms for assimilating the essential metals from the environment, often accumulating them to high levels against a concentration gradient. Because the metal ions are reactive, the assimilation pathways are regulated by supply and demand, and when supply does not meet demand, adaptive mechanisms for

Abbreviations: ATPase – adenosine 5'-triphosphatase; EDTA – ethylenediaminetetraacetic acid; PSI – Photosystem I; PSII – Photosystem II; SOD – superoxide dismutase.

conserving, re-distributing and prioritizing the metal nutrient come into play. Based on its amount in an organism, the metal is classified as a micronutrient (e.g. Fe), a trace nutrient (e.g. Cu) or an ultra-trace nutrient (e.g. Se, Mo or Mn). The amount required is organism-specific. Plants carry out different metabolism than do animals, and therefore their micronutrient requirements are distinct.

B. Trace Metal Deficiency Impacts the Chloroplast

The impact of metal nutrition on the chloroplast has long been recognized because the symptoms of metal nutrient deficiency, which invariably include some form of “chlorosis” or chlorophyll-deficiency, are easily visualized in the green organs. There is a substantial amount of descriptive older literature on Fe-, Cu- and Mn-deficient plants, which established the importance of these metals in various metabolic pathways (Marschner, 1995). More recent research has focused on understanding the cell biology of metal homeostasis, and for this purpose, microorganisms such as algae and cyanobacteria have been useful because of the facility with which the growth media can be metal-depleted or fortified, coupled with the possibility of monitoring a large homogeneous population of cells.

C. Metalloprotein Assembly—Thermodynamics vs. Kinetics

It is useful to emphasize that the fundamental biochemical principles that are taught in the context of the so-called “central metabolic pathways” apply also to the metabolism of the metal nutrients. For instance, a requirement for catalysis of metalloprotein assembly was not appreciated historically because metal-binding to apoproteins was known to be a thermodynamically favorable reaction, based on the fact that metalloproteins are usually more stable than their corresponding apoproteins. Also, in the case of FeS centers, the uncatalyzed reaction occurred readily and produced the correct cluster as long as the appropriate reducing conditions were provided (Malkin and Rabinowitz, 1966). Nevertheless, metalloprotein assembly has been documented for many proteins to be selective *in vivo* relative to the uncatalyzed *in vitro* metal reconstitution reactions. Not surprisingly, the rate of assembly *in vivo* is much faster than for the corresponding uncatalyzed *in vitro* reaction. The use of assembly factors *in vivo*, aside from accelerating a specific reaction, also provides a

means for regulation of metal cofactor utilization. This is relevant in a cell where a limiting micronutrient like iron may be required both for respiratory chain function as well as for photosynthesis. In this situation, iron delivery pathways to individual organelles allow assembly pathways to respond to metabolic demand. In a multicellular organism, there may also be differentiation of function and hence expression of particular metal-utilizing metabolic pathways in specialized organs or at particular developmental states, which suggests operation of inter-cellular signals for metal nutrient homeostasis. This area of metabolism is presently under-studied, especially in the context of chloroplast function in plants. The interested reader is referred to the work of Raven for an excellent treatment of variation in metal requirement in photosynthetic organisms in response to metabolic demand (Raven, 1988, 1990; Raven *et al.*, 1999).

Another point to consider is the use of equilibrium constants in describing intracellular metal distribution. This treatment is valid only in a system that is at equilibrium, which is distinctly not the case in a living cell. The assimilation and distribution of metal cofactors in most organisms requires an input of energy to maintain the system at a steady state away from equilibrium. This is grasped easily for metal transport (e.g. by metal-transporting P-type ATPases) but it applies also to other steps in metalloprotein assembly, including preparation of the apoprotein substrates (e.g. maintenance of ligand oxidation state) and formation of clusters (e.g. the Mn₄Ca complex involved in oxygen evolution). Theoretical calculations of concentrations of “free” metals or particular metal-ligand complexes based on equilibrium constants must be interpreted with caution because they may not reflect the true dynamic in a cell, which is in a non-equilibrium situation.

D. Fe, Cu and Mn

The metals that are well-studied in the context of chloroplast biogenesis and function are Fe, Cu and Mn because of their abundance in the photosynthetic apparatus (Raven *et al.*, 1999). Also, the corresponding metalloproteins are readily monitored in the holoform by spectroscopic methods, and changes in metalloprotein expression are therefore easy to visualize. Other important metals such as Zn are spectroscopically silent, and hence, much less is known about the biogenesis of Zn-enzymes in the chloroplast despite their prevalence and abundance. The discussion in this chapter is restricted to the impact of Fe, Cu and Mn nutrition

on chloroplast function, especially the photosynthetic apparatus.

II. Fe

A variety of iron enzymes occur in chloroplasts: heme proteins like cytochromes and P450s, soluble and membrane-bound two- and four-iron-sulfur proteins, di-iron enzymes and mononuclear-iron enzymes, and iron bound to ferritin (Jäger-Vottero *et al.*, 1997; Kerfeld and Krogmann, 1998; Merchant and Dreyfuss, 1998; Briat *et al.*, 1999; Froehlich *et al.*, 2001; Berthold and Stenmark, 2003; Gray *et al.*, 2004; Tian and DellaPenna, 2004). These enzymes function in electron transfer reactions of the photosynthetic apparatus and in various redox reactions in pathways that produce secondary metabolites and natural products. Because iron is an actively acquired nutrient that limits most life forms, organisms generally do not excrete iron when it is in intracellular excess beyond what is needed for maintenance of the iron enzymes, but rather store it. The reactivity of iron in an aerobic environment requires that it be stored in a protected form, as in the protein ferritin. When iron is required for *de novo* synthesis of iron-containing proteins, it can be mobilized from the stored form, and when it is released as iron-containing enzymes are degraded, the store can be re-built. Ferritin is therefore a key component of iron homeostasis.

A. Ferritin

The protein ferritin accounts for stored iron in a plant and is the source of iron for chloroplast development. The reader is referred to substantial reviews by Briat and Theil and their co-workers for details of ferritin chemistry and biology (Briat and Lobréaux, 1997; Briat *et al.*, 1999; Curie and Briat, 2003; Theil, 2003, 2004). The protein consists of 24 subunits that bind up to 4.5×10^3 atoms of iron as ferric-oxy-hydroxide within an internal core. This mineral core is built by movement of iron into the core and oxidation of ferrous to ferric ions by the ferroxidase activity of the ferritin subunits (Lawson *et al.*, 1989). Plant ferritins are distinguished from animal ferritins by being localized within an organelle, the plastid, and by their subunit composition. While animal ferritins consist of two types of related chains, H and L, plant ferritins have a single type of chain. The plant ferritin subunit, encoded by a multi-gene family, is more closely related to the H-chain but also has features of the L chains that facilitate iron

nucleation and yield a stable core (Van Wuytswinkel *et al.*, 1995; Wardrop *et al.*, 1999). The mineral core in plant ferritin is also unique in containing a high proportion of phosphate like bacterioferritins (Waldo *et al.*, 1995). The proteins encoded by the *FER* genes in plants and algae include an N-terminal plastid-targeting sequence in the precursor, and an N-terminal region in the mature protein that is a determinant of stability and protease susceptibility *in vitro* (Ragland *et al.*, 1990; van Wuytswinkel *et al.*, 1995; Wardrop *et al.*, 1999; La Fontaine *et al.*, 2002). Interestingly, plastid ferritin is quite distinct from bacterioferritin found in cyanobacteria, which indicates that chloroplast ferritin is a function acquired from the host rather than retained from the endosymbiont (Laulhere *et al.*, 1992).

Ferritin expression is determined by multiple signals because of the nutritional importance of iron, the variation in iron demand at different stages of growth, and the potential for toxicity of iron in an aerobic environment. Ferritin abundance is regulated by changes in RNA abundance through transcriptional regulation and polypeptide abundance through post-transcriptional mechanisms in response to multiple signals, including iron supply, developmental stage, and wounding (Lescure *et al.*, 1991; Lobréaux and Briat, 1991; Ragland and Theil, 1993; Fobis-Loisy *et al.*, 1996; Tarantino *et al.*, 2003). Therefore, it is important to monitor protein abundance for a picture of ferritin action *in vivo*, while RNA abundance only presents a picture of the potential or capacity for ferritin action. Recent studies in animals and plants suggest that ferritin can also be found in mitochondria under certain conditions (Levi *et al.*, 2001; Zancani *et al.*, 2004), which adds another layer of complexity in understanding the biology of ferritin. Another relevant aspect of ferritin function is the amount of iron in the mineral core, which can change during growth and development (e.g. van der Mark *et al.*, 1981), but this has not been studied systematically.

The four ferritin-encoding genes in *Arabidopsis*, *FER1*, *FER2*, *FER3* and *FER4*, show unique patterns of expression in response to iron nutrition, environmental stress and developmental stage, which reinforces the importance of plastid Fe homeostasis for plant growth and physiology (Petit *et al.*, 2001; Tarantino *et al.*, 2003). Ferritin accumulates at high levels in non-green plastids and decreases in abundance as they green, suggesting that ferritin is the source of iron found in heme- and other iron-containing proteins in the photosynthetic apparatus, although the direct movement of labeled Fe from ferritin to an Fe-containing enzyme has not been monitored. As the leaf gets older, the ferritin

content decreases until the organ is at the stage of senescence, when the ferritin content increases again, presumably to accommodate iron that is released from enzymes as the proteins of the chloroplast are degraded. The re-appearance of ferritin is attributed to increased gene expression and de novo synthesis (Tarantino *et al.*, 2003). Ferritin accumulation is also increased under conditions of oxidative stress, which presumably exacerbate the toxicity of iron (Briat *et al.*, 1999; Petit *et al.*, 2001).

Although ferritin accumulation involves post-transcriptional regulatory mechanisms, when a soybean *FER* cDNA was over-expressed in tobacco plants via a 35S promoter-driven construct, ferritin did over-accumulate in the mature plants (Van Wuytswinkel *et al.*, 1998). The plants accumulated more iron but displayed symptoms of iron-deficiency (i.e. inter-veinal chlorosis) even when the ferritin was targeted to the plastid. These studies emphasize the role of plastid ferritin in iron homeostasis and the function of ferritin as an iron “buffer” that keeps iron available in the cell but in a non-toxic form. This work also raised the question of how iron might be mobilized from ferritin. The more abundant ferritin in the over-expressing plants catalyzes and accommodates the over-chelation, but the fact that it is not available for normal chloroplast development indicates that the iron mobilization system can not circumvent the over-chelation. Because oxidation is involved in the deposition of iron in the ferritin core, it is generally assumed that reduction is required for iron mobilization and there are some experiments that support this notion (e.g. Bienfait and van den Briel, 1980). In that *in vitro* study, a connection between copper and ascorbate- and oxygen-dependent iron mobilization from ferritin was noted, suggesting perhaps a role for an enzyme like ascorbate oxidase. Nevertheless, an *in vivo* connection has not yet been established. The models considered for iron release from ferritin in animal cells favor either lysosomal degradation of the protein and/or unfolding of the iron cores, but mobilization of iron from the mineral would still depend on reduction (reviewed by Theil, 2004). Besides serving as a buffer for iron, ferritin is also an important store of Fe for the next generation and does accumulate in seeds (Masuda *et al.*, 2001).

B. Heme, FeS and Fe Cofactor Synthesis

During development of the chloroplast in germinating seedlings, the disappearance of ferritin is correlated with the appearance of iron-containing catalyzes such as

cytochromes that contain heme (or Fe-protoporphyrin IX) and iron-sulfur proteins (of either the Fe₂S₂ variety as in ferredoxin or the Fe₄S₄ variety as in PSI). It is possible that iron distribution to various enzymes is regulated based on physiological demand for, or importance of, particular metabolic pathways (see below). For an understanding of the principles underlying the regulation, it is first useful to have a description of the iron-utilizing pathways.

The biosynthesis of heme and FeS centers represent major iron utilizing pathways in the cell, and the organelles (mitochondria and plastids) contain an abundance of these redox cofactors. In *Saccharomyces cerevisiae*, inhibition of mitochondrial heme synthesis blocks transcriptional activation of the iron uptake genes, and disruption of iron-sulfur metabolism results in mitochondrial iron accumulation and hence altered cellular iron homeostasis, substantiating the importance of cofactor biosynthesis in iron homeostasis (e.g. Knight *et al.*, 1998; J Li *et al.*, 1999; Lange *et al.*, 2000; Crisp *et al.*, 2003).

1. Heme

In fungi and animal cells, the heme biosynthetic pathway is distributed between the cytosol and the mitochondrion. In plants, the entire pathway, from δ -aminolevulinate to Fe-protoporphyrin IX, is localized in plastids, but the last two enzymes—protoporphyrinogen oxidase and ferrochelatase—occur also in the mitochondrion (Chow *et al.*, 1997; Lermontova *et al.*, 1997; Beale, 1999; Watanabe *et al.*, 2001). In *Arabidopsis*, two ferrochelatase-encoding genes, *FC-I* and *FC-II*, are expressed under different conditions (Singh *et al.*, 2002). Interestingly, *FC-I*, whose product is probably localized to both plastids and mitochondria, showed increased expression in leaves upon wounding or upon treatment with salicylic acid, which suggested that an increased potential for heme synthesis, presumably to provide cofactors for induced P450s (see below), is part of the defense response. It is not presently known whether heme found outside the plastid, in peroxisomes (peroxidases), cytosol (hemoglobin), endoplasmic reticulum (P450s and non-heme oxygenases), or the cell wall (peroxidases), is derived from the plastid pool or the mitochondrial pool.

Iron deficiency impacts heme levels and reduces the content of cytochromes in the photosynthetic apparatus (Duggan and Gassman, 1974; Moseley *et al.*, 2002a). Because the tetrapyrrole pathway is feedback regulated by heme, synthesis of δ -aminolevulinate is promoted

under these conditions (Cornah *et al.*, 2003; Franklin *et al.*, 2003).

For many organisms, heme is a source of iron, especially under conditions of nutritional deficiency. Heme or Fe-protoporphyrin IX is oxidized by a mixed function-type oxidase reaction, which requires molecular oxygen and a reductant, to Fe-biliverdin (usually the α isoform) and CO. This reaction, catalyzed by a heme oxygenase, is driven by removal of product through the action of biliverdin IX α reductase (Franklin *et al.*, 2003). The step also releases iron bound to the tetrapyrrole. In animals, fungi and bacteria, a heme oxygenase is a key target of iron deficiency because it releases iron from heme for use in other iron-containing enzymes (e.g. Poss and Tonegawa, 1997; Protchenko and Philpott, 2003; Frankenberg-Dinkel, 2004; Skaar *et al.*, 2004).

The role of ferrochelatase as an iron-utilizing enzyme and heme oxygenase as an iron-releasing enzyme in plastid iron homeostasis is intriguing but not yet analyzed thoroughly in plants. The major role of the plastid heme oxygenases lies in the production of bilins for light harvesting and signaling (Willows *et al.*, 2000; M Terry *et al.*, 2002). In a red alga, the gene *pbsA* in the plastid genome, encoding a heme oxygenase, is transcriptionally activated by iron-deficiency (Richaud and Zabulon, 1997). In *Chlamydomonas* neither of the two typical heme oxygenase-encoding genes appears to be transcriptionally regulated by iron nutrition (J. Kropat and S. Merchant, unpublished results), and the question of regulation by iron has not yet been addressed for the various heme oxygenases of plants. Nor has the issue of whether plastid heme levels may signal organelle or cellular iron status, as evidently is the case for mitochondrial heme in yeast, been addressed.

2. FeS

a. Discovery and Function of Prototypical Bacterial Nif, Isc and Other Components

The biosynthesis of FeS centers requires mobilization of sulfur from cysteine, mobilization of Fe, assembly of the cluster and transfer of the cluster to apoenzymes (Merchant and Dreyfuss, 1998; Lill and Kispal, 2000; Frazzon and Dean, 2003). Although not all of the molecular events are understood, the necessary components have been defined in many organisms through classical and reverse genetic approaches. The first components were identified in the context

of nitrogenase function (called NifS and NifU) and, subsequently, related molecules encoded by *isc* genes in *Escherichia coli* were shown to be responsible for the assembly of clusters in various iron-sulfur proteins (Zheng *et al.*, 1993; Takahashi and Nakamura, 1999). Sulfur is mobilized from cysteine by NifS/IscS in a pyridoxal phosphate-dependent reaction catalyzed by a cysteine desulfurase, which yields alanine and sulfane sulfur. This activity is required also for the synthesis of other sulfur containing compounds in bacteria such as thiamine and thionucleosides. NifU, a three-domain protein, is responsible for building the FeS cluster (Yuvaniyama *et al.*, 2000). An N-terminal iron-binding domain related to IscU is the assembly scaffold and binds a “transient” cluster through three cysteine residues (Agar *et al.*, 2000). IscU interacts with the chaperone system, Hsc66 and Hsc20, encoded in the *isc* operon (Table 1) (Hoff *et al.*, 2000). The middle domain of NifU holds a permanent FeS cluster, presumably with redox function during cluster assembly. The C-terminal domain, called the NFU domain or CnfU, is involved in transferring the FeS cluster to apoproteins. The domain contains a CxxC motif of unknown function (Frazzon *et al.*, 2002). IscA is another cluster assembly component providing a scaffold for transient formation and binding of an FeS cluster, but how it relates to IscU function is not understood (Ollagnier-de-Choudens *et al.*, 2001; Cupp-Vickery *et al.*, 2004). A ferredoxin (itself containing an FeS cluster) is associated with the *isc* operon and is required for cluster biogenesis, presumably for reducing ferric to ferrous iron. This ferredoxin interacts physically with IscA (Ollagnier-de-Choudens *et al.*, 2001).

Interestingly, under anaerobic but not aerobic conditions, a simple system consisting solely of NifS- and NifU-homologues is sufficient for FeS cluster assembly in *E. coli* lacking both the *isc* and *suf* (see below) operons (Ali *et al.*, 2004). This suggests that the additional components of the Isc system evolved to accommodate an aerobic environment.

In a genetic screen for defects in FeS cluster metabolism in *Salmonella enterica*, two new assembly factors were identified, called ApbC and ApbE (Skovran and Downs, 2003). Phenotypic analysis suggests that these proteins are required for the maintenance (repair) or assembly of oxygen-labile FeS clusters in the enzymes ThiH (required for thiamine biosynthesis), succinate dehydrogenase (containing multiple FeS clusters) and aconitase. The specific role of the proteins is not known, but ApbC does catalyze ATP hydrolysis and the *apb* mutants (unlike *isc*

Table 1. Biochemical functions required for cluster biosynthesis in bacteria and organelles. A list of components and types of activities required for FeS cluster assembly in various organisms. See text for details.

Function	Gene
Sulfur mobilization from cysteine	bacterial IscS, NifS mitochondrial Nfs1
ATP dependent sulfur transfer?	bacterial and plastid SufS ³ +SufE
ATP-dependent Repair / synthesis of (oxygen labile Fe ₄ S ₄) clusters	bacterial and plastid SufC+SufB+SufD bacterial ApbC / Mrp cytosolic Cfd1p / Nbp35p plastid Hcf101
Cluster assembly and provision of scaffold for assembly	bacterial NifU-N terminus, IscU mitochondrial Isu1, Isu2
Cluster assembly	bacterial IscA mitochondrial Isa1, Isa2 plastid ISAI ³
Cluster transfer	bacterial and plastid SufA ⁴ bacterial NifU-C terminus with conserved CxxC motif mitochondrial Nfu1 plastid NFUs
Reductant	bacterial NifU-permanent cluster in central portion bacterial Fdx mitochondrial Yah1
Chaperones	bacterial ATP-dependent Hsp70, HscA, Hsc66 + bacterial J-type co-chaperone, HscB, Hsc20 mitochondrial Ssq1 + Jac1
Iron metabolism	mitochondrial frataxin

³ Also called CsdB in *E. coli* or NifS-Type II.

⁴ SufA is related to IscA and the nomenclature used depends on genic context (i.e. whether the gene occurs in a *suf* vs. *isc* operon).

mutants) can be suppressed by exogenous iron. These features are reminiscent of the essential P-loop ATPase Cfd1p in *S. cerevisiae*, which is involved in repair or synthesis of the aconitase cluster in the cytosol, and indeed, Cfd1p and ApbC share sequence similarity (Roy *et al.*, 2003). Cfd1p contains a conserved CxxCxxC motif, of which the first two cysteine residues are functionally important and conserved also in ApbC.

b. Eukaryotic Homologs of Nif and Isc Components Function in Mitochondria

The discovery of homologs in eukaryotes of the proteins encoded in the bacterial *nif/isc* operon led to the description of a related machinery in mitochondria, consisting of Nfs1 (related to NifS), Nfu1 (related to the C-terminal domain of NifU), Isu1/2 (related to the N-terminal domain of NifU), Isa1/2 (related to IscA), Yah1 (a mitochondrial ferredoxin) and the molecular chaperone system, Ssq1 plus Jac1 (related to Hsc66 and Hsc20) (Table 1) (Garland *et al.*, 1999; Kispal *et al.*, 1999; J Li *et al.*, 1999; Schilke *et al.*, 1999; L Jensen and Culotta, 2000; Lange *et al.*, 2000; Mühlhoff *et al.*, 2002). In *S. cerevisiae*, and probably animals as well, this machinery appears to be the source of

clusters for all FeS proteins in the cell (Lill and Kispal, 2000), although additional components, like Cfd1p, Nar1p and Nbp35p, are required for the synthesis of extra-mitochondrial clusters (Roy *et al.*, 2003; Balk *et al.*, 2004).

Nevertheless, several lines of evidence suggested that plastids make their own FeS clusters. First, isolated plastids could incorporate sulfur from cysteine into acid-labile clusters in ferredoxin in a reaction requiring ATP and NADPH (Takahashi *et al.*, 1986; Takahashi *et al.*, 1991a; Takahashi *et al.*, 1991b). Second, newly imported apo-ferredoxin could be converted into the holoform *in vitro* in isolated plastids in the absence of cytosol (HM Li *et al.*, 1990; Pilon *et al.*, 1992). And third, most of the iron in the plant cell is plastid-localized (see above). Therefore, it was concluded that plant cells must have at least two FeS assembly machineries, one in the mitochondrion and one in the plastid.

c. Plastid Components That are Nif/Isc-Related

Analysis of the *Arabidopsis* genome revealed two *nifS*-related genes. One, *AtNFS1*, is suggested to

encode a mitochondrially-targeted protein, and another, *AtNFS2*, a plastid-localized one with cysteine desulfurase activity (Kushnir *et al.*, 2001; Léon *et al.*, 2002). Subsequently, five NFU proteins, each containing the conserved C-terminal CxxC motif, and encoded by *AtNFU1* through *NFU5* were described, as well as a plastid-localized *IscA* homolog, *AtISA1* (Léon *et al.*, 2003; Yabe *et al.*, 2004). The NFU proteins were distinguished into two sub-types, the NFU1-3 type¹ being plant-specific and proposed to be plastid-localized based on immunodetection, GFP fusions and *in vitro* import studies, and NFU4 and NFU5 being mitochondrial. A T-DNA insert in the *NFU* gene on chromosome V, *CNFU2* or *NFU2* (encoding a plastid-type protein), resulted in decreased abundance of photosystem I, ferredoxin, sulfite reductase, and reduced stromal iron-sulfur cluster assembly activity (Touraine *et al.*, 2004; Yabe *et al.*, 2004). On the other hand, Fe_3S_4 glutamate synthase activity was not reduced, nor was the abundance of the Fe_2S_2 Rieske protein or subunits of the cytochrome *b₆f* complex reduced, suggesting distinct pathways for plastid iron-sulfur cluster assembly (Touraine *et al.*, 2004).

d. The More Recently Discovered Suf System Functions in FeS Cluster Assembly in Bacteria and Plastids

The Suf pathway was revealed through analysis of suppressors of *E. coli* strains in which the ISC machinery was deleted (Takahashi and Tokumoto, 2002). In the suppressed strain, the Suf pathway, which is important in bacteria under conditions of oxidative stress and iron limitation (Nachin *et al.*, 2003; Outten *et al.*, 2004; Wang *et al.*, 2004), is mis-expressed, allowing the SUF system to cover the loss of the ISC system. Homologs of the SUF components (called SufABCDEs) are found in many bacteria, archaea and plastid-containing eukaryotes (Ellis *et al.*, 2001), and the operation of the Suf pathway in *Arabidopsis* plastids was demonstrated recently (Xu and Møller, 2004). As mentioned above, anaerobic conditions facilitate cluster assembly. The corollary is that conditions that favour oxidation reactions make cluster maintenance and assembly more difficult. The use of the Suf system in cyanobacteria and chloroplasts perhaps represents an adaptation to

greater oxidative stress in this compartment relative to the mitochondrion.

SufC, a cytoplasmic ABC-type ATPase in bacteria, is a key component of the SUF system because of its high degree of conservation and the severity of phenotype associated with loss of function (Nachin *et al.*, 2003). SufC associates with SufB and SufD to form a complex that is required for “repair” of labile FeS clusters that are damaged during oxidative stress, which may mean energy-dependent insertion of Fe^{2+} into Fe_3S_4 centers.

SufS is related to NifS and in bacteria constitutes one subunit of a cysteine desulfurase. SufS also exhibits selenocysteine lyase activity, which is required for the synthesis of selenoproteins. The second subunit, SufE, enhances the cysteine desulfurase over the selenocysteine lyase activity (Loiseau *et al.*, 2003). The plastid SufS, called CpNifS or AtNFS2, also shows both cysteine desulfurase and selenocysteine lyase activity (Pilon-Smits *et al.*, 2002). Interestingly, the recombinant protein shows only a fraction (1 to 2%) of the activity of the endogenous protein in stromal extracts in the assembly of iron sulfur clusters of ferredoxin (Ye *et al.*, 2004). One possibility is that only a fraction of the recombinant protein is active. Another possibility, raised by the function of SufE in bacteria, is that the *in vivo* reaction involves other factors, such as a plastid SufE-homolog (Xu and Møller, 2004). The latter model is supported by the observation of a high molecular weight CpNifS-containing complex (Ye *et al.*, 2004). SufS and NifS are related, but the key difference may be the dual role of SufS in both Se and S metabolism, dependent on interaction with SufE. SufA is related to *IscA* and by analogy probably has a role in cluster assembly.

e. Directly Discovered Plastid Components

In a screen for mutants of *Arabidopsis* defective in the assembly of PSI, Meurer and co-workers identified HCF101 as a candidate FeS cluster assembly factor in the plastid stroma (Lezhneva *et al.*, 2004; Stöckel and Oelmüller, 2004). They proposed a role for HCF101 in the assembly of Fe_4S_4 clusters as opposed to Fe_2S_2 clusters based on a drastic decrease in the abundance of PSI reaction center polypeptides PsaA and PsaB, attributable to degradation of the apoproteins as a result of a post-translational block in assembly. A less dramatic increase in the peripheral subunits argued for an effect of HCF101 on cofactor biogenesis and, more specifically Fe_4S_4 centers, owing to a 50% decreased abundance of ferredoxin thioredoxin reductase but not of ferredoxin or the Rieske FeS protein.

¹ The nomenclature of Leon *et al.* (2003) is used here owing to precedence. The gene names for the mitochondrial forms are *atNFU1* and *atNFU2* and for the plastid forms *atCNFU1* through *atCNFU3* in the work of Yabe *et al.* (2004).

The sequence relationship between HCF101 and ApbC (Lezhneva *et al.*, 2004), and also between HCF101 and Cfd1p, solidifies its role in FeS biogenesis in the plastid, as does its iron-dependent expression (Stöckel and Oelmüller, 2004), but what specifically that role might be is unclear. HCF101 also contains a conserved CxxC motif (Lezhneva *et al.*, 2004). As appropriate for a gene encoding a PSI assembly factor, *HCF101* is expressed in green organs (Stöckel and Oelmüller, 2004). The *Arabidopsis* genome encodes three different HCF101-like proteins—HCF101, HCF101-L1 and HCF101-L2, but L1 and L2 have not been characterized functionally as yet.

HCF101 is highly conserved with homologs in all kingdoms of life. These have been classified into four groups (Lezhneva *et al.*, 2004). HCF101 itself belongs to Class 1. The Class 2 form (L1 in *Arabidopsis*) is proposed to function in mitochondria based on the presence of an apparent N-terminal pre-sequence. The Class 3 form (L2 in *Arabidopsis*) functions in the cytosol and nucleus based on the location of the yeast representative, Nbp35p (Hausmann *et al.*, 2005), and the Class 4 form is in the cytosol. It is possible also that one of the *Arabidopsis* homologs is a redundant factor in the plastid (accounting perhaps for the weak impact of loss of HCF101 on ferredoxin thioredoxin reductase).

It is likely that a continued classical genetic approach to the study of plastid FeS cluster biogenesis could reveal new components besides those related to the products of the well-studied *isc* operon and the more recently discovered *suf* operon. A recent publication suggests that *Arabidopsis* APO1 may be involved in Fe₄S₄ cluster biosynthesis (Amann *et al.*, 2004), but the pleiotropic impact of loss of APO1 on membrane structure and the absence of homologs in *Chlamydomonas* and cyanobacteria suggest that APO1 may be more generally involved in a development-specific aspect of thylakoid biogenesis.

3. Hydrogenase Fe Cluster

Besides heme and FeS centers, there are other iron-containing cofactors in enzymes, including mono- and di-iron enzymes as well as uncharacterized iron sites (Fox, 1998; Plank *et al.*, 2001; Berthold and Stenmark, 2003; Hausinger, 2004). One of these is the bi-nuclear Fe center of hydrogenase in *Chlamydomonas* (Happe *et al.*, 1994). Genetic analysis of hydrogenase-minus mutants revealed two new enzymes, HydE and HydG belonging to the “Radical SAM” family (Sofia *et al.*, 2001), which are required for production of active

hydrogenase (Posewitz *et al.*, 2004). The restricted occurrence of HydE and HydG homologs in Fe-hydrogenase containing prokaryotes is consistent with the proposed function in hydrogenase assembly. By analogy to the biosynthesis of the nitrogenase cluster, the authors of this work suggest that HydE and HydG may be involved in mobilization of iron for assembly of the so-called H-cluster of the [Fe] hydrogenase, which also requires CN, CO and the di(thio-methyl)amine ligand, but they do not rule out a function in in situ generation of the iron-coordinating ligands. Heterologous expression of hydrogenase in *E. coli* requires only the HydE and HydG factors in addition to the gene for the apoprotein, indicating that these are probably the most critical assembly factors.

C. Transport of Iron into Chloroplasts

If ferritin is assembled with iron in the plastid, then there must be a mechanism for iron transport into the chloroplast. Also, under conditions of iron deficiency, it may be necessary to re-allocate iron from one compartment to another (such as the mitochondrion where iron is required for the function of respiratory enzymes), and intracellular transporters are expected to be central to organelle metal homeostasis. However, such molecules have not yet been discovered, although the Nramp transporters are candidates. The Nramp proteins, originally identified in mammals as iron transporters that affect resistance to microbial infection, are considered to be broad specificity, divalent cation transporters (Gunshin *et al.*, 1997). They are encoded in plant and other eukaryotic genomes as multi-gene families with members displaying distinct patterns of expression in response to divalent cation nutrition, suggesting that they may have metal-specific roles *in vivo*. While they are generally considered to be assimilatory transporters, they are found also in intracellular membranes in *S. cerevisiae* (reviewed by Van Ho *et al.*, 2002), and it is possible that they function to deliver metal ions into and out of organelles in plants. For instance, NRAMP3 in *Arabidopsis* localizes to the vacuolar membrane (Thomine *et al.*, 2003).

In *S. cerevisiae*, members of the carrier family—Mrs3p and Mrs4p—appear to be involved in iron transport across the mitochondrial inner membrane (Foury and Roganti, 2002; Mühlhoff *et al.*, 2003; Kunji, 2004; Lesuisse *et al.*, 2004). Although the *Arabidopsis* genome encodes 58 members of the mitochondrial carrier family, the probability that Mrs3p and Mrs4p homologues would function in chloroplasts is low because the plastid inner envelope transporters tend to be

distinct from the mitochondrial carriers (Flügge *et al.*, 2003; Picault *et al.*, 2004).

D. Fe-Deficiency Impacts the Photosynthetic Apparatus

The abundance of the photosynthetic apparatus in chloroplasts and the numerous iron-containing redox-active proteins therein, plus the loss of chlorophyll proteins as a marker for iron-deficiency, meant that studies of the impact of iron-deficiency on plastid biochemistry have focused on photosynthesis (e.g. N Terry, 1983; N Terry and Abadía, 1986). Nevertheless, the discovery of di-iron enzymes in desaturation and other fatty acid modification reactions, and in carotenoid synthesis (Cunningham and Gantt, 1998; Shanklin and Cahoon, 1998), and the recent molecular identification of cytochrome P450 enzymes functioning in the biosynthesis of carotenoids, oxylipins and other isoprenoid-derived compounds, indicates the importance of iron in many other physiological processes including the production of defense metabolites (Froehlich *et al.*, 2001; Helliwell *et al.*, 2001; Tian and DellaPenna, 2004).

General principles concerning the impact of iron-deficiency on photosynthesis in plants have not yet emerged because it is difficult to directly compare individual studies with different plant material at different stages of growth and under various conditions of other nutrients. Therefore, our understanding of iron-deficiency adaptation of the photosynthetic apparatus comes largely from studies of microorganisms like cyanobacteria and green algae where iron nutrition can be readily and uniformly controlled (Moseley *et al.*, 2002a; Michel and Pistorius, 2004). Some well-documented changes in response to iron-deficiency in cyanobacteria include the replacement of iron-containing ferredoxin by iron-free flavodoxin, a decrease in the ratio of PSI to PSII from about 4:1 to 1:1, and the *de novo* synthesis of a new antenna for photosystem I consisting of the IsiA polypeptide or CP43' (Laudenbach *et al.*, 1988; Laudenbach and Straus, 1988; La Roche *et al.*, 1996; Bibby *et al.*, 2001; Boekema *et al.*, 2001). PSI is also a prime target in iron-deficient plants, presumably because of its high iron content (three Fe₄S₄ clusters), but the other two adaptations are not known to occur in chloroplasts (Nishio *et al.*, 1985). The *Chlamydomonas* genome appears to encode several different chloroplast-targeted ferredoxins. Two of these genes show a reciprocal pattern of expression dependent on iron nutrition, but the physiological function of this pattern is not known (N. Fischer and J.-D. Rochaix, personal communication; A. Terauchi and S. Merchant, unpublished

results). Because ferredoxin is the source of electrons for many biosynthetic pathways in the plastid, it is possible that synthesis of alternate forms of ferredoxin may determine allocation of reducing power under iron-deficient conditions.

In a recent study with *Chlamydomonas*, a distinction was made between iron deficiency vs. iron-limitation (La Fontaine *et al.*, 2002; Moseley *et al.*, 2002a). *Iron-deficient* cells are defined as those that are not chlorotic but where the assimilator iron uptake genes, *FOX1*, *FTR1* and *FEA1*, are fully induced, whereas *iron-limited* cells are defined as symptomatic (i.e. chlorotic) and the rate of cell division is reduced. In *Chlamydomonas*, a progressive modification of the photosynthetic apparatus was observed. In marginally iron-deficient cells, the LHCI antenna was found to be physically and functionally uncoupled from PSI and this was correlated with an altered association of the PSI-K polypeptide (Fig. 1), which functions to facilitate the transfer of excitation energy from the peripheral




Fig. 1. The PSI-LHCI interaction is dependent on Fe nutrition. In Fe-replete cells, LHCI is physically and functionally associated with PSI in order to optimize energy transfer. As the cells anticipate iron-deficiency (i.e. in the situation where iron uptake is enhanced), the photosynthetic apparatus is modified to decrease excitation energy input into the PSI reaction center. This occurs via a loosening of the interaction between LHCI and PSI through a change in the association of PSI-K with PSI, and also by a change in the polypeptide composition of the LHCI complex (indicated by the change in shape of LHCI from rectangular to rhomboidal). The rationale for this modification is to avoid oxidative stress resulting from loss of iron from labile Fe₄S₄ clusters in PSI. When iron is limiting cell division, despite the expression of high affinity transporters, the abundances of PSI and the LHCI antenna are down-regulated by induced proteolysis to conserve iron (indicated with the scissors symbols). The plastid iron status may be sensed by flux through the tetrapyrrole pathway. For regulating chlorophyll protein abundance, the level of chlorophyll synthesized is relevant, and this is sensitive to iron nutrition because the aerobic oxidative cyclase (CHL27) is an iron-containing enzyme. For regulating heme proteins, the level of heme may be relevant, and this is clearly dependent on iron availability at the step catalyzed by ferrochelatase.

antenna to the reaction center (P Jensen *et al.*, 2000). This response was suggested to function as a protective mechanism to avoid photo-oxidative damage resulting from damaged or improperly assembled FeS clusters in PSI. And indeed, iron-deficiency or loss of LHCI antenna can “rescue” the light sensitivity of a *Chlamydomonas* strain lacking PsaF (Hippler *et al.*, 2000; Moseley *et al.*, 2002a).

As iron-deficient *Chlamydomonas* cells progress to iron limitation, specific subunits of the LHCI antenna proteins are degraded, and the abundance of both photosystems and the cytochrome complex is reduced to about 5% of the level maintained in iron-replete cells coincident with a decreased rate of cell division. The abundance of the ATP synthase is unchanged by iron deficiency. Interestingly, the chlorophyll content on a per cell basis is maintained at about 50% of the level found in iron-replete cells, with most of the chlorophyll associated with LHCII complexes that are not associated with a reaction center. This complex may serve as a reservoir of chlorophyll for de novo synthesis of photosystems during the re-greening process initiated by iron nutrition. De novo chlorophyll synthesis does occur even in iron limited cells, but the new pigment must be selectively allocated to the LHCII in -Fe cells.

E. Fe-Sensing

The impact of Fe-deficiency on the biosynthesis of chlorophyll (or chlorophyll proteins) was noted decades ago and is a classic symptom of Fe-deficiency (Bogorad *et al.*, 1958; Machold, 1971; Spiller and Terry, 1980; N Terry, 1980). The aerobic oxidative cyclase or CHL27, now known to be a di-iron enzyme, was suggested to be a key target of iron-deficiency (Spiller *et al.*, 1982; Tottey *et al.*, 2003). The loss of chlorophyll proteins in iron-deficiency was therefore attributed to reduced flux through the chlorophyll biosynthetic pathway (Spiller and Terry, 1980). However, as mentioned above, iron-deficiency chlorosis represents a specific re-programming of chlorophyll protein synthesis rather than a general decrease in all chlorophyll proteins. Because a *Chlamydomonas* mutant with reduced cyclase activity recapitulated parts of the program initiated by marginally iron-deficient cells, it was suggested that the cyclase might represent an iron sensor in the plastid (Moseley *et al.*, 2002a). According to this model, the occupancy of the active site iron in CHL27 would be proportional to iron availability in the plastid. As the cell, and hence the organelle, becomes iron-deficient, the rate of chlorophyll synthesis is decreased, leading to restriction of chlorophyll for the de novo synthesis of chlorophyll proteins. If there is a hier-

archical allocation of chlorophyll to particular apoproteins (and there is considerable evidence for this), then the program of chlorophyll-protein complex accumulation can be linked to plastid iron status. In the case of PSI, it appears that the stability of PsaK is very sensitive to iron-nutrition status as well as chlorophyll biosynthesis. The model therefore suggests that occupancy of the pigment sites in PsaK determines the association of this polypeptide with PSI, and hence the functional association of the LHCI antenna with PSI (Fig. 1). Nevertheless, a causal connection between chlorophyll binding to PsaK and its assembly with PSI has not yet been established. An attractive aspect of this model is that it distinguishes between iron sensing in the plastid vs. iron sensing in the nucleus to control the expression of iron assimilatory genes. On the other hand, the mechanism is clearly relevant only in the context of the photosynthetic apparatus. Whether other plastid types have mechanisms for signaling and responding to iron nutrition is not known. One can envision a similar mechanism for regulation of heme protein accumulation in all plastid types via the action of ferrochelatase, but so far there are no studies that test this idea.

III. Cu

Three abundant plastid proteins that contain copper are CuZnSOD in the plastid stroma and plastocyanin and polyphenol oxidase in the thylakoid lumen (Jackson *et al.*, 1978; Kieselbach *et al.*, 1998). These proteins are not present in all chloroplasts; the chlorophyte algae (such as *Chlamydomonas*) contain only plastocyanin, *Arabidopsis* has CuZnSOD and plastocyanin, and tomato and spinach have all three proteins. The relative proportion of each protein in a given plant cell depends of course on the organ, the developmental state and environmental conditions (e.g. Last and Gray, 1989; Perl-Treves and Galun, 1991; Thygesen *et al.*, 1995; Thipyapong *et al.*, 1997). The use and distribution of copper within plastid compartments and the impact of deficiency on plastid function is therefore likely to vary. Cyanobacteria, like the green algae, do not have CuZnSOD or polyphenol oxidase, but on the other hand, they do have a respiratory copper-containing oxidase.

A. Cu-Protein Assembly

1. Transporters

Once copper is taken up into the plant cell, presumably by a member of the COPT1 family of transporters

(Sancenón *et al.*, 2003; Sancenón *et al.*, 2004), it needs to be distributed to organelles for the biosynthesis of copper-enzymes like cytochrome oxidase, CuZnSOD, and plastocyanin, and to other compartments for the biosynthesis of various intra- or extra-cellular multi-copper oxidases such as ascorbate oxidase and laccase. Some members of the so-called CPx-type heavy-metal transporting ATPases are responsible for intracellular copper distribution (reviewed by Williams *et al.*, 2000). Three of these enzymes have been characterized in *Arabidopsis*. RAN1 is responsible for loading copper in the secretory pathway for the biosynthesis of the ethylene receptor, while PAA1 and PAA2 function to deliver copper to the chloroplast for CuZnSOD and plastocyanin (Hirayama *et al.*, 1999; Shikanai *et al.*, 2003; Abdel-Ghany *et al.*, 2005).

PAA1 and PAA2 carry N-terminal MxCxxC metal-binding domains, CPC ion-transduction motifs, classical P-type ATPase phosphorylation and phosphatase domains, and ATP-binding sites. PAA1 is proposed to localize to the envelope membranes and PAA2 to the thylakoid membrane based on *in vitro* import experiments and localization of GFP fusion proteins (Shikanai *et al.*, 2003; Abdel-Ghany *et al.*, 2005). This distribution is analogous to the localization of two copper transporting P-type ATPases in cyanobacteria, CtaA and PacS. The former is located in the cell membrane and functions in copper acquisition, whereas the latter is located in the thylakoid membrane and functions to deliver copper to the lumen (Kanamaru *et al.*, 1994; Phung *et al.*, 1994; Tottey *et al.*, 2001). The phenotypes of *paa1* and *paa2* mutants are entirely consistent with this model. Plants carrying mutations in *PAA1* show reduced abundance of both plastocyanin and CuZnSOD whereas *paa2* plants have less plastocyanin (Shikanai *et al.*, 2003; Abdel-Ghany *et al.*, 2005). Although copper transporting activity has not been shown for either PAA1 or PAA2, the impact of copper nutrition on the phenotype is consistent with their function in copper transport. Specifically, the phenotype of *paa2* alleles was exacerbated in medium containing low copper and suppressed in medium containing extra copper. Also, while both *paa1* and *paa2* mutants had normal leaf copper content, when metal content was analyzed after sub-cellular fractionation, *paa1* chloroplasts were found to have less copper than those from wild-type, and *paa2* showed less copper in the thylakoid fractions but essentially normal copper content in intact chloroplasts. PAA1 is expressed in all organ types, reflecting a need for copper in all plastid types, while PAA2 expression was detected only in green organs, consistent

with a function in photosynthesis (Abdel-Ghany *et al.*, 2005).

Interestingly, the genome of *Cyanidioschizon merolae* lacks a PAA1 or PAA2 homolog, but this is not incompatible with the fact that the organism also lacks a gene for plastocyanin and must use only a *c*-type cytochrome for photosynthesis (see below) (Hanikenne *et al.*, 2005).

2. Chaperones

The concept of a copper chaperone for delivery of copper between proteins developed through genetic analysis of copper homeostasis in *S. cerevisiae* and subsequent analysis of the function of homologous proteins in other organisms (O'Halloran and Culotta, 2000). Yeast Atx1p (homologs known as HAH1, Atox1 and CCH) is a small protein containing a metal-binding site that interacts specifically with the metal-binding site on Ccc2p (a P-type ATPase). By analogy, a stromal chaperone for copper delivery from the envelope to the thylakoid, and perhaps another in the lumenal compartment for delivery from PAA2 to apoplastocyanin, is predicted. In cyanobacteria, an Atx1-related molecule, identified through a two-hybrid interaction with the metal-binding domains of PacS and CtaA, was shown to function as a copper chaperone for plastocyanin and cytochrome oxidase assembly (Tottey *et al.*, 2002). Proteins carrying candidate copper chaperone motifs can be identified in the *Arabidopsis* genome, but these have not yet been analyzed functionally. It is also possible that because of the small size of Atx1-like copper chaperones, some candidate molecules have not been predicted accurately or they have escaped detection because of the less significant BLAST (similarity) scores.

A small copper-binding protein, related to a copper homeostasis factor in bacteria called CutA, was shown to be chloroplast-localized (Burkhead *et al.*, 2003). Its physiological function has not been deduced, but it may well function in copper trafficking and recycling (see below).

B. Cu Deficiency and Regulation by Cu

1. *Chlamydomonas*

Because plastocyanin is the most abundant copper protein in a photosynthetic cell, it is a prime target in the face of copper-deficiency. In cyanobacteria and green algae, there is a well-regulated "back up" system, in which a heme protein, called cytochrome *c*₆,

is induced in copper-deficiency to compensate for the loss of plastocyanin in the electron transfer chain (Wood, 1978; Sandmann *et al.*, 1983; Merchant, 1998). Accordingly, copper is not essential for photosynthesis in these organisms. The regulatory events have been studied most thoroughly in the *Chlamydomonas* model (Merchant, 1998). The *CYC6* gene for cytochrome c_6 is associated with copper-response elements that serve as binding sites for a transcriptional activator, Crr1, in copper-deficient cells (Quinn and Merchant, 1995; Quinn *et al.*, 2000; Eriksson *et al.*, 2004). In this situation, plastocyanin is rapidly degraded because of the reduced thermodynamic stability and increased protease-susceptibility of the apo-protein vs. the holo-protein, and the Crr1-dependent expression of a degrading activity (Merchant and Bogorad, 1986; HH Li and Merchant, 1995; Eriksson *et al.*, 2004). In *crr1* mutants, apo- and some holo-plastocyanin accumulate even in copper-deficient cells, owing presumably to the lack of the protease. The *crr1* mutation, therefore, by affecting the “salvage” of copper from plastocyanin, has an impact on respiratory growth in addition to its impact on photosynthesis owing to loss of *CYC6* expression (S. Tottey, S. Nakamoto, J. Kropat and S. Merchant, unpublished results).

Besides regulating plastocyanin and cytochrome c_6 abundance, Crr1 also controls the expression of *CPX1* and *CHL27A/CHL27B*, encoding oxygen-dependent enzymes (coproporphyrinogen oxidase and aerobic oxidative cyclase) in the tetrapyrrole biosynthetic pathway (Eriksson *et al.*, 2004). *CPX1* encodes a plastid-targeted isoform that is about 10- to 20-fold up-regulated in copper-deficiency, while the expression of *CPX2*, encoding possibly a mitochondrial isoform, is unaffected by copper (Hill and Merchant, 1995; J. Kropat and S. Merchant, unpublished results). Copper nutrition and Crr1 reciprocally regulate the expression of *CHL27A* and *CHL27B* (Moseley *et al.*, 2000; Moseley *et al.*, 2002b). Both isozymes are plastid-localized but they may be differently distributed between the envelope and thylakoid membranes within the plastid (M. Allen and S. Merchant, unpublished results). The rationale for *CPX1* and *CHL27A/CHL27B* regulation by copper is not known, but it does point to a previously unrecognized connection between copper and the tetrapyrrole pathway.

2. *Arabidopsis*

Copper-deficiency has not been studied systematically in *Arabidopsis* but the work on PAA1 and PAA2 function revealed that the standard medium for *Arabidopsis*

growth in the laboratory is probably slightly copper-deficient (Abdel-Ghany *et al.*, 2005). The addition of copper to that medium stimulates the accumulation of both plastocyanin and CuZnSOD with a more noticeable effect for SOD. The authors concluded that under conditions of copper limitation there is preferential allocation of copper to plastocyanin (for which there is no substitute in *Arabidopsis*) vs. CuZnSOD (for which there is a substitute). In fact, the expression of FeSOD is increased to compensate for loss of CuZnSOD.

3. *Redistribution of Copper*

Several lines of evidence indicate that metals can be redistributed from the chloroplast to other organelles or even secreted from the cells. When copper-replete *Chlamydomonas* cells become deficient, copper is re-allocated from plastocyanin in the chloroplast to cytochrome oxidase in the mitochondrion (S. Tottey, S. Nakamoto, J. Kropat and S. Merchant, unpublished results). In vascular plants, the copper content of senescent tissue decreases. This process is correlated with an increase in the content of a copper chaperone, CCH, in the vascular tissue (Mira *et al.*, 2001). These processes probably require the action of transporters, chaperones or copper-binding proteins to move copper from a stable intracellular site in a protein, but the relevant molecules have not yet been identified and the process has not been subject to genetic analysis.

IV. Mn

Mn is nutritionally essential for all living organisms (Frieden, 1985; Marschner, 1995). It functions as a redox catalyst because it can occur stably in a cell in many different oxidation states, and this is its role in Mn-containing SOD and in PSII. Mn^{2+} can also activate water to generate a strong nucleophile for hydrolytic reactions (as in the enzyme arginase) or it can stabilize a leaving group (as in the nucleotide products of a glycosyl transferase reaction), but its role in these types of reactions in the plastid are not specifically described in the literature.

A. *Manganese Transport*

The bulk of the manganese in a photosynthetic cell is found in PSII in the chloroplast lumen. The mechanism of assembly of this cluster is not well understood even though PSII biogenesis has been subject to considerable genetic dissection in both cyanobacteria

and *Chlamydomonas* (Pakrasi, 1995). Pakrasi and co-workers approached this problem in the *Synechocystis* model and discovered the MntABC system for manganese ion uptake into bacterial cells. In more recent work, they show that cyanobacteria contain two pools of manganese, a storage pool that is released upon treatment with EDTA but whose maintenance is energy-dependent, and a second pool in PSII that is derived from the storage pool (Keren *et al.*, 2002). By analogy, there must be mechanisms for Mn^{2+} transport into the chloroplast across the inner envelope membrane plus a mechanism for transport across the thylakoid membrane. The identity of the transporters in the chloroplast is unknown. The MntABC system appears to be strictly bacterial, indicating the operation of another system for chloroplasts.

The Nramp proteins (reviewed by Williams *et al.*, 2000; Forbes and Gros, 2001) are excellent candidates for a manganese delivery system to plastids. These molecules are proton-coupled divalent cation transporters that show broad substrate specificity in many *in vitro* experiments but it is likely that some members of the gene family are Mn^{2+} -specific *in vivo*. The bacterial homologs of the Nramps, called MntH, indeed appear to be Mn^{2+} selective (Kehres and Maguire, 2003) and Nramp homologs in *Chlamydomonas* do show increased expression in response to manganese-deficiency (M. Allen, S. Tottey, J. Kropat, J. del Campo and S. Merchant, unpublished results). Plant genomes contain multiple Nramp homologs with functionally distinct roles based on sub-cellular location, organ-specific pattern of expression, metal specificity, and pH sensitivity (Belouchi *et al.*, 1997; Curie *et al.*, 2000; Thomine *et al.*, 2000; Thomine *et al.*, 2003). While some members of the family are likely involved in iron homeostasis, others could function in manganese metabolism. But the role of plant *NRAMP* expression and function in manganese nutrition has received less attention.

A possibility for manganese acquisition by the plastid, hinted at by the intracellular organelle localization of Nramp homologs Smf1p and Smf2p in *S. cerevisiae* (reviewed by Van Ho *et al.*, 2002) is that one or more Nramps may be involved. In this context, it is worth noting that mitochondria also have a significant manganese requirement (e.g. for MnSOD) and the question of allocation of manganese to plastids vs. mitochondria in plants has not been addressed. In *S. cerevisiae*, a member of the carrier family has been proposed as a facilitator for mitochondrial manganese acquisition for MnSOD biogenesis (Luk *et al.*, 2003). It is not known whether Mtm1p is actually a Mn^{2+} transporter. The

mitochondrial carriers are evolutionarily distinct from most of the known plastid inner envelope translocators, and so it does not necessarily follow that a homolog of *S. cerevisiae* *MTM1* would function in plastid Mn-protein assembly. The diversity of Mn^{2+} transporters known in nature—MntA, MntH and perhaps the Mtm1p carrier—leaves open the possibility that a completely novel molecule operates in the plastid.

A recent comparative analysis of algal genomes revealed members of the cation diffusion facilitator family (called MTP proteins in plants) that may be manganese transporters, and it is suggested that one or more of these molecules could be plastid-localized (Hanikenne *et al.*, 2005).

B. Manganese Deficiency

The importance of manganese in the photochemical reactions of photosynthesis was recognized half a century ago because of the impact of Mn deficiency on phototrophic growth and oxygen evolution in algae (Pirson, 1955). The symptoms of Mn-deficiency in plants are noted as leaf discoloration, which implies an impact at the level of the chloroplast, but the biochemical consequences of deficiency have not been investigated. Mn-deficient *Chlamydomonas* cells show loss of PSII and MnSOD activity and a sensitivity to peroxides but not paraquat or Rose Bengal (M. Allen, S. Tottey, J. Kropat, J. del Campo and S. Merchant, unpublished results). Whether the oxidative stress occurs at the level of plastid or mitochondrion redox metabolism is not known.

V. Questions for Future Investigation

The metabolism of the transition elements is intimately inter-related. For instance, in many organisms, including algae, fungi and mammals (although not plants), a copper-containing enzyme is required for high affinity iron uptake (Askwith and Kaplan, 1998; La Fontaine *et al.*, 2002). Therefore, copper-deficiency generates secondarily an iron-deficiency. A connection between copper and zinc metabolism is known in humans, where excess zinc in the diet blocks copper intake (Kumar *et al.*, 2003). Recently we noted a role for manganese in iron assimilation in *Chlamydomonas* (M. Allen, S. Tottey, J. Kropat, J. del Campo and S. Merchant, unpublished results). The use of microarray and proteomic approaches to study metal homeostasis at a whole genome level is ideal for the discovery of such inter-relationships.

In a recent microarray study on iron-deficient yeast, the concept of metabolic re-modeling was noted where certain iron-utilizing pathways are down-regulated in favor of parallel pathways that are less iron-dependent (Shakoury-Elizeh *et al.*, 2004). This phenomenon is well known in the context of the photosynthetic apparatus where flavodoxin can substitute for ferredoxin or cytochrome *c*₆ for plastocyanin, and for the SODs where Mn-SOD is up-regulated to compensate for the loss of CuZnSOD in the copper-deficient rat (Hutber *et al.*, 1977; Wood, 1978; Merchant and Bogorad, 1987; Bottin and Lagoutte, 1992; Lai *et al.*, 1994). It is possible that there are back-up systems for other metalloenzymes in nature and these may also be discovered through whole genome analyses.

Acknowledgments

The Department of Agriculture, the Department of Energy, and the National Institutes of Health have supported research in my laboratory on trace metal nutrition in *Chlamydomonas*. The present members of the group have made important contributions to discussions of the ideas presented in this chapter.

References

- Abdel-Ghany S, Müller-Moulé P, Niyogi KK, Pilon M and Shikanai T (2005) Two P-type ATPases are required for copper delivery in *Arabidopsis thaliana* chloroplasts. *Plant Cell* 17: 1233–1251
- Agar JN, Yuvaniyama P, Jack RF, Cash VL, Smith AD, Dean DR and Johnson MK (2000) Modular organization and identification of a mononuclear iron-binding site within the NifU protein. *J Biol Inorg Chem* 5: 167–177
- Ali V, Shigeta Y, Tokumoto U, Takahashi Y and Nozaki T (2004) An intestinal parasitic protist, *Entamoeba histolytica*, possesses a non-redundant nitrogen fixation-like system for iron-sulfur cluster assembly under anaerobic conditions. *J Biol Chem* 279: 16863–16874
- Amann K, Lezhneva L, Wanner G, Herrmann RG and Meurer J (2004) ACCUMULATION OF PHOTOSYSTEM ONE1, a member of a novel gene family, is required for accumulation of [4Fe-4S] cluster-containing chloroplast complexes and antenna proteins. *Plant Cell* 16: 3084–3097
- Askwith C and Kaplan J (1998) Iron and copper transport in yeast and its relevance to human disease. *Trends Biochem Sci* 23: 135–138
- Balk J, Pierik AJ, Netz DJ, Mühlenhoff U and Lill R (2004) The hydrogenase-like Nar1p is essential for maturation of cytosolic and nuclear iron-sulphur proteins. *EMBO J* 23: 2105–2115
- Beale SI (1999) Enzymes of chlorophyll biosynthesis. *Photosynth Res* 60: 43–73
- Belouchi A, Kwan T and Gros P (1997) Cloning and characterization of the *OsNramp* family from *Oryza sativa*, a new family of membrane proteins possibly implicated in the transport of metal ions. *Plant Mol Biol* 33: 1085–1092
- Berthold DA and Stenmark P (2003) Membrane-bound diiron carboxylate proteins. *Annu Rev Plant Biol* 54: 497–517
- Bibby TS, Nield J and Barber J (2001) Iron deficiency induces the formation of an antenna ring around trimeric photosystem I in cyanobacteria. *Nature* 412: 743–745
- Bienfait HF and van den Briel ML (1980) Rapid mobilization of ferritin iron by ascorbate in the presence of oxygen. *Biochim Biophys Acta* 631: 507–510
- Boekema EJ, Hifney A, Yakushevskaya AE, Piotrowski M, Keegstra W, Berry S, Michel KP, Pistorius EK and Kruijff J (2001) A giant chlorophyll-protein complex induced by iron deficiency in cyanobacteria. *Nature* 412: 745–748
- Bogorad L, Pires G, Swift H and McIlrath WJ (1958) The structure of chloroplasts in leaf tissue of iron deficient *Xanthium*. *Brookhaven Symp Biol* 11: 132–137
- Bottin H and Lagoutte B (1992) Ferredoxin and flavodoxin from the cyanobacterium *Synechocystis* sp PCC 6803. *Biochim Biophys Acta* 1101: 48–56
- Briat J-F and Lobréaux S (1997) Iron transport and storage in plants. *Trends Plant Sci* 2: 187–193
- Briat JF, Lobréaux S, Grignon N and Vansuyt G (1999) Regulation of plant ferritin synthesis: how and why. *Cell Mol Life Sci* 56: 155–166
- Burkhead JL, Abdel-Ghany SE, Morrill JM, Pilon-Smits EA and Pilon M (2003) The *Arabidopsis thaliana* CUTA gene encodes an evolutionarily conserved copper binding chloroplast protein. *Plant J* 34: 856–867
- Chow KS, Singh DP, Roper JM and Smith AG (1997) A single precursor protein for ferrochelatase-I from *Arabidopsis* is imported *in vitro* into both chloroplasts and mitochondria. *J Biol Chem* 272: 27565–27571
- Cornah JE, Terry MJ and Smith AG (2003) Green or red: what stops the traffic in the tetrapyrrole pathway? *Trends Plant Sci* 8: 224–230
- Crisp RJ, Pollington A, Galea C, Jaron S, Yamaguchi-Iwai Y and Kaplan J (2003) Inhibition of heme biosynthesis prevents transcription of iron uptake genes in yeast. *J Biol Chem* 278: 45499–45506
- Cunningham FX and Gantt E (1998) Genes and enzymes of carotenoid biosynthesis in plants. *Annu Rev Plant Physiol Plant Mol Biol* 49: 557–583
- Cupp-Vickery JR, Silberg JJ, Ta DT and Vickery LE (2004) Crystal structure of IscA, an iron-sulfur cluster assembly protein from *Escherichia coli*. *J Mol Biol* 338: 127–137
- Curie C and Briat JF (2003) Iron transport and signaling in plants. *Annu Rev Plant Biol* 54: 183–206
- Curie C, Alonso JM, Le Jean M, Ecker JR and Briat JF (2000) Involvement of NRAMP1 from *Arabidopsis thaliana* in iron transport. *Biochem J* 347: 749–755
- Duggan J and Gassman M (1974) Induction of porphyrin synthesis in etiolated bean leaves by chelators of iron. *Plant Physiol* 53: 206–215
- Ellis KE, Clough B, Saldanha JW and Wilson RJ (2001) Nifs and Sufs in malaria. *Mol Microbiol* 41: 973–981
- Eriksson M, Moseley JL, Tottey S, del Campo JA, Quinn JM, Kim Y and Merchant S (2004) Genetic dissection of nutritional copper signaling in *Chlamydomonas* distinguishes regulatory and target genes. *Genetics* 168: 795–807

- Flügge U-I, Häusler RE, Ludewig F and Fischer K (2003) Functional genomics of phosphate antiport systems of plastids. *Physiol Plant* 118: 475–482
- Fobis-Loisy I, Aussel L and Briat JF (1996) Post-transcriptional regulation of plant ferritin accumulation in response to iron as observed in the maize mutant *ys1*. *FEBS Lett* 397: 149–54
- Forbes JR and Gros P (2001) Divalent-metal transport by NRAMP proteins at the interface of host-pathogen interactions. *Trends Microbiol* 9: 397–403
- Foury F and Roganti T (2002) Deletion of the mitochondrial carrier genes *MRS3* and *MRS4* suppresses mitochondrial iron accumulation in a yeast frataxin-deficient strain. *J Biol Chem* 277: 24475–24483
- Fox BG (1998) Catalysis by non-heme iron. In: Sinnott M (ed) *Comprehensive Biological Catalysis*, pp 261–348. Academic Press, London
- Frankenberg-Dinkel N (2004) Bacterial heme oxygenases. *Antioxid Redox Signal* 6: 825–834
- Franklin KA, Linley PJ, Montgomery BL, Lagarias JC, Thomas B, Jackson SD and Terry MJ (2003) Misregulation of tetrapyrrole biosynthesis in transgenic tobacco seedlings expressing mammalian biliverdin reductase. *Plant J* 35: 717–728
- Frazzon J and Dean DR (2003) Formation of iron-sulfur clusters in bacteria: an emerging field in bioinorganic chemistry. *Curr Opin Chem Biol* 7: 166–173
- Frazzon J, Fick JR and Dean DR (2002) Biosynthesis of iron-sulphur clusters is a complex and highly conserved process. *Biochem Soc Trans* 30: 680–685
- Frieden E (1985) New perspectives on the essential trace elements. *J Chem Ed* 62: 917–923
- Froehlich JE, Itoh A and Howe GA (2001) Tomato allene oxide synthase and fatty acid hydroperoxide lyase, two cytochrome P450s involved in oxylipin metabolism, are targeted to different membranes of chloroplast envelope. *Plant Physiol* 125: 306–317
- Garland SA, Hoff K, Vickery LE and Culotta VC (1999) *Saccharomyces cerevisiae* *ISU1* and *ISU2*: members of a well-conserved gene family for iron-sulfur cluster assembly. *J Mol Biol* 294: 897–907
- Gray J, Wardzala E, Yang M, Reinbothe S, Haller S and Pauli F (2004) A small family of LLS1-related non-heme oxygenases in plants with an origin amongst oxygenic photosynthesizers. *Plant Mol Biol* 54: 39–54
- Gunshin H, Mackenzie B, Berger UV, Gunshin Y, Romero MF, Boron WF, Nussberger S, Gollan JL and Hediger MA (1997) Cloning and characterization of a mammalian proton-coupled metal-ion transporter. *Nature* 388: 482–488
- Hanikenne M, Krämer U, Demoulin V and Baurain D (2005) A comparative inventory of metal transporters in the green alga *Chlamydomonas reinhardtii* and the red alga *Cyanidioschizon merolae*. *Plant Physiol* 137: 428–446
- Happe T, Mosler B and Naber JD (1994) Induction, localization and metal content of hydrogenase in the green alga *Chlamydomonas reinhardtii*. *Eur J Biochem* 222: 769–774
- Hausinger RP (2004) Fe(II)/ α -ketoglutarate-dependent hydroxylases and related enzymes. *Crit Rev Biochem Mol Biol* 39: 21–68
- Hausmann A, Netz DJA, Balk J, Pierik AJ, Mühlhoff U and Lill R (2005) The eukaryotic P loop NTPase Nbp35: an essential component of the cytosolic and nuclear iron-sulfur protein assembly machinery. *Proc Natl Acad Sci USA* 102: 3266–3271
- Helliwell CA, Sullivan JA, Mould RM, Gray JC, Peacock WJ and Dennis ES (2001) A plastid envelope location of *Arabidopsis* ent-kaurene oxidase links the plastid and endoplasmic reticulum steps of the gibberellin biosynthesis pathway. *Plant J* 28: 201–208
- Hill KL and Merchant S (1995) Coordinate expression of coproporphyrinogen oxidase and cytochrome c6 in the green alga *Chlamydomonas reinhardtii* in response to changes in copper availability. *EMBO J* 14: 857–865
- Hippler M, Biehler K, Krieger-Liszczay A, van Dillewijn J and Rochaix JD (2000) Limitation in electron transfer in photosystem I donor side mutants of *Chlamydomonas reinhardtii*. Lethal photo-oxidative damage in high light is overcome in a suppressor strain deficient in the assembly of the light harvesting complex. *J Biol Chem* 275: 5852–5859
- Hirayama T, Kieber JJ, Hirayama N, Kogan M, Guzman P, Nourizadeh S, Alonso JM, Dailey WP, Dancis A and Ecker JR (1999) RESPONSIVE-TO-ANTAGONIST1, a Menkes/Wilson disease-related copper transporter, is required for ethylene signaling in *Arabidopsis*. *Cell* 97: 383–393
- Hoff KG, Silberg JJ and Vickery LE (2000) Interaction of the iron-sulfur cluster assembly protein IscU with the Hsc66/Hsc20 molecular chaperone system of *Escherichia coli*. *Proc Natl Acad Sci USA* 97: 7790–7795
- Hutber GN, Hutson KG and Rogers LJ (1977) Effect of iron deficiency on levels of two ferredoxins and flavodoxin in a cyanobacterium. *FEMS Microbiol Lett* 1: 193–196
- Jackson C, Dench J, Moore AL, Halliwell B, Foyer CH and Hall DO (1978) Subcellular localisation and identification of superoxide dismutase in the leaves of higher plants. *Eur J Biochem* 91: 339–344
- Jäger-Vottero P, Dorne A-J, Jordanov J, Douce R and Joyard J (1997) Redox chains in chloroplast envelope membranes: spectroscopic evidence for the presence of electron carriers, including iron-sulfur centers. *Proc Natl Acad Sci USA* 94: 1597–1602
- Jensen LT and Culotta VC (2000) Role of *Saccharomyces cerevisiae* *ISA1* and *ISA2* in iron homeostasis. *Mol Cell Biol* 20: 3918–3927
- Jensen PE, Gilpin M, Knoetzel J and Scheller HV (2000) The PSI-K subunit of photosystem I is involved in the interaction between light-harvesting complex I and the photosystem I reaction center core. *J Biol Chem* 275: 24701–24708
- Kaim W and Schwederski B (1994) *Bioinorganic Chemistry: Inorganic Elements in the Chemistry of Life—An Introduction and Guide*. John Wiley and Sons, New York
- Kanamaru K, Kashiwagi S and Mizuno T (1994) A copper-transporting P-type ATPase found in the thylakoid membrane of the cyanobacterium *Synechococcus* species PCC7942. *Mol Microbiol* 13: 369–377
- Kehres DG and Maguire ME (2003) Emerging themes in manganese transport, biochemistry and pathogenesis in bacteria. *FEMS Microbiol Rev* 27: 263–290
- Keren N, Kidd MJ, Penner-Hahn JE and Pakrasi HB (2002) A light-dependent mechanism for massive accumulation of manganese in the photosynthetic bacterium *Synechocystis* sp. PCC 6803. *Biochemistry* 41: 15085–15092
- Kerfeld CA and Krogmann DW (1998) Photosynthetic cytochromes *c* in cyanobacteria, algae and plants. *Annu Rev Plant Physiol Plant Mol Biol* 49: 397–425

- Kieselbach T, Hagman K, Andersson B and Schröder WP (1998) The thylakoid lumen of chloroplasts. Isolation and characterization. *J Biol Chem* 273: 6710–6716
- Kispal G, Csere P, Prohl C and Lill R (1999) The mitochondrial proteins Atm1p and Nfs1p are essential for biogenesis of cytosolic Fe/S proteins. *EMBO J* 18: 3981–3989
- Knight SA, Sepuri NB, Pain D and Dancis A (1998) Mt-Hsp70 homolog, Ssc2p, required for maturation of yeast frataxin and mitochondrial iron homeostasis. *J Biol Chem* 273: 18389–18393
- Kumar N, Gross JB Jr and Ahlskog JE (2003) Myelopathy due to copper deficiency. *Neurology* 61: 273–274
- Kunji ER (2004) The role and structure of mitochondrial carriers. *FEBS Lett* 564: 239–244
- Kushnir S, Babiychuk E, Storozhenko S, Davey MW, Papenbrock J, De Rycke R, Engler G, Stephan UW, Lange H, Kispal G, Lill R and Van Montagu M (2001) A mutation of the mitochondrial ABC transporter Stal1 leads to dwarfism and chlorosis in the *Arabidopsis* mutant *starik*. *Plant Cell* 13: 89–100
- La Fontaine S, Quinn JM, Nakamoto SS, Page MD, Göhre V, Moseley JL, Kropat J and Merchant S (2002) Copper-dependent iron assimilation pathway in the model photosynthetic eukaryote *Chlamydomonas reinhardtii*. *Eukaryot Cell* 1: 736–757
- La Roche J, Boyd PW, McKay RML and Geider RJ (1996) Flavodoxin as an in situ marker for iron stress in phytoplankton. *Nature* 382: 802–805
- Lai CC, Huang WH, Askari A, Wang Y, Sarvazyan N, Klevay LM and Chiu TH (1994) Differential regulation of superoxide dismutase in copper-deficient rat organs. *Free Radic Biol Med* 16: 613–620
- Lange H, Kaut A, Kispal G and Lill R (2000) A mitochondrial ferredoxin is essential for biogenesis of cellular iron-sulfur proteins. *Proc Natl Acad Sci USA* 97: 1050–1055
- Last DI and Gray JC (1989) Plastocyanin is encoded by a single-copy gene in the pea haploid genome. *Plant Mol Biol* 12: 655–666
- Laudenbach DE and Straus NA (1988) Characterization of a cyanobacterial iron stress-induced gene similar to *psbC*. *J Bacteriol* 170: 5018–5026
- Laudenbach DE, Reith ME and Straus NA (1988) Isolation, sequence analysis and transcriptional studies of the flavodoxin gene from *Anacystis nidulans* R2. *J Bacteriol* 170: 258–265
- Laulhere JP, Laboure AM, Van Wuytswinkel O, Gagnon J and Briat JF (1992) Purification, characterization and function of bacterioferritin from the cyanobacterium *Synechocystis* PCC 6803. *Biochem J* 281: 785–793
- Lawson DM, Treffry A, Artymiuk PJ, Harrison PM, Yewdall SJ, Luzzago A, Cesareni G, Levi S and Arosio P (1989) Identification of the ferroxidase centre in ferritin. *FEBS Lett* 254: 207–210
- Léon S, Touraine B, Briat JF and Lobréaux S (2002) The *AtNFS2* gene from *Arabidopsis thaliana* encodes a NifS-like plastidial cysteine desulphurase. *Biochem J* 366: 557–564
- Léon S, Touraine B, Ribot C, Briat JF and Lobréaux S (2003) Iron-sulphur cluster assembly in plants: distinct NFU proteins in mitochondria and plastids from *Arabidopsis thaliana*. *Biochem J* 371: 823–830
- Lermontova I, Kruse E, Mock HP and Grimm B (1997) Cloning and characterization of a plastidial and a mitochondrial isoform of tobacco protoporphyrinogen IX oxidase. *Proc Natl Acad Sci USA* 94: 8895–8900
- Lescure AM, Proudhon D, Pesey H, Ragland M, Theil EC and Briat JF (1991) Ferritin gene transcription is regulated by iron in soybean cell cultures. *Proc Natl Acad Sci USA* 88: 8222–8226
- Lesuisse E, Lyver ER, Knight SA and Dancis A (2004) Role of *YHMI*, encoding a mitochondrial carrier protein, in iron distribution of yeast. *Biochem J* 378: 599–607
- Levi S, Corsi B, Bosisio M, Invernizzi R, Volz A, Sanford D, Arosio P and Drysdale J (2001) A human mitochondrial ferritin encoded by an intronless gene. *J Biol Chem* 276: 24437–24440
- Lezhneva L, Amann K and Meurer J (2004) The universally conserved HCF101 protein is involved in assembly of [4Fe-4S]-cluster-containing complexes in *Arabidopsis thaliana* chloroplasts. *Plant J* 37: 174–185
- Li HH and Merchant S (1995) Degradation of plastocyanin in copper-deficient *Chlamydomonas reinhardtii*. *J Biol Chem* 270: 23504–23510
- Li HM, Theg SM, Bauerle CM and Keegstra K (1990) Metal-ion-center assembly of ferredoxin and plastocyanin in isolated chloroplasts. *Proc Natl Acad Sci USA* 87: 6748–6752
- Li J, Kogan M, Knight SA, Pain D and Dancis A (1999) Yeast mitochondrial protein, Nfs1p, coordinately regulates iron-sulfur cluster proteins, cellular iron uptake and iron distribution. *J Biol Chem* 274: 33025–33034
- Lill R and Kispal G (2000) Maturation of cellular Fe-S proteins: an essential function of mitochondria. *Trends Biochem Sci* 25: 352–356
- Lobréaux S and Briat JF (1991) Ferritin accumulation and degradation in different organs of pea (*Pisum sativum*) during development. *Biochem J* 274: 601–606
- Loiseau L, Ollagnier-de-Choudens S, Nachin L, Fontecave M and Barras F (2003) Biogenesis of Fe-S cluster by the bacterial Suf system: SufS and SufE form a new type of cysteine desulfurase. *J Biol Chem* 278: 38352–38359
- Luk E, Carroll M, Baker M and Culotta VC (2003) Manganese activation of superoxide dismutase 2 in *Saccharomyces cerevisiae* requires *MTM1*, a member of the mitochondrial carrier family. *Proc Natl Acad Sci USA* 100: 10353–10357
- Machold O (1971) Lamellar proteins of green and chlorotic chloroplasts as affected by iron deficiency and antibiotics. *Biochim Biophys Acta* 238: 324–331
- Malkin R and Rabinowitz JC (1966) The reconstitution of clostridial ferredoxin. *Biochem Biophys Res Commun* 23: 822–827
- Marschner H (1995) Mineral Nutrition of Higher Plants. Academic Press, London
- Masuda T, Goto F and Yoshihara T (2001) A novel plant ferritin subunit from soybean that is related to a mechanism in iron release. *J Biol Chem* 276: 19575–19579
- Merchant S (1998) Synthesis of metalloproteins involved in photosynthesis: plastocyanin and cytochromes. In: Rochaix J-D, Goldschmidt-Clermont M and Merchant S (eds) *The Molecular Biology of Chloroplasts and Mitochondria in Chlamydomonas*, pp 597–611. Kluwer Academic Publishers, Dordrecht, the Netherlands
- Merchant S and Bogorad L (1986) Rapid degradation of apoplastocyanin in Cu(II)-deficient cells of *Chlamydomonas reinhardtii*. *J Biol Chem* 261: 15850–15853
- Merchant S and Bogorad L (1987) Metal ion regulated gene expression: use of a plastocyanin-less mutant of *Chlamydomonas*

- reinhardtii* to study the Cu(II)-dependent expression of cytochrome *c-552*. *EMBO J* 6: 2531–2535
- Merchant S and Dreyfuss BW (1998) Posttranslational assembly of photosynthetic metalloproteins. *Annu Rev Plant Physiol Plant Mol Biol* 49: 25–51
- Michel KP and Pistorius EK (2004) Adaptation of the photosynthetic electron transport chain in cyanobacteria to iron deficiency: the function of *IdiA* and *IsiA*. *Physiol Plant* 120: 36–50
- Mira H, Martínez-García F and Peñarrubia L (2001) Evidence for the plant-specific intercellular transport of the *Arabidopsis* copper chaperone CCH. *Plant J* 25: 521–528
- Moseley J, Quinn J, Eriksson M and Merchant S (2000) The *Crd1* gene encodes a putative di-iron enzyme required for photosystem I accumulation in copper deficiency and hypoxia in *Chlamydomonas reinhardtii*. *EMBO J* 19: 2139–2151
- Moseley JL, Allinger T, Herzog S, Hoerth P, Wehinger E, Merchant S and Hippler M (2002a) Adaptation to Fe-deficiency requires remodeling of the photosynthetic apparatus. *EMBO J* 21: 6709–6720
- Moseley JL, Page MD, Alder NP, Eriksson M, Quinn J, Soto F, Theg SM, Hippler M and Merchant S (2002b) Reciprocal expression of two candidate di-iron enzymes affecting photosystem I and light-harvesting complex accumulation. *Plant Cell* 14: 673–688
- Mühlenhoff U, Richhardt N, Gerber J and Lill R (2002) Characterization of iron-sulfur protein assembly in isolated mitochondria. A requirement for ATP, NADH and reduced iron. *J Biol Chem* 277: 29810–29816
- Mühlenhoff U, Stadler JA, Richhardt N, Seubert A, Eickhorst T, Schweyen RJ, Lill R and Wiesenberger G (2003) A specific role of the yeast mitochondrial carriers MRS3/4p in mitochondrial iron acquisition under iron-limiting conditions. *J Biol Chem* 278: 40612–40620
- Nachin L, Loiseau L, Expert D and Barras F (2003) SufC: an unorthodox cytoplasmic ABC/ATPase required for [Fe-S] biogenesis under oxidative stress. *EMBO J* 22: 427–437
- Nishio JN, Abadía J and Terry N (1985) Chlorophyll-proteins and electron transport during iron nutrition-mediated chloroplast development. *Plant Physiol* 78: 296–299
- O'Halloran TV and Culotta VC (2000) Metallochaperones, an intracellular shuttle service for metal ions. *J Biol Chem* 275: 25057–25060
- Ollagnier-de-Choudens S, Mattioli T, Takahashi Y and Fontecave M (2001) Iron-sulfur cluster assembly: characterization of *IscA* and evidence for a specific and functional complex with ferredoxin. *J Biol Chem* 276: 22604–22607
- Outten FW, Djaman O and Storz G (2004) A *suf* operon requirement for Fe-S cluster assembly during iron starvation in *Escherichia coli*. *Mol Microbiol* 52: 861–872
- Pakrasi HB (1995) Genetic analysis of the form and function of photosystem I and photosystem II. *Annu Rev Genet* 29: 755–776
- Perl-Treves R and Galun E (1991) The tomato Cu, Zn superoxide dismutase genes are developmentally regulated and respond to light and stress. *Plant Mol Biol* 17: 745–760
- Petit JM, Briat JF and Lobreaux S (2001) Structure and differential expression of the four members of the *Arabidopsis thaliana* ferritin gene family. *Biochem J* 359: 575–582
- Phung LT, Ajlani G and Haselkorn R (1994) P-type ATPase from the cyanobacterium *Synechococcus* 7942 related to the human Menkes and Wilson disease gene products. *Proc Natl Acad Sci USA* 91: 9651–9654
- Picault N, Hodges M, Palmieri L and Palmieri F (2004) The growing family of mitochondrial carriers in *Arabidopsis*. *Trends Plant Sci* 9: 138–146
- Pilon M, de Kruijff B and Weisbeek PJ (1992) New insights into the import mechanism of the ferredoxin precursor into chloroplasts. *J Biol Chem* 267: 2548–2556
- Pilon-Smits EA, Garifullina GF, Abdel-Ghany S, Kato S, Mihara H, Hale KL, Burkhead JL, Esaki N, Kurihara T and Pilon M (2002) Characterization of a NifS-like chloroplast protein from *Arabidopsis*. Implications for its role in sulfur and selenium metabolism. *Plant Physiol* 130: 1309–1318
- Pirson A (1955) Functional aspects in mineral nutrition of green plants. *Annu Rev Plant Physiol* 6: 71–114
- Plank DW, Gengenbach BG and Gronwald JW (2001) Effect of iron on activity of soybean multi-subunit acetyl-coenzyme A carboxylase. *Physiol Plant* 112: 183–194
- Poschwitz MC, King PW, Smolinski SL, Zhang L, Seibert M and Ghirardi ML (2004) Discovery of two novel radical *S*-adenosylmethionine proteins required for the assembly of an active [Fe] hydrogenase. *J Biol Chem* 279: 25711–25720
- Poss KD and Tonegawa S (1997) Heme oxygenase 1 is required for mammalian iron reutilization. *Proc Natl Acad Sci USA* 94: 10919–10924
- Protchenko O and Philpott CC (2003) Regulation of intracellular heme levels by *HMX1*, a homologue of heme oxygenase, in *Saccharomyces cerevisiae*. *J Biol Chem* 278: 36582–36587
- Quinn JM and Merchant S (1995) Two copper-responsive elements associated with the *Chlamydomonas* *Cyc6* gene function as targets for transcriptional activators. *Plant Cell* 7: 623–638
- Quinn JM, Barraco P, Eriksson M and Merchant S (2000) Coordinate copper- and oxygen-responsive *Cyc6* and *Cpx1* expression in *Chlamydomonas* is mediated by the same element. *J Biol Chem* 275: 6080–6089
- Ragland M and Theil EC (1993) Ferritin (mRNA, protein) and iron concentrations during soybean nodule development. *Plant Mol Biol* 21: 555–560
- Ragland M, Briat JF, Gagnon J, Laulhere JP, Massenet O and Theil EC (1990) Evidence for conservation of ferritin sequences among plants and animals and for a transit peptide in soybean. *J Biol Chem* 265: 18339–18344
- Raven JA (1988) The iron and molybdenum use efficiencies of plant growth with different energy, carbon and nitrogen sources. *New Phytol* 109: 279–287
- Raven JA (1990) Predictions of Mn and Fe use efficiencies of phototrophic growth as a function of light availability for growth and of C assimilation pathway. *New Phytol* 116: 1–18
- Raven JA, Evans MCW and Korb RE (1999) The role of trace metals in photosynthetic electron transport in O₂-evolving organisms. *Photosynth Res* 60: 111–150
- Richaud C and Zabulon G (1997) The heme oxygenase gene (*pbsA*) in the red alga *Rhodella violacea* is discontinuous and transcriptionally activated during iron limitation. *Proc Natl Acad Sci USA* 94: 11736–11741
- Roy A, Solodovnikova N, Nicholson T, Antholine W and Walden WE (2003) A novel eukaryotic factor for cytosolic Fe-S cluster assembly. *EMBO J* 22: 4826–4835
- Sancenón V, Puig S, Mira H, Thiele DJ and Peñarrubia L (2003) Identification of a copper transporter family in *Arabidopsis thaliana*. *Plant Mol Biol* 51: 577–587

- Sancenón V, Puig S, Mateu-Andrés I, Dorcey E, Thiele DJ and Peñarrubia L (2004) The *Arabidopsis* copper transporter COPT1 functions in root elongation and pollen development. *J Biol Chem* 279: 15348–15355
- Sandmann G, Reck H, Kessler E and Boger P (1983) Distribution of plastocyanin and soluble plastidic cytochrome *c* in various classes of algae. *Arch Microbiol* 134: 23–27
- Schilke B, Voisine C, Beinert H and Craig E (1999) Evidence for a conserved system for iron metabolism in the mitochondria of *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA* 96: 10206–10211
- Shakoury-Elizeh M, Tiedeman J, Rashford J, Ferea T, Demeter J, Garcia E, Rolfes R, Brown PO, Botstein D and Philpott CC (2004) Transcriptional remodeling in response to iron deprivation in *Saccharomyces cerevisiae*. *Mol Biol Cell* 15: 1233–1243
- Shanklin J and Cahoon EB (1998) Desaturation and related modifications of fatty acids. *Annu Rev Plant Physiol Plant Mol Biol* 49: 611–641
- Shikanai T, Müller-Moulé P, Munekage Y, Niyogi KK and Pilon M (2003) PAA1, a P-type ATPase of *Arabidopsis*, functions in copper transport in chloroplasts. *Plant Cell* 15: 1333–1346
- Singh DP, Cornah JE, Hadingham S and Smith AG (2002) Expression analysis of the two ferrochelatase genes in *Arabidopsis* in different tissues and under stress conditions reveals their different roles in haem biosynthesis. *Plant Mol Biol* 50: 773–788
- Skaar EP, Gaspar AH and Schneewind O (2004) IsdG and IsdI, heme-degrading enzymes in the cytoplasm of *Staphylococcus aureus*. *J Biol Chem* 279: 436–443
- Skovran E and Downs DM (2003) Lack of the ApbC or ApbE protein results in a defect in Fe-S cluster metabolism in *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 185: 98–106
- Sofia HJ, Chen G, Hetzler BG, Reyes-Spindola JF and Miller NE (2001) Radical SAM, a novel protein superfamily linking unresolved steps in familiar biosynthetic pathways with radical mechanisms: functional characterization using new analysis and information visualization methods. *Nucleic Acids Res* 29: 1097–1106
- Spiller S and Terry N (1980) Limiting factors in photosynthesis. II. Iron stress diminishes photochemical capacity by reducing the number of photosynthetic units. *Plant Physiol* 65: 121–125
- Spiller SC, Castelfranco AM and Castelfranco PA (1982) Effects of iron and oxygen on chlorophyll biosynthesis I. In vivo observations on iron and oxygen-deficient plants. *Plant Physiol* 69: 107–111
- Stöckel J and Oelmüller R (2004) A novel protein for photosystem I biogenesis. *J Biol Chem* 279: 10243–10251
- Takahashi Y and Nakamura M (1999) Functional assignment of the ORF2-*iscS-iscU-iscA-hscB-hscA-fox-ORF3* gene cluster involved in the assembly of Fe-S clusters in *Escherichia coli*. *J Biochem* 126: 917–926
- Takahashi Y and Tokumoto U (2002) A third bacterial system for the assembly of iron-sulfur clusters with homologs in archaea and plastids. *J Biol Chem* 277: 28380–28383
- Takahashi Y, Mitsui A, Hase T and Matsubara H (1986) Formation of the iron-sulfur cluster of ferredoxin in isolated chloroplasts. *Proc Nat Acad Sci USA* 83: 2434–2437
- Takahashi Y, Mitsui A, Fujita Y and Matsubara H (1991a) Roles of ATP and NADPH in formation of the Fe-S cluster of spinach ferredoxin. *Plant Physiol* 95: 104–110
- Takahashi Y, Mitsui A and Matsubara H (1991b) Formation of the Fe-S cluster of ferredoxin in lysed spinach chloroplast. *Plant Physiol* 95: 97–103
- Tarantino D, Petit JM, Lobreaux S, Briat JF, Soave C and Murgia I (2003) Differential involvement of the IDRS cis-element in the developmental and environmental regulation of the *AtFer1* ferritin gene from *Arabidopsis*. *Planta* 217: 709–716
- Terry N (1980) Limiting factors in photosynthesis. I. Use of iron stress to control photosynthetic capacity in vivo. *Plant Physiol* 65: 114–120
- Terry N (1983) Limiting factors in photosynthesis. IV. Iron stress-mediated changes on light-harvesting and electron transport capacity and its effects on photosynthesis *in vivo*. *Plant Physiol* 71: 855–860
- Terry N and Abadia J (1986) Function of iron in chloroplasts. *J Plant Nutr* 9: 609–646
- Terry MJ, Linley PJ and Kohchi T (2002) Making light of it: the role of plant haem oxygenases in phytochrome chromophore synthesis. *Biochem Soc Trans* 30: 604–609
- Theil EC (2003) Ferritin: at the crossroads of iron and oxygen metabolism. *J Nutr* 133: 1549S–1553S
- Theil EC (2004) Iron, ferritin and nutrition. *Annu Rev Nutr* 24: 327–343
- Thipyapong P, Joel DM and Steffens JC (1997) Differential expression and turnover of the tomato polyphenol oxidase gene family during vegetative and reproductive development. *Plant Physiol* 113: 707–718
- Thomine S, Wang R, Ward JM, Crawford NM and Schroeder JI (2000) Cadmium and iron transport by members of a plant metal transporter family in *Arabidopsis* with homology to *Nramp* genes. *Proc Nat Acad Sci USA* 97: 4991–4996
- Thomine S, Lelievre F, Debarbieux E, Schroeder JI and Barbier-Brygoo H (2003) AtNRAMP3, a multispecific vacuolar metal transporter involved in plant responses to iron deficiency. *Plant J* 34: 685–695
- Thygesen PW, Dry IB and Robinson SP (1995) Polyphenol oxidase in potato. A multigene family that exhibits differential expression patterns. *Plant Physiol* 109: 525–531
- Tian L and DellaPenna D (2004) Progress in understanding the origin and functions of carotenoid hydroxylases in plants. *Arch Biochem Biophys* 430: 22–29
- Tottey S, Rich PR, Rondet SA and Robinson NJ (2001) Two Menkes-type ATPases supply copper for photosynthesis in *Synechocystis* PCC 6803. *J Biol Chem* 276: 19999–20004
- Tottey S, Rondet SA, Borrelly GP, Robinson PJ, Rich PR and Robinson NJ (2002) A copper metallochaperone for photosynthesis and respiration reveals metal-specific targets, interaction with an importer, and alternative sites for copper acquisition. *J Biol Chem* 277: 5490–5497
- Tottey S, Block MA, Allen M, Westergren T, Albriex C, Scheller HV, Merchant S and Jensen PE (2003) *Arabidopsis* CHL27, located in both envelope and thylakoid membranes, is required for the synthesis of protochlorophyllide. *Proc Nat Acad Sci USA* 100: 16119–16124
- Touraine B, Boutin JP, Marion-Poll A, Briat JF, Peltier G and Lobréaux S (2004) Nfu2: a scaffold protein required for [4Fe-4S] and ferredoxin iron-sulphur cluster assembly in *Arabidopsis* chloroplasts. *Plant J* 40: 101–111
- van der Mark F, de Lange T and Bienfait HF (1981) The role of ferritin in developing primary bean leaves under various light conditions. *Planta* 153: 338–342

- Van Ho A, Ward DM and Kaplan J (2002) Transition metal transport in yeast. *Annu Rev Microbiol* 56: 237–261
- Van Wuytswinkel O, Savino G and Briat JF (1995) Purification and characterization of recombinant pea-seed ferritins expressed in *Escherichia coli*: influence of N-terminus deletions on protein solubility and core formation *in vitro*. *Biochem J* 305: 253–261
- Van Wuytswinkel O, Vansuyt G, Grignon N, Fourcroy P and Briat JF (1998) Iron homeostasis alteration in transgenic tobacco overexpressing ferritin. *Plant J* 17: 93–97
- Waldo GS, Wright E, Whang ZH, Briat JF, Theil EC and Sayers DE (1995) Formation of the ferritin iron mineral occurs in plastids. *Plant Physiol* 109: 797–802
- Wang T, Shen G, Balasubramanian R, McIntosh L, Bryant DA and Golbeck JH (2004) The *sufR* gene (sll0088 in *Synechocystis* sp. strain PCC 6803) functions as a repressor of the *sufBCDS* operon in iron-sulfur cluster biogenesis in cyanobacteria. *J Bacteriol* 186: 956–967
- Wardrop AJ, Wicks RE and Entsch B (1999) Occurrence and expression of members of the ferritin gene family in cowpeas. *Biochem J* 337: 523–530
- Watanabe N, Che FS, Iwano M, Takayama S, Yoshida S and Isogai A (2001) Dual targeting of spinach protoporphyrinogen oxidase II to mitochondria and chloroplasts by alternative use of two in-frame initiation codons. *J Biol Chem* 276: 20474–20481
- Williams LE, Pittman JK and Hall JL (2000) Emerging mechanisms for heavy metal transport in plants. *Biochim Biophys Acta* 1465: 104–126
- Willows RD, Mayer SM, Foulk MS, DeLong A, Hanson K, Chory J and Beale SI (2000) Phytobilin biosynthesis: the *Synechocystis* sp. PCC 6803 heme oxygenase-encoding *ho1* gene complements a phytochrome-deficient *Arabidopsis thaliana* *hy1* mutant. *Plant Mol Biol* 43: 113–120
- Wood PM (1978) Interchangeable copper and iron proteins in algal photosynthesis. Studies on plastocyanin and cytochrome *c*-552 in *Chlamydomonas*. *Eur J Biochem* 87: 9–19
- Xu XM and Møller SG (2004) AtNAP7 is a plastidic SufC-like ATP-binding cassette/ATPase essential for *Arabidopsis* embryogenesis. *Proc Natl Acad Sci USA* 101: 9143–9148
- Yabe T, Morimoto K, Kikuchi S, Nishio K, Terashima I and Nakai M (2004) The *Arabidopsis* chloroplastic NifU-like protein CnfU, which can act as an iron-sulfur cluster scaffold protein, is required for biogenesis of ferredoxin and photosystem I. *Plant Cell* 16: 993–1007
- Ye H, Garifullina GF, Abdel-Ghany SE, Zhang L, Pilon-Smits EAH and Pilon M (2004) The chloroplast NifS-like protein of *Arabidopsis thaliana* is required for iron-sulfur cluster formation in ferredoxin. *Planta* 220: 602–608
- Yuvaniyama P, Agar JN, Cash VL, Johnson MK and Dean DR (2000) NifS-directed assembly of a transient [2Fe-2S] cluster within the NifU protein. *Proc Natl Acad Sci USA* 97: 599–604
- Zancani M, Peresson C, Biroccio A, Federici G, Urbani A, Murgia I, Soave C, Micali F, Vianello A and Macri F (2004) Evidence for the presence of ferritin in plant mitochondria. *Eur J Biochem* 271: 3657–3664
- Zheng L, White RH, Cash VL, Jack RF and Dean DR (1993) Cysteine desulfurase activity indicates a role for NIFS in metallocluster biosynthesis. *Proc Natl Acad Sci USA* 90: 2754–2758