


# Synthesis of $\alpha$ -Diketones from Alkylaryl- and Diarylalkynes Using Mercuric Salts

Michael E. Jung\* and Gang Deng

Department of Chemistry and Biochemistry, University of California, Los Angeles, 405 Hilgard Avenue, Los Angeles, California 90095, United States

**S** Supporting Information

**ABSTRACT:** Both alkylarylalkynes and diarylalkynes **1** are converted into the  $\alpha$ -diketones **2** in good yield by the use of mercuric salts, e.g., mercuric nitrate hydrate or mercuric triflate, in the presence of water. Other mercuric salts, e.g., sulfate, chloride, acetate, or trifluoroacetate, do not provide the diketone product. A possible mechanism is proposed.




The oxidation of alkylarylalkynes and diarylalkynes **1** to furnish the corresponding  $\alpha$ -diketones **2** is well-known in organic chemistry. A very large number of oxidants have been used for this process. For example,  $\text{KMnO}_4$  has been employed often for this transformation<sup>1</sup> as has  $\text{RuO}_4$  (often generated in situ or immobilized).<sup>2</sup> There are also several reports of the use of various DMSO-based oxidations, usually with an added electrophile<sup>3</sup> or in the presence of a palladium catalyst<sup>4</sup> for the formation of **2** from **1**. Finally, a large variety of other metal-based<sup>5</sup> and nonmetal-based<sup>6</sup> oxidations have been reported. For a project involving the synthesis of androgen receptor antagonists, we had need of a good method for converting alkylarylalkynes into  $\alpha$ -diketones. We report here that methodology and its application to the conversion of several disubstituted alkynes **1** to the corresponding  $\alpha$ -diketones **2**.

We hoped if it might be possible to intercept the well-known mechanism<sup>7</sup> for mercuric-catalyzed hydration of an alkyne (Scheme 1) by reaction of the  $\alpha$ -mercurioketone intermediate **D** with another equivalent of the mercuric salt.

We examined this process using 1,2-diphenylacetylene **1a** as the substrate. Thus treatment of **1a** with 2 equiv of mercuric

nitrate hydrate at 22 °C in aqueous THF for 12 h gave the expected  $\alpha$ -diketone, benzil **2a**, in 52% yield (Scheme 2). We


**Scheme 2. Oxidation of 1a To Give 2a**



carried out several experiments on this test reaction to find the best set of conditions. Carrying out the reaction under air or under argon gave the same results, so oxygen is not required for the process. The use of anhydrous THF with an added equivalent of water afforded good yields. Other solvents worked well, e.g., methanol, DME, dioxane, acetonitrile, acetone, acetic acid, DMSO, and especially DMF. The best yields were obtained in methanol with 1 equiv of added water (22 °C, 20 min, 84%) and in DMF (22 °C, 24 h, 90%). The use of other mercuric salts, e.g., sulfate, chloride, acetate, and trifluoroacetate, did not give any **2a**. However, the use of mercuric triflate,  $\text{Hg}(\text{OTf})_2$ , also produced good yields of the  $\alpha$ -diketone **2a**.

We then applied this method to the synthesis of a wide variety of  $\alpha$ -diketones using the following set of conditions, namely treatment of the alkyne **1** with 2 equiv of mercuric nitrate hydrate in DMF in air at 22 °C. The results are shown in Table 1. Arylalkynes generally gave good yields of the expected  $\alpha$ -diketone products, e.g., **2b,d-f,m,n**, with yields ranging from 47 to 82%. In addition, most of the diarylacetylenes gave quite good yields of product. The presence of halogens (**1c,i,j,n**) did not hinder the oxidation nor did the more oxidizable or labile functionalities, such as phenols (**1l** and **1m**), esters (**1k** and **1o**), amines (**1e**, **1p**, and **1q**), or nitriles (**1f**), all of which gave reasonable yields of the desired products **2**. Substrates with electron-donating substituents (methoxy, **2g**, or methyl, **2h**) gave good yields, while the substrate having a 4-trifluoromethyl

**Scheme 1. Mercuric Nitrate Catalyzed Hydration of an Alkyne**



Received: February 26, 2014

Published: April 1, 2014

Table 1. Oxidation of Alkynes with Mercuric Nitrate Hydrate

entry	Ar	R	time	product	byproduct	yield %
1	Ph	Ph	24 h	<b>2a</b>	none	90
2	Ph	Me	1 h	<b>2b</b>	none	78
3	3-FC <sub>6</sub> H <sub>4</sub>	Ph	2 h	<b>2c</b>	none	91
4	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	Me	20 min	<b>2d</b>	none	47
5	4-NH <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Bu	20 min	<b>2e</b>	none	79
6	4-(NCC(Me) <sub>2</sub> NH)C <sub>6</sub> H <sub>4</sub>	Bu	20 min	<b>2f</b>	none	67
7	4-MeOC <sub>6</sub> H <sub>4</sub>	Ph	2 h	<b>2g</b>	none	76
8	4-MeC <sub>6</sub> H <sub>4</sub>	Ph	2 h	<b>2h</b>	none	69
9	2-ClC <sub>6</sub> H <sub>4</sub>	Pr	1 h	<b>2i</b>	none	57
10	3-FC <sub>6</sub> H <sub>4</sub>	Pr	1 h	<b>2j</b>	none	78
11	2-(MeOOC)C <sub>6</sub> H <sub>4</sub>	Ph	1 h	<b>2k</b>	none	66
12	4-HOC <sub>6</sub> H <sub>4</sub>	Ph	2 h	<b>2l</b>	none	81
13	4-HOC <sub>6</sub> H <sub>4</sub>	Pr	15 min	<b>2m</b>	none	64
14	4-ClC <sub>6</sub> H <sub>4</sub>	Pr	30 min	<b>2n</b>	none	82
15	4-(AcO)C <sub>6</sub> H <sub>4</sub>	Ph	2 h	<b>2o</b>	none	56
16	4-(Bn <sub>2</sub> N)C <sub>6</sub> H <sub>4</sub>	Ph	2 h	<b>2p</b>	none	86
17	4-(BnNH)C <sub>6</sub> H <sub>4</sub>	Ph	2 h	<b>2q</b>	none	51
18			12 h		none	69
19	2-(NO <sub>2</sub> )C <sub>6</sub> H <sub>4</sub> <b>1s</b>	Ph	48 h (50 °C)			33/54
20	2-(NO <sub>2</sub> )C <sub>6</sub> H <sub>4</sub> <b>1t</b>	Pr	2 h			0/61%
21		Ph	1 h			69/20
22			20 min			49/38
23			1 h			32/37


group **1d** gave the lowest yield among the successful substrates, perhaps due to lower electron density in the alkyne. The 1,4-di(propynyl)benzene **1r** afforded the bis( $\alpha$ -diketone) **2r** in good yield. A few substrates did not work well in this reaction, giving mixtures of products. Thus, the 2-nitrophenyl substrate **1s** gave the expected  $\alpha$ -diketone **2s** in 33% yield along with the expected 3-benzoylanthranil **3s** in 54% yield. The analogous 2-nitrophenyl substrate with a propyl group on the end of the alkyne, **1t**, gave none of the  $\alpha$ -diketone and only the anthranil

**3t** in 61% yield. This cyclization of a 1-(2-nitrophenyl)alkynes such as **1s** and **1t** is well-known<sup>8</sup> and is usually carried out by treatment with either transition metals or Lewis acids. We also attempted the oxidation on two different heterocyclic alkyne substrates. Thus, the protected 4-pentynyl imidazole **1u** gave a relatively good yield of the desired  $\alpha$ -diketone **2u** (69%) accompanied by the simple hydration product **3u** in 20% yield. The 3-pentynylpyridine **1v** likewise gave the desired  $\alpha$ -diketone **2v** as the major product (49%) along with the opposite

hydration product **3v** in 38% yield.<sup>9</sup> Finally the symmetrical (2-nitrophenyl)acetylene **1w** did not give any of the desired  $\alpha$ -diketone product **2w** but rather the two byproducts, the simple hydration product **3w** in 32% yield and the anthranil **3w'** in 37% yield.<sup>10</sup>

We believe that the mechanism involves the steps shown in Scheme 3, namely attack of the alkyne **1** on the mercuric nitrate

**Scheme 3. Proposed Mechanism of Oxidation with Mercuric Nitrate**




to give the cyclic mercuronium ion **A**, which is then attacked by water to give **B**, which loses a proton to give the  $\alpha$ -mercurio enol **C**. Tautomerization would then give the  $\alpha$ -mercurio ketone **D**. Up to this point, this is the same mechanism as for the simple hydration of the alkyne (as shown in Scheme 1). The key step is the attack of nitrate on the  $\alpha$ -mercurio ketone **D**, with activation by the second equivalent of mercuric nitrate, to generate the  $\alpha$ -nitrato ketone **H** and mercurous nitrate. The final step is the reductive elimination of nitrous acid from **H** to give the observed  $\alpha$ -diketone product **2**.

Perhaps the most unusual step in this proposed mechanism is the conversion of the  $\alpha$ -mercurio ketone **D** to the  $\alpha$ -nitrato ketone **H**, but this step has precedent in the literature since a similar conversion of 2-methoxy-1,2-diphenylethyl mercuric nitrate to the 2-methoxy-1,2-diphenylethyl nitrate is known.<sup>11</sup> There is also good precedent for the final step, since the conversion of  $\alpha$ -nitrato ketones to  $\alpha$ -diketones is well-known.<sup>12</sup>

This oxidation also proceeds, although less well, with 2 equiv of anhydrous mercuric triflate and 2 equiv of water in THF. We propose a very similar mechanism for the formation of the  $\alpha$ -diketone **2** in this reaction (Scheme 4), namely the attack of triflate ion on the corresponding  $\alpha$ -mercurio ketone **D'** activated by mercuric triflate to give the  $\alpha$ -sulfonyloxy ketone

**Scheme 4. Proposed Mechanism of Oxidation with Mercuric Triflate**



**H'**. Elimination of trifluoromethanesulfinate from **H'** would then give the  $\alpha$ -diketone **2**. This last step, the elimination of sulfinate to give ketones, is well precedented in the literature.<sup>13</sup>

In summary, we have developed a new method for the oxidation of alkylaryllalkynes and diarylalkynes **1** to give  $\alpha$ -diketones **2** with mercuric salts. The reaction is limited to salts that can undergo facile subsequent elimination, namely nitrates and triflates. The use of this process in synthesis is underway and will be reported in due course.

## ■ ASSOCIATED CONTENT

### Supporting Information

Experimental procedures and proton and carbon NMR for all new compounds and those prepared by routes different from those in the literature. This material is free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [jung@chem.ucla.edu](mailto:jung@chem.ucla.edu).

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We thank the National Institutes of Health for support of this research. We also acknowledge a SPOR grant in Prostate Cancer for support. This material is based upon work supported by the National Science Foundation under equipment grant no. CHE-1048804. The LC-MS used in this project was supported by grant no. S10RR025631 from the National Center for Research Resources. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center for Research Resources or the National Institutes of Health. The NMR spectrometers were supported by the National Science Foundation under equipment grant no. CHE-1048804.

## ■ REFERENCES

- (1) (a) Lee, D. G.; Chang, V. S. *Synthesis* **1978**, 462. (b) Srinivasan, N. S.; Lee, D. G. *J. Org. Chem.* **1979**, *44*, 1574. (c) Lee, D. G.; Chang, V. S. *J. Org. Chem.* **1979**, *44*, 2726. For more recent examples, see: (d) Nowak, P.; Cole, D. C.; Aulabaugh, A.; Bard, J.; Chopra, R.; Cowling, R.; Fan, K. Y.; Hu, B.; Jacobsen, S.; Jani, M.; Jin, G.; Lo, M.-C.; Malamas, M. S.; Manas, E. S.; Narasimhan, R.; Reinhart, P.; Robichaud, A. J.; Stock, J. R.; Subrath, J.; Svenson, K.; Turner, J.; Wagner, E.; Zhou, P.; Ellingboe, J. W. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 632. (e) Mutoh, K.; Shima, K.; Yamaguchi, T.; Kobayashi, M.; Abe, J. *Org. Lett.* **2013**, *15*, 2938. (f) Trosien, S.; Waldvogel, S. R. *Org. Lett.* **2012**, *14*, 2976. (g) Reddy, M. A.; Thomas, A.; Mallesham, G.; Sridhar, B.; Rao, V. J.; Bhanuprakash, K. *Tetrahedron Lett.* **2011**, *52*, 6942.
- (2) (a) Gopal, H.; Gordon, A. J. *Tetrahedron Lett.* **1971**, *12*, 2941. (b) Mueller, P.; Godoy, J. *Helv. Chim. Acta* **1981**, *64*, 2531. (c) Carling, R. W.; Holmes, A. B. *Tetrahedron Lett.* **1986**, *27*, 6133. For more recent examples, see: (d) Bensulong, S.; Boonsombat, J.; Ruchirawat, S. *Tetrahedron* **2013**, *69*, 9335. (e) Xu, Y.; Wan, X. *Tetrahedron Lett.* **2013**, *54*, 642. (f) Ren, W.; Liu, J.; Chen, L.; Wan, X. *Adv. Synth. Catal.* **2010**, *352*, 1424. (g) Magnus, P.; Stent, M. A. H. *Org. Lett.* **2005**, *7*, 3853. (h) Kobayashi, S.; Miyamura, H.; Akiyama, R.; Ishida, T. *J. Am. Chem. Soc.* **2005**, *127*, 9251.
- (3) (a) Wolfe, S.; Pilgrim, W. R.; Garrard, T. F.; Chamberlain, P. C. *J. Chem.* **1971**, *49*, 1099. (b) Yusybov, M. S.; Filimonov, V. D. *Synthesis* **1991**, 131. For more recent examples, see: (c) Griebenow, N.; Meyer, T. *Synlett* **2010**, 2639. (d) Jones, D. J.; Purushothaman, B.; Ji, S.; Holmes, A. B.; Wong, W. W. H. *Chem. Commun.* **2012**, *48*, 8066.

(e) Giraud, A.; Provot, O.; Peyrat, J.-F.; Alami, M.; Brion, J.-D. *Tetrahedron* **2008**, *64*, 4287. (f) Niu, M.; Fu, H.; Jiang, Y.; Zhao, Y. *Synthesis* **2008**, 2879. (g) Wan, Z.; Jones, C. D.; Mitchell, D.; Pu, J. Y.; Zhang, T. Y. *J. Org. Chem.* **2006**, *71*, 826.

(4) (a) Yusubov, M. S.; Filimonov, V. D.; Vasilyeva, V. P.; Chi, K.-W. *Synthesis* **1995**, 1234. (b) Mori, S.; Takubo, M.; Yanase, T.; Maegawa, T.; Monguchi, Y.; Sajiki, H. *Adv. Synth. Catal.* **2010**, *352*, 1630. (c) Saga, Y.; Motoki, R.; Makino, S.; Shimizu, Y.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2010**, *132*, 7905. (d) Gao, A.; Yang, F.; Li, J.; Wu, Y. *Tetrahedron* **2012**, *68*, 4950. (e) Sawama, Y.; Takubo, M.; Mori, S.; Monguchi, Y.; Sajiki, H. *Eur. J. Org. Chem.* **2011**, 3361.

(5) (a) Ballistreri, F. P.; Failla, S.; Spina, E.; Tomaselli, G. A. *J. Org. Chem.* **1989**, *54*, 947. (b) Zhu, Z.; Espenson, J. H. *J. Org. Chem.* **1995**, *60*, 7728. (c) Ren, W.; Xia, Y.; Ji, S.-J.; Zhang, Y.; Wan, X.; Zhao, J. *Org. Lett.* **2009**, *11*, 1841. (d) Enthaler, S. *ChemCatChem* **2011**, *3*, 1929. (e) Xu, C.-F.; Xu, M.; Jia, Y.-X.; Li, C.-Y. *Org. Lett.* **2011**, *13*, 1556. (f) Liu, Y.; Chen, X.; Zhang, J.; Xu, Z. *Synlett* **2013**, *24*, 1371.

(6) (a) Clayton, M. D.; Marcinow, Z.; Rabideau, P. W. *Tetrahedron Lett.* **1998**, *39*, 9127. (b) Rogatchov, V. O.; Filimonov, V. D.; Yusubov, M. S. *Synthesis* **2001**, 1001. (c) Chu, J.-H.; Chen, Y.-J.; Wu, M.-J. *Synthesis* **2009**, 2155. (d) Santoro, S.; Battistelli, B.; Gjoka, B.; Si, C.-w. S.; Testaferri, L.; Tiecco, M.; Santi, C. *Synlett* **2010**, 1402. (e) Tingoli, M.; Mazzella, M.; Panunzi, B.; Tuzi, A. *Eur. J. Org. Chem.* **2011**, 399.

(7) (a) Kutscheroff, M. *Chem. Ber.* **1881**, *14*, 1540. (b) Kutscheroff, M. *Chem. Ber.* **1884**, *17*, 13. (c) Kutscheroff, M. G. *Chem. Ber.* **1909**, *42*, 2759. For a good review, see: (d) Hintermann, L.; Labonne, A. *Synthesis* **2007**, 1121. For practical procedures, see: (e) Stacy, G. W.; Mikulec, R. A. *Org. Synth. Coll.* **1963**, *Vol. IV*, 13. (f) Palomo, C.; Oiárbide, M.; Aizpurua, J. M.; González, A.; Garcia, J. M.; Landa, C.; Odriozola, I.; Linden, A. *J. Org. Chem.* **1999**, *64*, 8193. (g) Corominas, A.; Montana, A. M. *Synth. Commun.* **2013**, *43*, 2062.

(8) (a) Asao, N.; Sato, K.; Yamamoto, Y. *Tetrahedron Lett.* **2003**, *44*, 5675. (b) Li, X.; Incarvito, C. D.; Vogel, T.; Crabtree, R. H. *Organometallics* **2005**, *24*, 3066. (c) Jadhav, A. M.; Bhunia, S.; Liao, H.-Y.; Liu, R.-S. *J. Am. Chem. Soc.* **2011**, *133*, 1769. (d) Ramana, C. V.; Patel, P.; Vanka, K.; Miao, B.; Degterev, A. *Eur. J. Org. Chem.* **2010**, 5955.

(9) It is likely that the protonated pyridine (or the mercuric complex at N) causes the normal regiochemistry of hydration to be reversed.

(10) This is a known product of the acid promoted rearrangement of 1,2-di(*o*-nitrophenyl)ethanol. Bakke, J. M. *Acta Chem. Scand. B* **1975**, *29*, 1063.

(11) Shearer, D. A.; Wright, G. F. *Can. J. Chem.* **1955**, *33*, 1002.

(12) (a) Emmons, W. D.; Freeman, J. P. *J. Am. Chem. Soc.* **1955**, *77*, 4415. (b) For a similar elimination to give aldehydes from benzyl nitrates, see: McKillop, A.; Ford, M. E. *Synth. Commun.* **1974**, 45.

(13) (a) Hoffman, R. V.; Wilson, A. L.; Kim, H.-O. *J. Org. Chem.* **1990**, *55*, 1267. (b) Jung, M. E.; Lam, P. S. Y.; Mansuri, M. M.; Speltz, L. M. *J. Org. Chem.* **1985**, *50*, 1087. (c) Creary, X. *Acc. Chem. Res.* **1985**, *18*, 3. (d) Maas, G.; Lorenz, W. *J. Org. Chem.* **1984**, *49*, 2273. (e) Creary, X. *J. Org. Chem.* **1980**, *45*, 2419.