

Development of Novel Mitochondrial Pyruvate Carrier Inhibitors to Treat Hair Loss

Xiaoguang Liu, Aimee A. Flores, Lisa Situ, Wen Gu, Hui Ding, Heather R. Christofk, William E. Lowry, and Michael E. Jung*

Cite This: *J. Med. Chem.* 2021, 64, 2046–2063


Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information



ABSTRACT: Herein, we report the synthesis and evaluation of novel analogues of UK-5099 both *in vitro* and *in vivo* for the development of mitochondrial pyruvate carrier (MPC) inhibitors to treat hair loss. A comprehensive understanding of the structure–activity relationship was obtained by varying four positions of the hit compound, namely, the alkyl group on the N1 position, substituents on the indole core, various aromatic and heteroaromatic core structures, and various Michael acceptors. The major discovery was that the inhibitors with a 3,5-bis(trifluoromethyl)benzyl group at the N1 position were shown to have much better activity than JXL001 (UK-5099) to increase cellular lactate production. Additionally, analogue JXL069, possessing a 7-azaindole heterocycle, was also shown to have significant MPC inhibition activity, which further increases the chemical space for drug design. Finally, more than 10 analogues were tested on shaved mice by topical treatment and promoted obvious hair growth on mice.

INTRODUCTION

Hair loss, also known as alopecia or baldness, is a very common health problem. Fifty percent of males and 25% of females by age 50 suffer pattern hair loss.¹ Common types of hair loss include male-pattern hair loss, female-pattern hair loss, alopecia areata, and a thinning of hair known as telogen effluvium.¹ Minoxidil (Rogaine) and finasteride (Propecia) are the two Food and Drug Administration (FDA)-approved medicines to treat hair loss but only 30–40% of patients experience hair growth with the requirement of repeated treatment, and serious side effects have been reported.² Surgical procedures to treat baldness are expensive and can be painful. Therefore, a better therapy for hair loss is highly desired.

Hair follicle growth occurs in cycles. Each cycle consists of a long growing phase (anagen), a short transitional phase (catagen), and a short resting phase (telogen).³ The ability of the hair follicles to maintain this cycle depends on the presence of hair follicle stem cells (HFSCs).⁴ The HFSCs are typically inactive but can quickly “wake up” and actively divide when a new hair growth cycle is initiated.^{5,6} When HFSCs fail to

activate, hair loss occurs. Our collaborators, Lowry, Christofk, and co-workers, discovered that the activation of HFSCs can be achieved by stimulating the activity of lactate dehydrogenase (LDH), the enzyme catalyzing the reduction of pyruvate to lactate.⁷ Genetic deletion of mitochondrial pyruvate carrier (MPC) or pharmacological inhibition of MPC via treatment with UK-5099 resulted in increased LDH activity in HFSCs.⁷ When the MPC is knocked down or pharmacologically inhibited, pyruvate in the cytoplasm cannot enter the mitochondria and instead is converted to other metabolites, such as lactate by lactate dehydrogenase (LDH), which results in the increase of LDH activity and the activation of HFSCs. Overall, our collaborators discovered a novel mechanism to

Received: September 8, 2020

Published: February 3, 2021



promote hair growth and MPC has been proven to be a promising therapeutic target for the treatment of hair loss.


MPC, located on the inner mitochondrial membrane, is a transporter protein that transports pyruvate from the cytoplasm to the mitochondria. Although great efforts have been taken to study MPC, its molecular structure is still not clear.^{8,9} The MPC protein family consists of three members, MPC1, MPC2, and MPC3. The MPC1–MPC2 complex, an oligomeric structure of about 150 kD, is an essential component of the mitochondrial pyruvate carrier in yeast, flies, and mammals.⁹ UK-5099 is a known covalent inhibitor of MPC, which reversibly reacts with a cysteine on the protein with a K_i value of 0.05 μM .¹⁰ UK-5099 has been reported to promote lactate production rate as a result of MPC inhibition in various settings.^{11,12} Our collaborators, Lowry, Christofk, and co-workers, demonstrated that the topical treatment of mice in the telogen phase with UK-5099 (20 μM) led to a robust acceleration of the hair cycle.⁷ Therefore, UK-5099 is a good hit compound for developing novel MPC inhibitors as a treatment of hair loss.

Herein, we describe an extensive structure–activity relationship (SAR) study of a series of (*E*)-2-cyano-3-(1*H*-indol-3-yl)acrylic acids and their structural analogues. We have developed a general synthetic route to these compounds that allows one to prepare many analogues rapidly. All of these new compounds have been tested for their ability to promote cellular lactate production *in vitro*. The compounds that enhanced cellular lactate production rate were then tested on mice topically for their ability to accelerate hair growth. We have identified more than 10 compounds with potent MPC inhibitory activity and the ability to promote hair growth in mice.

RESULTS AND DISCUSSION

Chemistry. The synthesis of the analogues, which are labeled **JXL001**–**JXL096**, was carried out as shown in the various schemes. The synthesis of **JXL001** is shown in **Scheme 1**. Copper-mediated *N*-arylation of indole-3-carboxaldehyde **1**

Scheme 1. Synthesis of JXL001




with iodobenzene followed by Knoevenagel condensation with ethyl 2-cyanoacetate provided the ester version of UK-5099, **JXL004**. Finally, aqueous LiOH-mediated saponification delivered the desired compound **JXL001**. Following the same synthetic route with substituted indole-3-carboxaldehydes and substituted iodobenzenes, various analogues were synthesized rapidly (**Table 1**).

Analogue **JXL020** was synthesized by the synthetic route shown in **Scheme 2**. Benzylation of indole-3-carboxaldehyde **1**

Table 1. Analogues Synthesized by Following Scheme 1


analogue	R ¹	R ²	R ³
JXL001	H	OH	Ph
JXL002	H	OH	H
JXL003	H	OEt	H
JXL004	H	OEt	Ph
JXL005	6-Cl	OEt	Ph
JXL006	H	OEt	2-OMeC ₆ H ₄
JXL007	H	OEt	4-OMeC ₆ H ₄
JXL012	H	OH	4-OMeC ₆ H ₄
JXL013	6-Cl	OH	Ph
JXL014	H	OH	2-OMeC ₆ H ₄
JXL026	4-Cl	OH	Ph
JXL027	6-F	OH	Ph
JXL028	7-Cl	OH	Ph
JXL029	5-F	OH	Ph
JXL035	4-Cl	OH	Ph
JXL093	4-Cl	OEt	Ph

Scheme 2. Synthesis of JXL020



with 3,5-bis(trifluoromethyl)benzyl bromide generated compound **3** in good yield. Knoevenagel condensation with ethyl 2-cyanoacetate provided **JXL011**. Finally, aqueous LiOH-mediated saponification afforded analogue **JXL020**. When substituted indole-3-carboxaldehydes and various benzyl bromides were used, various analogues were synthesized rapidly by following the same synthetic route as for **JXL020**. The analogues are listed in **Table 2**. Based on the result of our SAR study, the analogues of **JXL020** with the 3,5-bis(trifluoromethyl)benzyl group at the N1 position were shown to have much better activity than **JXL001** in promoting cellular lactate production. A series of analogues with a 3,5-bis(trifluoromethyl)benzyl group at the N1 position were

Table 2. Analogues Synthesized by Following Scheme 2




analogue	R ¹	R ²	R ⁴
JXL008	H	OEt	4-F
JXL009	H	OEt	3,4-di-F
JXL010	H	OEt	3,5-di-F
JXL011	H	OEt	3,5-bis(CF ₃)
JXL015	H	OH	4-F
JXL016	H	OH	3,4-di-F
JXL017	H	OH	3,5-di-F
JXL018	H	OEt	H
JXL019	H	OH	H
JXL020	H	OH	3,5-bis(CF ₃)
JXL036	5-F	OH	3,4-di-F
JXL037	5-F	OH	3,5-di-F
JXL039	6-F	OH	3,4-di-F
JXL040	6-F	OH	3,5-di-F

synthesized by following a similar synthetic route as shown for JXL020 (Table 3).

The analogue JXL079 with a substituted homobenzyl group at the N1 position was synthesized as shown in Scheme 3a. The homobenzyl mesylate 7 was synthesized from the commercially available compound 3,5-bis(trifluoromethyl)benzyl bromide by nucleophilic addition of its derived

Table 3. Analogues with 3,5-Bis(trifluoromethyl)benzyl Group at the N1 Position



analogue	R ¹	R ²	analogue	R ¹	R ²
JXL038	5-F	OH	JXL062	6-Br	OH
JXL041	6-F	OH	JXL063	6-Cl	OH
JXL050	4-Cl	OH	JXL064	7-Cl	OH
JXL051	4-Br	OH	JXL065	4-Br	OEt
JXL052	4-F	OH	JXL066	4-F	OEt
JXL053	7-F	OH	JXL068	H	<i>Or</i> -Bu
JXL054	5-Cl	OH	JXL078	H	OMe
JXL055	4-CN	OH	JXL081	H	morpholine
JXL056	4-CO ₂ H	OH	JXL089	5-Cl	OEt
JXL057	4-OBn	OH	JXL090	4-CN	OEt
JXL058	6-OBn	OH	JXL091	4-CO ₂ Me	OEt
JXL059	7-OBn	OH	JXL092	4-CO ₂ Me	OH
JXL060	4-OMe	OH	JXL094	H	NH ₂
JXL061	5-Br	OH			

Grignard reagent to carbon dioxide followed by diisobutylaluminum hydride (DIBAL) reduction and mesylation. Nucleophilic substitution of compound 7 with indole 8 generated compound 9. Upon treatment with trifluoroacetic acid, the *tert*-butyl group was cleaved to form the desired analogue JXL079. The amide analogue JXL080 was synthesized by an *N,N'*-dicyclohexylcarbodiimide (DCC) coupling reaction of JXL003 and compound 6 (Scheme 3b).

In addition to analogues with the indole core, analogues with other aromatic and heteroaromatic cores were also synthesized. 4-Pyridinecarboxaldehyde and substituted benzaldehydes were used to synthesize analogues following a synthetic route similar to that used for JXL001 (Scheme 1). The analogues are listed in Table 4. Some analogues with nitrogen-containing heterocycles were also synthesized using a synthetic route similar to that used for JXL069 in Scheme 4. Benzoylation of 7-azaindole 4 with KOH as the base generated compound 5 with good yield and regioselectivity. The Vilsmeier–Haack reaction followed by Knoevenagel condensation with ethyl 2-cyanoacetate provided JXL082. Finally, aqueous LiOH-mediated saponification furnished the analogue JXL069. Analogues with a 4-azaindole, a 6-azaindole, and a substituted 7-azaindole core were successfully synthesized and are listed in Table 5. The Vilsmeier–Haack reaction failed to generate the 3-carboxyaldehydes of 4-azaindole and 6-azaindole, so the formylation was carried out using hexamethylenetetramine (HMTA) in acetic acid (Duff reaction).

The analogues with substituents at the 3-position of the acrylates, JXL083, JXL084, and JXL085, were synthesized, as shown in Scheme 5. Aldehyde 3 was oxidized by a Pinnick oxidation to the carboxylic acid 10, which was further converted to acyl chloride 11. Deprotonation of ethyl 2-cyanoacetate by lithium diisopropylamide (LDA) and nucleophilic addition to compound 11 generated analogue JXL083. Analogue JXL084 was synthesized by acetylation of JXL083 and JXL085 was synthesized by chlorination of JXL083.

Analogues with various Michael acceptors were synthesized by Knoevenagel condensations (Figure 1). The bis-nitrile JXL021 was synthesized by the condensation with malononitrile and JXL025 was synthesized by the condensation with diethyl malonate followed by LiOH-mediated monosaponification. JXL086 and JXL095 were generated by the condensation of aldehydes 3 and 12, respectively, with diethyl (cyanomethyl)phosphonate. JXL096 was synthesized from JXL086 by trimethylsilyl bromide (TMSBr)-mediated cleavage of the ethyl groups. Analogues with thiazolidinedione (TZD) structures were synthesized similarly as JXL024. Condensation of various aldehydes with either thiazolidine-2,4-dione or 2-iminothiazolidin-4-one in reflux acetic acid afforded the analogues listed in Figure 2.

Biological Evaluation: SAR Analysis. When the MPC is inhibited, pyruvate in the cytoplasm cannot enter the mitochondria and instead is converted to other metabolites, such as lactate by lactate dehydrogenase (LDH). Since our collaborators previously found that LDH activity correlated with hair follicle stem cell activation, and genetic deletion of LDH activity via LDH α knockout in hair follicle stem cells blocked their activation,⁷ we were particularly interested in using rerouting of pyruvate toward lactate production as a readout for MPC inhibition in this study. The use of cell-based lactate production as a readout of MPC inhibition was a practical first-level readout for screening new compounds, but

Scheme 3. Synthesis of JXL079 and JXL080




Table 4. Analogues with Benzene Core and JXL022

analogue	R ⁵	analogue	R ⁵
JXL030	2-F	JXL045	3-CF ₃ , 4-CF ₃
JXL032	4-F	JXL046	2-Cl, 3-CF ₃
JXL034	4-CF ₃	JXL047	3-CF ₃ , 4-F
JXL042	3-F, 4-Me	JXL048	H
JXL043	3-F, 4-F	JXL049	4-OH
JXL044	2-F, 4-F		

that it is certainly not synonymous with MPC inhibition. Our hit compound, UK-5099, is known to be a covalent inhibitor. To confirm its mechanism of action, the analogue of UK-5099 without the alkene double bond was synthesized and tested in the cell-based assay. The reduced UK-5099 showed no effect on lactate production rate, meaning that it did not inhibit MPC (data not shown). This result supports that UK-5099 is a covalent inhibitor.

A dose–response experiment of the commercial UK-5099 and in-house-prepared JXL001 with MCF10A cells was conducted to confirm their inhibition of MPC by measuring the cells' lactate production rate (data not shown). This result indicated that both samples promoted lactate production in a dose-dependent pattern. Because both samples provided the highest lactate production rate at 10 μ M and the same concentration of UK-5099 was used in the earlier *in vivo* study to detect an effect on hair growth,⁷ a concentration of 10 μ M was chosen for the evaluation of the activity of the analogues to

promote cellular lactate production. The data of the *in vitro* test is shown in Figure 3.

Here, we used the (*E*)-2-cyano-3-(1-phenyl-1H-indol-3-yl)acrylic acid (JXL001) as a reference compound and designed new compounds to vary the four most easily altered positions, namely, (1) the alkyl group at N1, (2) the substituents on the indole core, (3) various aromatic and heteroaromatic cores, and (4) various Michael acceptors. Our goal was to identify candidates for the treatment of hair loss with significant improvement in potency.

Analogues at the N1 Position (R3). We envisioned that modifying the *N*-phenyl group would be a good starting point for conducting an SAR by medicinal chemistry. Several *N*-aryl-substituted analogues are listed in Table 1. Analogue JXL002 without the phenyl group showed little inhibitory effect on MPC, implying that the phenyl group is important for UK-5099's biological activity. Similarly, compound JXL003 was worse than JXL004 in terms of stimulating lactate production. Analogues with *N*-2-methoxyphenyl and *N*-4-methoxyphenyl groups (JXL006 and JXL007, esters; JXL014 and JXL012, carboxylic acids) showed similar or slightly better activity than JXL001 to promote the cellular lactate production rate. Due to the harsh reaction conditions and low reaction yields for the *N*-arylation step, we switched our attention to analogues with *N*-benzyl groups.

We then investigated the *N*-benzyl analogues because the homologated analogues possess high structural similarity with JXL001. A series of analogues with different benzyl groups were synthesized and tested (Table 2). Although most of these analogues showed better activity in increasing lactate production rate, the analogue with the 3,5-bis-(trifluoromethyl)benzyl group, JXL020, was shown to have

Scheme 4. Synthesis of JXL069





Table 5. Analogues with Azaindole Cores



analogue	X	Y	Z	R ¹	R ²
JXL069	C	CH	N	H	OH
JXL072	N	CH	CH	N/A	OH
JXL073	C	N	CH	H	OH
JXL076	C	CH	N	Cl	OH
JXL077	C	CH	N	Br	OH
JXL082	C	CH	N	H	OEt
JXL087	C	CH	N	Cl	OEt
JXL088	C	CH	N	Br	OEt

the highest ability to inhibit MPC. JXL079, the homologue of JXL020, was shown to have lower potency than JXL020 in inhibition of MPC. Additionally, the amide JXL080 did not promote the cellular lactate production. Up to this point, the 3,5-bis(trifluoromethyl)benzyl group was the best moiety at the N1 position of the indole core.

Analogues with Substituents on the Indole Ring.

Taking JXL020 as the new hit compound, the influence of substituents on the indole core was explored (the analogues are listed in Table 3). Comparing the four analogues containing 4-F, 5-F, 6-F, and 7-F (JXL052, JXL038, JXL041, and JXL053) with JXL020, JXL052 (the 4-F analogue) and JXL053 (the 7-F analogue) showed better activity than JXL020 to increase the lactate production rate, while the other two analogues could hardly promote the lactate

production. For chloro-substituted analogues (JXL050, JXL054, JXL063, and JXL064), only the 4-Cl analogue (JXL050) showed better activity than JXL020 to increase the lactate production rate. Similarly, the analogue with 4-Br (JXL051) was better than JXL020, while analogues with 5-Br and 6-Br were worse than JXL020. Among the three benzyloxy-substituted indole analogues (JXL057, JXL058, and JXL059), only the 7-OBn analogue JXL059 showed better activity than JXL020 but the 4-OBn analogue failed to increase its biological activity in promoting lactate production rate, probably due to the steric effect of the large substituent. Based on the analysis, more analogues with small substituents at the 4-position of indole were synthesized and evaluated. The analogue with 4-CN (JXL055) was better than JXL020 and the analogue with 4-OMe (JXL060) was similar to JXL020 in terms of promoting lactate production. The analogue with 4-COOH (JXL056) showed much worse activity than JXL020 in inhibition of MPC, perhaps due to its high polarity that may make it hard to penetrate the cell membrane. Based on this data, it seems that large substituents on the indole ring are harmful to the inhibitory activity for MPC and small substituents at the four-position are beneficial for the desired activity.

Analogues with Aromatic and Heteroaromatic Cores Other than the Indole Core.

Analogues with 4-pyridine and many other substituted benzene cores (Table 4) were shown to have poorer activity than JXL001 in promoting cellular lactate production, implying that the indole core structure is important for the biological activity.

We further evaluated other aromatic cores to identify novel MPC inhibitors (Table 5). Analogues containing azaindole cores were synthesized according to Scheme 4. To our delight, the 7-azaindole analogue (JXL069) was much more potent than the 4-azaindole analogue (JXL072), the 6-azaindole analogue (JXL073), and the hit compound JXL020 with an indole core. The improved inhibitory activity may result from a new ligand–protein interaction or the electronic effect of the azaindole structure. The 4-Cl-7-azaindole and 4-Br-7-azaindole

Scheme 5. Synthesis of JXL083, JXL084, and JXL085





Figure 1. Analogues with various Michael acceptors.




Figure 2. Analogues containing the TZD group.

analogues (**JXL076** and **JXL077**) showed similar activity as **JXL020** in promoting lactate production rate. This new azaindole structure broadened the chemical space for further SAR study.

Analogues with Various Michael Acceptors. We envisioned that manipulating the Michael acceptor moiety might be important for the potency and selectivity of the

analogues. Modification of this group can change the molecule's lipophilicity and electrophilicity. To test this idea, analogues with different alkyl groups were synthesized (Table 3). The methyl ester (**JXL078**) and the *tert*-butyl ester analogues (**JXL068**) showed a dramatic loss of potency compared with the corresponding carboxylic acid analogue




Figure 3. *In vitro* evaluation of analogues of JXL001–JXL096. Human breast epithelial (MCF10A) cells were treated with 10 μ M analogue for 24–30 h, and the cellular lactate production rate was measured, relative to vehicle (dimethyl sulfoxide, DMSO) treatment and normalized to cell number. Data were compiled from sequential batches of experiments, each with their own DMSO controls and JXL001 and JXL020 treatments for comparison. Error bars indicate +1 s.d. Representative experiments are shown, with most analogues tested at least twice. Shading indicates adjusted *p*-values calculated by performing ordinary one-way analysis of variance (ANOVA) followed by Dunnett's post-test, comparing each analogue to DMSO.

(JXL020), while the ethyl ester (JXL011) showed similar potency as JXL020. Based on this discovery, many other ethyl ester analogues were tested. To our delight, most of them were shown to have similar potency to their corresponding carboxylic acid analogues, such as JXL065 and JXL051, JXL066 and JXL052, and JXL082 and JXL069.

The analogues with malononitrile (JXL021) and dicarboxylate (JXL025) were shown to have poorer activity than JXL001 in increasing the lactate production rate (Figure 1). Analogues containing the TZD groups in Figure 2 failed to show promising activity in promoting lactate production. Amide analogues (JXL081 and JXL094) showed very strong ability to promote lactate production; however, substantial cell death was observed (data was not shown). The indole core-based diethyl phosphonate analogue (JXL086; Figure 1) promoted lactate production more than JXL001. However, the corresponding phosphonic acid analogue (JXL096) failed to promote lactate production rate, perhaps due to its poor cell permeability. The 7-azaindole core-based diethyl phosphonate analogue (JXL095) showed even stronger potency than JXL086 in promoting lactate production. Overall, adjusting the Michael acceptors showed a dramatic effect on the biological activity.

IC₅₀ Determination for Several Analogues. The test of mitochondrial pyruvate uptake would directly confirm MPC inhibition. However, given the structural similarity between UK-5099 (which does not inhibit pyruvate carboxylase (PC) or pyruvate dehydrogenase (PDH) activity) and the new analogues, it is reasonable that decreased pyruvate-driven respiration is a result of MPC inhibition. Since the MPC enables pyruvate-fueled respiration, measuring how well the UK-5099 analogues block the oxygen consumption rate (OCR) can be an assay of MPC inhibition.¹³ The pyruvate-

fueled OCR assay we used on permeabilized cells for measuring IC₅₀ of our MPC inhibitors is the most direct functional assay available for measuring MPC inhibition. In this assay, the only substrate for mitochondrial respiration given to the permeabilized cells is pyruvate. The MPC inhibitors block the pyruvate-fueled OCRs in a dose-dependent manner. We were able to convince ourselves that these values were direct since our MPC inhibitors had no effect on succinate-fueled OCR, suggesting a direct effect on the MPC and not some other component of the electron transport chain. Analogues JXL001, JXL020, and JXL069 were tested for their ability to inhibit the pyruvate-fueled respiration of permeabilized cells, and IC₅₀ values were determined (Figure 4). JXL001 showed an IC₅₀ value of around 140 nM. JXL020 and JXL069 showed much lower IC₅₀ values (16.6 and 42.8 nM) than JXL001, which implies that they are more potent than JXL001 in inhibiting MPC.

***In Vivo* Evaluation of MPC Inhibitors.** To determine the efficacy of the compounds on hair growth, C57BL/6J mice were shaved at postnatal day 50 during the telogen phase of the hair cycle. Compounds were suspended in lotion and applied topically to the shaved back skin of the mice every other day for 2 weeks. As seen in Figure 5, all of the analogues that showed the ability to promote lactate production in the *in vitro* assay were also able to stimulate hair growth over the course of 2 weeks. JXL020, the analogue with 3,5-bis-(trifluoromethyl)benzyl group at the N1 position, strongly promoted hair growth. The four-substituted analogues, JXL035, JXL050, JXL051, and JXL052, were also shown to have a strong ability to accelerate hair growth on mice. This data showed that most of the MPC inhibitors active *in vitro* could promote hair growth in mice. We tried to compare the *in vivo* efficacy of the MPC inhibitors with different *in vitro*




Figure 4. IC₅₀ measurement of JXL001, JXL020, and JXL069. MCF10A cells were treated with analogues to measure oxygen consumption rate (OCR) under different conditions. State 3 is the measurement of respiration in permeabilized cells in the presence of pyruvate, malate, and ADP. IC₅₀ values are indicated in bold. Error bars denote ±1 standard error of mean (SEM; *n* = 2).

potencies on the MPC inhibition. However, the hair growth assay is not very quantitative with respect to timing. All of the MPC inhibitors perform better than the controls, but it is difficult to run enough animals together to see statistically significant data to distinguish between the activities of the various MPC inhibitors.

CONCLUSIONS

This study reports the synthesis and evaluation of novel analogues of UK-5099 both *in vitro* and *in vivo* for the development of MPC inhibitors to treat hair loss. We modified

four sites around the core structure and tested the activity of those analogues in promoting cellular lactate production to establish a structure–activity relationship for this system. The substituent at the N1 position is crucial for the inhibition of MPC. After screening various aryl and substituted benzyl groups, the analogue containing a 3,5-bis(trifluoromethyl)-benzyl group at the N1 position, JXL020, was shown to have much better ability than JXL001 to promote cellular lactate production *in vitro* (IC₅₀ = 16.6 nM) and accelerate hair growth on mice. Modification of substituents on the indole core structure revealed that small substituents, such as F, Cl, Br, and CN, at the C4 position were beneficial for the biological activity. In addition, analogues with different aromatic and heteroaromatic core structures were synthesized and tested. Analogue JXL069, having a 7-azaindole heterocycle, was also shown to have significantly better activity than the analogues containing a 4-azaindole or a 6-azaindole core structure. This discovery further increases the chemical space for the development of MPC inhibitors. Furthermore, different types of Michael acceptors were synthesized and tested for their ability to increase lactate production rate. Analogues with the ethyl ester functional group were shown to have similar activity to inhibit MPC as their corresponding carboxylic acid analogues, such as ester JXL011 and acid JXL020. This finding might provide more opportunities in drug formulation since the esters have different solubilities than the carboxylic acids. Analogues containing amides (JXL081 and JXL094) and diethyl phosphonates (JXL086 and JXL095) showed potent MPC inhibition. Finally, more than 10 analogues were tested on mice topically to promote hair growth. The analogues containing the 3,5-bis(trifluoromethyl)benzyl group at the N1 position (JXL020, JXL050, JXL051, and JXL052) stimulated hair growth dramatically. With the novel and potent MPC inhibitors in hand, the pharmacokinetic and safety tests of these compounds are ongoing with the final goal of developing drugs to treat hair growth.




Figure 5. Evaluation of analogues *in vivo*. On each row, the mice were treated at the same time with the corresponding MPC inhibitors. Animals treated topically with the analogue (20 μ M) show pigmentation and hair growth, indicative of the entry into anagen, after 8 days of treatment. Full anagen, indicated by a full coat of hair, is achieved after 14 days of treatment. Mice treated topically with vehicle control do not show pigmentation nor hair growth even after 12 days of treatment.

EXPERIMENTAL SECTION

General. Toluene was distilled from sodium under an argon atmosphere. Dichloromethane was distilled from calcium hydride under an argon atmosphere. All other solvents or reagents were purified according to literature procedures. ^1H NMR spectra were recorded on Bruker spectrometers at 500 MHz and are reported relative to deuterated solvent signals. Data for ^1H NMR spectra are reported as follows: chemical shift (δ ppm), multiplicity, coupling constant (Hz), and integration. Splitting patterns are designated as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; and br, broad. ^{13}C NMR spectra were recorded on Bruker spectrometers at 100 MHz. Data for ^{13}C NMR spectra are reported as follows: chemical shift (δ ppm), multiplicity, and coupling constant (Hz). Splitting patterns are designated the same way as in ^1H NMR. High-resolution mass spectrometry (HRMS) was taken on a Waters LCT Premier with ACQUITY UPLC with an autosampler. Mass spectra were recorded from 70 to 2000 Da. The purity of compounds was analyzed with an Agilent 1200 HPLC system equipped with a G1312A binary pump, a G1314A autosampler, a G1314A VWD, and an Agilent SB-Aq C18 column (1.8 μm , 4.6 \times 50 mm 2). Mass spectra were recorded using an Agilent 6130 LC/MS system equipped with an ESI source. Method: solution A: H_2O w/0.1% formic acid and solution B: ACN w/0.1% formic acid; increase of B from 10 to 99%, 0.5–2 min; 99%, 2–17 min; flow rate 1.0 mL/min; UV 254 nm. The purity of all final compounds was determined to be >95% by analytical high-performance liquid chromatography (HPLC) analysis except the compounds **JXL024**, **JXL025**, and **JXL056** with around 90% purity. All solvents were liquid chromatography-tandem mass spectrometry (LC–MS/MS) grade and purchased from Fisher Scientific.

Experimental Details for the Synthesis of JXL001 and Related Analogues. To the solution of indole-3-carboxaldehyde (2.8 mmol, 411 mg) in dry dimethylformamide (DMF; 6 mL) were added Cu_2O (0.3 equiv, 0.84 mmol, 120 mg), K_2CO_3 (2.0 equiv, 5.6 mmol, 774 mg), and iodobenzene (2.0 equiv, 5.6 mmol, 624 μL) sequentially. The reaction was stirred and refluxed for 24 h, at which point thin-layer chromatography (TLC) indicated that the reaction was completed. After it was cooled to 21 $^\circ\text{C}$, the reaction mixture was filtrated through a celite pad eluting with ethyl acetate. The filtrate was washed with saturated NaCl solution, and the organic phase was dried by sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes/ethyl acetate = 8:1) to provide the desired product. Yield: 89%, 550.7 mg.

To the solution of 1-phenyl-indole-3-carbaldehyde (1 mmol, 221 mg) in ethanol (1 mL) were added ethyl 2-cyanoacetate (1.3 equiv, 1.3 mmol, 140 μL) and L-proline (40 mol %, 0.4 mmol, 58 mg). The reaction was stirred at 21 $^\circ\text{C}$ for 12 h and a yellow solid precipitated gradually. After completion of the reaction, ice-cold water (2 mL) was added into the reaction vial. The solid was separated by Buchner funnel filtration and washed with water (2 mL \times 3) and dried to afford the desired product. Yield: 95%, 300 mg.

To the solution of (*E*)-ethyl 2-cyano-3-(1-phenyl-1*H*-indol-3-yl)acrylate (0.32 mmol, 100 mg) in tetrahydrofuran (THF; 2 mL) was added 0.5 N LiOH solution (3 equiv, 0.6 mmol, 1.2 mL). The reaction mixture was stirred at 21 $^\circ\text{C}$ for 1 h. After reaction completion shown by TLC, THF was evaporated. Concentrated HCl was added dropwise to acidify the reaction mixture until the pH was lower than 1; meanwhile, a yellow solid precipitated. Ice-cold water (5 mL) was added to the reaction mixture, and the solid was separated by Buchner funnel filtration and washed with water (5 mL \times 3). After drying by vacuum, the solid was washed with 2 mL of solvent mixture (hexanes/ethyl acetate = 5:1) 5–10 times and monitored by TLC until nonpolar impurities disappear (the nonpolar compound was the retro-aldol condensation product, which can be recovered from the filtrate). Finally, the purity of the product was checked by NMR. Yield: 65%, 60 mg.

The following compounds were synthesized by a route similar to that described for **JXL001**: **JXL002**, **JXL003**, **JXL004**, **JXL005**, **JXL006**, **JXL007**, **JXL012**, **JXL013**, **JXL014**, **JXL021**, **JXL025**, **JXL026**, **JXL027**, **JXL028**, **JXL029**, **JXL035**, and **JXL093**.

Experimental Details for the Synthesis of JXL020. To the solution of indole-3-carboxaldehyde (3 mmol, 435 mg) in dry DMF (6 mL) were added 3,5-bis(trifluoromethyl)benzyl bromide (1.2 equiv, 3.6 mmol, 660 μL) and KOH (1.2 equiv, 3.6 mmol, 200 mg) at 0 $^\circ\text{C}$. The reaction mixture was stirred at 21 $^\circ\text{C}$ for 2 h. After the reaction completion shown by TLC, water (6 mL) was added into the reaction vial. The reaction mixture was extracted by dichloromethane (15 mL \times 3). The combined organic layer was dried by sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes/ethyl acetate = 8:1) to provide the desired product. Yield: 90%, 1001.7 mg.

To the solution of 1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indole-3-carbaldehyde (1 mmol, 371 mg) in ethanol (1 mL) were added ethyl 2-cyanoacetate (1.3 equiv, 1.3 mmol, 140 μL) and L-proline (40 mol %, 0.4 mmol, 58 mg). The reaction was stirred at 21 $^\circ\text{C}$ for 12 h and a yellow solid precipitated gradually. After completion of the reaction, ice-cold water (2 mL) was added into the reaction vial. The solid was separated by Buchner funnel filtration and washed with water (2 mL \times 3) and dried to afford the desired product. Yield: 93%, 433 mg.

To the solution of (*E*)-ethyl 3-(1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylate (0.21 mmol, 100 mg) in THF (2 mL) was added 0.5 N LiOH solution (3 equiv, 0.4 mmol, 0.8 mL). The reaction mixture was stirred at 21 $^\circ\text{C}$ for 1 h. After the reaction completion shown by TLC, THF was evaporated. Concentrated HCl was added dropwise to acidify the reaction mixture until pH was lower than 1; meanwhile, a yellow solid precipitated. Ice-cold water (5 mL) was added to the reaction mixture, and the solid was separated by Buchner funnel filtration and washed with water (5 mL \times 3). After drying by vacuum, the solid was washed with 2 mL of solvent mixture (hexanes/ethyl acetate = 5:1) 5–10 times and monitored by TLC until nonpolar impurities disappear (the nonpolar compound was the retro-aldol condensation product, which can be recovered from the filtrate). Finally, the purity of the product was checked by NMR. Yield: 55%, 52 mg.

The following compounds were synthesized by a route similar to that described for **JXL020**: **JXL008**, **JXL009**, **JXL010**, **JXL011**, **JXL015**, **JXL016**, **JXL017**, **JXL018**, **JXL019**, **JXL036**, **JXL037**, **JXL038**, **JXL039**, **JXL040**, **JXL041**, **JXL050**, **JXL051**, **JXL052**, **JXL053**, **JXL054**, **JXL055**, **JXL056**, **JXL057**, **JXL058**, **JXL059**, **JXL060**, **JXL061**, **JXL062**, **JXL063**, **JXL064**, **JXL065**, **JXL066**, **JXL068**, **JXL078**, **JXL081**, **JXL089**, **JXL090**, and **JXL091**.

Experimental Details for the Synthesis of JXL024. To the solution of 1-phenyl-1*H*-indole-3-carbaldehyde (0.4 mmol, 90 mg) in AcOH (3 mL) were added thiazolidine-2,4-dione (1 equiv, 0.4 mmol, 46.8 mg) and NaOAc (3 equiv, 98 mg). The reaction mixture was stirred at reflux for 24 h. After it was cooled to 21 $^\circ\text{C}$, the reaction mixture was filtered by vacuum filtration and washed with AcOH (3 mL \times 3) and water (5 mL \times 3). After drying by vacuum, the desired product was produced. Yield: 34%, 44 mg.

The following compounds were synthesized by a route similar to that described for **JXL024**: **JXL023**, **JXL067**, **JXL070**, **JXL072**, **JXL074**, and **JXL075**.

Experimental Details for the Synthesis of JXL022. To the solution of 4-pyridinecarboxaldehyde (1 mmol, 107 mg) in ethanol (1 mL) were added ethyl 2-cyanoacetate (1.3 equiv, 1.3 mmol, 140 μL) and L-proline (40 mol %, 0.4 mmol, 58 mg). The reaction was stirred at 21 $^\circ\text{C}$ for 12 h, and a yellow solid precipitated gradually. After completion of the reaction, ice-cold water (2 mL) was added into the reaction vial. The solid was separated by Buchner funnel filtration and washed with water (2 mL \times 3) and dried to afford the desired product, ethyl (*E*)-2-cyano-3-(pyridin-4-yl)acrylate, which was used for the next step without further purification.

To the solution of (*E*)-2-cyano-3-(pyridin-4-yl)acrylate (0.21 mmol, 42.4 mg) in THF (2 mL) was added 0.5 N LiOH solution (3 equiv, 0.4 mmol, 0.8 mL). The reaction mixture was stirred at 21 $^\circ\text{C}$ for 1 h. After the reaction completion shown by TLC, THF was evaporated. Concentrated HCl was added dropwise to acidify the reaction mixture until pH was lower than 1; meanwhile, a yellow solid precipitated. Ice-cold water (5 mL) was added to the reaction mixture, and the solid was separated by Buchner funnel filtration and

washed with water (5 mL \times 3). After drying by vacuum, the solid was washed with 2 mL of solvent mixture (hexanes/ethyl acetate = 5:1) 5–10 times and monitored by TLC until nonpolar impurities disappear. Finally, the purity of the product was checked by NMR. Yield: 64%, 23.4 mg.

The following compounds were synthesized by a route similar to that described for **JXL022**: **JXL030**, **JXL031**, **JXL032**, **JXL033**, **JXL034**, **JXL042**, **JXL43**, **JXL044**, **JXL045**, **JXL046**, **JXL047**, **JXL048**, and **JXL049**.

Experimental Details for the Synthesis of JXL069. To the solution of 7-azaindole **4** (1 mmol, 118 mg) in dry DMF (2 mL) were added 3,5-bis(trifluoromethyl)benzyl bromide (1.2 equiv, 1.2 mmol, 220 μ L) and KOH (1.2 equiv, 1.2 mmol, 67 mg) at 0 °C. The reaction mixture was stirred at 21 °C for 2 h. After the reaction completion shown by TLC, water (6 mL) was added to the reaction vial. The reaction mixture was extracted by dichloromethane (15 mL \times 3). The combined organic layer was dried by sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes/ethyl acetate = 12:1) to provide the desired benzylation product **5**.

POCl₃ (1 mmol, 90 μ L) was added dropwise to DMF (2 mL) at 0 °C under argon. After stirring for 10 min, a solution of compound **2** (1 mmol, 334 mg) in DMF (2 mL) was added slowly with stirring. The mixture was kept at 21 °C overnight. The reaction was quenched by adding water (5 mL) at 0 °C and then extracted with dichloromethane (10 mL \times 3). The combined organic layer was dried by sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes/ethyl acetate = 4:1) to provide the desired aldehyde **12**.

To the solution of aldehyde **3** (1 mmol, 372 mg) in ethanol (2 mL) were added ethyl 2-cyanoacetate (1.3 equiv, 1.3 mmol, 140 μ L) and *L*-proline (40 mol %, 0.4 mmol, 58 mg). The reaction was stirred at 21 °C for 12 h, and a yellow solid precipitated gradually. After completion of the reaction, ice-cold water (2 mL) was added into the reaction vial. The solid was separated by Buchner funnel filtration and washed with water (2 mL \times 3) and dried to afford the desired product **JXL082**.

To the solution of **JXL082** (0.21 mmol, 100 mg) in THF (2 mL) was added 0.5 N LiOH solution (3 equiv, 0.4 mmol, 0.8 mL). The reaction mixture was stirred at 21 °C for 1 h. After the reaction completion shown by TLC, THF was evaporated. Concentrated HCl was added dropwise to acidify the reaction mixture until pH was lower than 1; meanwhile, a yellow solid precipitated. Ice-cold water (5 mL) was added to the reaction mixture, and the solid was filtrated by a PYREX Hirsch-Type Funnel with a fritted disc and washed with water (5 mL \times 3). After drying by vacuum, the solid was washed with 2 mL of solvent mixture (hexanes/ethyl acetate = 5:1) 5–10 times and monitored by TLC until nonpolar impurities disappeared (the nonpolar compound was the retro-Aldol condensation product, which can be recovered from the filtrate). Finally, the purity of the product **JXL069** was checked by NMR.

The following compounds were synthesized by a route similar to that described for **JXL069**: **JXL072**, **JXL073**, **JXL076**, **JXL077**, **JXL082**, **JXL087**, and **JXL088**.

Experimental Details for the Synthesis of JXL079. A flask containing Mg powder (10 mmol, 240 mg) and a stir bar was sealed and vacuumed and refilled with argon three times. Anhydrous diethyl ether (32 mL) and bis(trifluoromethyl)benzyl bromide (8 mmol, 1.46 mL) were added to the reaction flask. The reaction mixture was stirred and refluxed for 30 min, and then ground dry ice powder (5 g) was added into the reaction flask. After 1 h, the reaction was complete as shown by TLC. The extra Mg powder was filtered off, and the solvent was evaporated under vacuum. HCl (1 N, 20 mL) was added to the residue, and the precipitate was filtered and dried to provide the desired carboxylic acid.

A flask containing a stir bar was sealed, vacuumed, and refilled with argon three times. Anhydrous dichloromethane (20 mL) and DIBAL (1 M in hexanes, 6 mmol, 6 mL) were added to the flask. The crude carboxylic acid (2 mmol, 544 mg) dissolved in dry dichloromethane (10 mL) was added to the reaction flask at –78 °C. After 2 h, the

reaction was complete as shown by TLC and it was then quenched by adding sat. ammonium chloride (10 mL). The resulting mixture was extracted with dichloromethane (20 mL \times 3), and the organic phases were combined and evaporated on a rotary evaporator. The residue was purified by flash column chromatography (hexanes/ethyl acetate = 10:1) to afford the desired product 2-(3,5-bis(trifluoromethyl)phenyl)ethan-1-ol (yield: 90%, 464 mg).

To a solution of 2-(3,5-bis(trifluoromethyl)phenyl)ethan-1-ol (0.2 mmol, 51.6 mg) in dichloromethane (2 mL) were added triethylamine (0.22 mmol, 31 μ L) and mesyl chloride (MsCl, 0.2 mmol, 17 μ L) at 0 °C. After stirring for 1 h, the reaction was complete as shown by TLC. The solvent was removed by flowing air over the open flask. The residue was purified by flash column chromatography (hexanes/ethyl acetate = 10:1) to afford the desired product 3,5-bis(trifluoromethyl)phenethyl methanesulfonate (yield: 82%, 55.1 mg).

A flask containing NaH (60%, 0.22 mmol, 8.8 mg) and a stir bar was sealed, vacuumed, and refilled with argon three times. Anhydrous THF (3 mL) and a solution of *tert*-butyl (*E*)-2-cyano-3-(1*H*-indol-3-yl)acrylate (0.2 mmol, 53.6 mg) in THF (2 mL) were added into the reaction flask. The reaction mixture was stirred for 30 min, and then 3,5-bis(trifluoromethyl)phenethyl methanesulfonate (0.164 mmol, 55.1 mg) in 2 mL of THF was added. The reaction was stirred for 24 h and quenched by sat. NH₄Cl solution. The resulting mixture was extracted with dichloromethane (4 mL \times 3), and the organic phases were combined and evaporated on a rotary evaporator. The residue was purified by flash column chromatography (hexanes/ethyl acetate = 10:1) to afford the desired product *tert*-butyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)phenethyl)-1*H*-indol-3-yl)-2-cyano-acrylate (yield: 68%, 69 mg).

To a solution of methyl *tert*-butyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)phenethyl)-1*H*-indol-3-yl)-2-cyanoacrylate (0.1 mmol, 50.8 mg) in dichloromethane (2 mL) was added trifluoroacetic acid (3 equiv, 0.3 mmol, 34 μ L). The reaction mixture was stirred at 21 °C for 30 min, and a yellow solid precipitated. After the reaction was complete as shown by TLC, the reaction solvent was evaporated by flowing air over the open flask. The solid was washed with 2 mL of solvent mixture (hexanes/ethyl acetate = 5:1) 5–10 times and monitored by TLC until all of the nonpolar impurities disappeared. Finally, the purity of the product was checked by NMR. Yield: 87%, 39 mg.

Experimental Details for the Synthesis of JXL080. To a solution of ethyl (*E*)-2-cyano-3-(1*H*-indol-3-yl)acrylate (0.5 mmol, 112 mg) in dichloromethane (5 mL) were added 2-(3,5-bis(trifluoromethyl)phenyl)acetic acid (0.55 mmol, 150 mg), 4-(*N,N*-dimethylamino)pyridine (DMAP; catalytic amount, 6 mg), and DCC (0.5 mmol, 103 mg) at 0 °C. The mixture was allowed to reach 21 °C and stirred overnight. The white precipitate was filtered, and the resulting solution was concentrated in vacuum. The solid was purified by flash column chromatography (hexanes/ethyl acetate = 10:1) to afford the desired product (yield: 78%, 192.6 mg).

Experimental Details for the Synthesis of JXL083. To a solution of 1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indole-3-carbaldehyde (10 mmol, 3.71 g) in acetone (60 mL) were added 2-methyl-2-butene (9 mL), NaH₂PO₄ (3 equiv, 4.4 g), and NaClO₂ (6.6 mmol, 6 g) in 6 mL of water. The reaction mixture was stirred at 21 °C for 24 h. After the reaction was complete as shown by TLC, the reaction solvent was evaporated on a rotary evaporator. The crude material was dissolved in ethyl acetate (30 mL) and water (30 mL) and extracted with ethyl acetate (30 mL \times 3). The organic phase was combined, dried with sodium sulfate, and evaporated on a rotary evaporator. The solid was purified by flash column chromatography (hexanes/ethyl acetate = 2:1) to afford the desired product 1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indole-3-carboxylic acid (yield: 89%, 3.44 g).

A 100 mL round-bottom flask with a stir bar containing the carboxylic acid (5 mmol, 1935 mg) from the previous step was sealed, vacuumed, and refilled with argon three times. To the flask was added 50 mL of dichloromethane and oxalyl chloride (25 mmol, 2.1 mL) dropwise. The reaction mixture was stirred at 21 °C for 1.5 h. The reaction solvent was evaporated by vacuum, and the resulting compound was used for the next step.

A 100 mL round-bottom flask with a stir bar was sealed, vacuumed, and refilled with argon three times. Diisopropylamine (5.5 mmol, 765 μL) and THF (10 mL) were added into the flask and it was cooled to $-78\text{ }^\circ\text{C}$. *n*-BuLi (2.5 M in hexanes, 5 mmol, 2 mL) was added slowly to the flask. After the mixture had stirred for 30 min, a solution of ethyl 2-cyanoacetate (5 mmol, 590 μL) in THF (10 mL) was added slowly to the flask. After the mixture had stirred for 1 h, a solution of acyl chloride (5 mmol from the previous step) in THF (5 mL) was added slowly to the reaction mixture. After 1 h, the reaction was quenched by adding an aqueous 1 M HCl solution (10 mL) and extracted with ethyl acetate (10 mL \times 3). The organic phase was combined, dried with sodium sulfate, and evaporated by a rotary evaporator. The solid was purified by flash column chromatography (hexanes/ethyl acetate = 10:1) to afford the desired product ethyl (Z)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyano-3-hydroxyacrylate (**JXL083**) (yield: 80%, 1.93 g).

Experimental Details for the Synthesis of JXL084. To a solution of **JXL083** (0.5 mmol, 240 mg) in dichloromethane (10 mL) were added pyridine (0.5 mmol, 40 μL) and acetyl chloride (1.0 mmol, 84 μL). The reaction mixture was stirred for 1 h, and TLC indicated that the reaction was complete. The reaction solvent was evaporated by flowing air over the open flask. The residue was purified by flash column chromatography (hexanes/ethyl acetate = 10:1) to afford the desired product ethyl (Z)-3-acetoxy-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylate (**JXL084**) (yield: 86%, 225 mg).

Experimental Details for the Synthesis of JXL085. To a solution of **JXL083** (0.5 mmol, 240 mg) in dichloromethane (10 mL) were added triethylamine (1.0 mmol, 139.5 μL) and phosphoryl chloride (0.55 mmol, 520 μL). The reaction mixture was stirred for 1 h at reflux, and TLC indicated that the reaction was complete. The reaction solvent was evaporated by flowing air over the open flask. The residue was purified by flash column chromatography (hexanes/ethyl acetate = 10:1) to afford the desired product ethyl (Z)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-3-chloro-2-cyanoacrylate (**JXL085**) (yield: 84%, 210 mg).

Experimental Details for the Synthesis of JXL086. To the solution of 1-(3,5-bis(trifluoromethyl)benzyl)-1H-indole-3-carboxaldehyde (1 mmol, 371 mg) in ethanol (3 mL) were added diethyl cyanomethylphosphate (1.3 equiv, 1.3 mmol, 204 μL) and L-proline (40 mol%, 0.4 mmol, 58 mg). The reaction was stirred at $50\text{ }^\circ\text{C}$ for 24 h. After completion of the reaction as indicated by TLC, the reaction solvent was evaporated by flowing air over the open flask. The solid was purified by flash column chromatography (hexanes/ethyl acetate = 2:1) to afford the desired product **JXL086** (yield: 90%, 477 mg).

Experimental Details for the Synthesis of JXL096. A solution of **JXL086** (30 mg, 0.057 mmol) in dichloromethane (2 mL) was cooled to $0\text{ }^\circ\text{C}$ and bromotrimethylsilane (40 μL , 0.3 mmol) was added dropwise under argon. The mixture was warmed to $21\text{ }^\circ\text{C}$ and stirred for 12 h. The solvent was evaporated under vacuum, and the resulting residue was then dissolved in methanol (2 mL). The mixture was stirred at $21\text{ }^\circ\text{C}$ for 2 h. Evaporation of all volatiles under vacuum gave the phosphoric acid **JXL096** (yield: 92%, 25 mg).

Characterization Data for JXL001–JX096. (E)-2-Cyano-3-(1-phenyl-1H-indol-3-yl)acrylic Acid (**JXL001/UK-5099**). ^1H NMR (500 MHz, DMSO- d_6) δ 8.59 (s, 1H), 8.56 (s, 1H), 8.06 (m, 1H), 7.65 (m, 4H), 7.53 (m, 2H), 7.34 (m, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.5, 145.6, 137.7, 136.3, 133.6, 130.5, 128.9, 128.0, 125.0, 124.9, 123.3, 119.9, 118.4, 111.9, 96.7. HRMS (ESI, m/z) calcd for $\text{C}_{18}\text{H}_{12}\text{N}_2\text{O}_2$ ($[\text{M} - \text{H}]^-$): 287.0821. Found: 287.0826.

(E)-2-Cyano-3-(1H-indol-3-yl)acrylic Acid (**JXL002**). ^1H NMR (500 MHz, DMSO- d_6) δ 12.48 (s, 1H), 8.51 (s, 1H), 8.49 (s, 1H), 7.91 (d, $J = 6.5$ Hz, 1H), 7.53 (d, $J = 7.0$ Hz, 1H), 7.23 (m, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 165.0, 146.5, 136.5, 132.4, 127.3, 123.9, 122.4, 118.9, 118.8, 113.2, 110.2, 94.0. HRMS (ESI, m/z) calcd for $\text{C}_{12}\text{H}_8\text{N}_2\text{O}_2$ ($[\text{M} - \text{H}]^-$): 211.0508. Found: 211.0511.

Ethyl (E)-2-Cyano-3-(1H-indol-3-yl)acrylate (**JXL003**). ^1H NMR (500 MHz, CDCl_3) δ 12.55 (s, 1H), 8.53 (s, 1H), 8.52 (s, 1H), 7.92 (d, $J = 7.6$ Hz, 1H), 7.53 (d, $J = 7.8$ Hz, 1H), 7.26 (app. t, $J = 7.4$ Hz, 1H), 7.22 (app. t, $J = 7.4$ Hz, 1H), 4.24 (q, $J = 7.0$ Hz, 2H), 1.26 (t, $J = 7.0$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 163.6, 147.0, 136.6, 133.0, 127.3, 124.0, 122.5, 118.9, 118.4, 113.3, 110.3, 92.6, 61.8, 14.5. HRMS (ESI, m/z) calcd for $\text{C}_{14}\text{H}_{12}\text{N}_2\text{O}_2$ ($[\text{M} - \text{H}]^-$): 239.0820. Found: 239.0818.

Ethyl (E)-2-Cyano-3-(1-phenyl-1H-indol-3-yl)acrylate (**JXL004**). ^1H NMR (500 MHz, CDCl_3) δ 8.71 (s, 1H), 8.66 (s, 1H), 7.90 (d, $J = 7.2$ Hz, 1H), 7.54 (m, 6H), 7.36 (m, 2H), 4.39 (q, $J = 7.1$ Hz, 2H), 1.42 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 163.6, 145.6, 137.8, 136.4, 133.2, 129.9, 128.5, 124.8, 124.4, 123.0, 118.5, 117.9, 111.6, 111.5, 95.4, 62.0, 14.3. HRMS (ESI, m/z) calcd for $\text{C}_{20}\text{H}_{16}\text{N}_2\text{O}_2$ ($[\text{M} + \text{H}]^+$): 317.1290. Found: 317.1269.

Ethyl (E)-3-(6-Chloro-1-phenyl-1H-indol-3-yl)-2-cyanoacrylate (**JXL005**). ^1H NMR (500 MHz, CDCl_3) δ 8.67 (s, 1H), 8.58 (s, 1H), 7.81 (d, $J = 8.5$ Hz, 1H), 7.60 (m, 2H), 7.52 (m, 4H), 7.34 (d, $J = 8.4$ Hz, 1H), 4.39 (q, $J = 7.1$ Hz, 2H), 1.41 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 163.4, 145.1, 137.3, 136.8, 133.5, 130.5, 130.0, 128.8, 126.8, 124.8, 123.6, 119.5, 117.5, 111.6, 111.4, 96.4, 62.1, 14.2. HRMS (ESI, m/z) calcd for $\text{C}_{20}\text{H}_{15}\text{ClN}_2\text{O}_2$ ($[\text{M} + \text{H}]^+$): 351.0900. Found: 351.0907.

Ethyl (E)-2-Cyano-3-(1-(2-methoxyphenyl)-1H-indol-3-yl)acrylate (**JXL006**). ^1H NMR (500 MHz, CDCl_3) δ 8.67 (s, 1H), 8.66 (s, 1H), 7.89 (d, $J = 7.8$ Hz, 1H), 7.49 (app. t, $J = 8.6$ Hz, 1H), 7.41 (d, $J = 7.6$ Hz, 1H), 7.35 (app. t, $J = 7.3$ Hz, 1H), 7.30 (t, $J = 7.5$ Hz, 1H), 7.23 (d, $J = 8.1$ Hz, 1H), 7.13 (m, 2H), 4.39 (q, $J = 7.1$ Hz, 2H), 3.81 (s, 3H), 1.41 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 163.8, 154.2, 146.0, 137.2, 135.2, 130.3, 127.8, 126.0, 124.0, 122.6, 120.8, 118.2, 118.0, 112.3, 111.8, 111.0, 94.7, 61.8, 55.7, 14.3. HRMS (ESI, m/z) calcd for $\text{C}_{21}\text{H}_{18}\text{N}_2\text{O}_3$ ($[\text{M} + \text{H}]^+$): 347.1396. Found: 347.1403.

Ethyl (E)-2-Cyano-3-(1-(4-methoxyphenyl)-1H-indol-3-yl)acrylate (**JXL007**). ^1H NMR (500 MHz, CDCl_3) δ 8.65 (s, 1H), 8.64 (s, 1H), 7.89 (d, $J = 7.2$ Hz, 1H), 7.44 (m, 3H), 7.35 (m, 2H), 7.07 (d, $J = 8.8$ Hz, 2H), 4.38 (q, $J = 7.1$ Hz, 2H), 3.90 (s, 3H), 1.41 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 163.8, 159.5, 145.6, 136.8, 133.5, 130.6, 128.3, 126.2, 124.2, 122.9, 118.4, 118.0, 115.0, 111.5, 111.1, 94.9, 61.9, 55.6, 14.2. HRMS (ESI, m/z) calcd for $\text{C}_{21}\text{H}_{18}\text{N}_2\text{O}_3$ ($[\text{M} + \text{H}]^+$): 347.1396. Found: 347.1401.

Ethyl (E)-2-Cyano-3-(1-(4-fluorobenzyl)-1H-indol-3-yl)acrylate (**JXL008**). ^1H NMR (500 MHz, CDCl_3) δ 8.60 (app. s, 2H), 7.85 (d, $J = 6.8$ Hz, 1H), 7.32 (m, 3H), 7.15 (m, 2H), 7.03 (app. t, 2H), 5.39 (s, 2H), 4.37 (q, $J = 7.1$ Hz, 2H), 1.40 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 163.7, 162.5 (d, $J_{\text{C-F}} = 247.7$ Hz), 145.7, 136.1, 133.8, 130.9, 128.6, 128.5, 124.0, 122.7, 118.6, 118.0, 116.0 (d, $J_{\text{C-F}} = 21.9$ Hz), 110.9, 110.4, 94.6, 61.9, 50.7, 14.2. HRMS (ESI, m/z) calcd for $\text{C}_{21}\text{H}_{17}\text{FN}_2\text{O}_2$ ($[\text{M} + \text{H}]^+$): 347.1352. Found: 347.1351.

Ethyl (E)-2-Cyano-3-(1-(3,4-difluorobenzyl)-1H-indol-3-yl)acrylate (**JXL009**). ^1H NMR (500 MHz, CDCl_3) δ 8.60 (s, 1H), 8.59 (s, 1H), 7.86 (d, $J = 7.8$ Hz, 1H), 7.33 (m, 2H), 7.28 (s, 1H), 7.13 (m, 1H), 6.95 (m, 1H), 6.89 (m, 1H), 5.39 (s, 2H), 4.38 (q, $J = 7.1$ Hz, 2H), 1.40 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 163.7, 150.7 (dd, $J = 251.2, 13.2$ Hz), 150.2 (dd, $J = 250.4, 12.6$ Hz), 145.7, 136.1, 133.7, 132.3, 128.6, 124.3, 122.9, 122.7, 120.0, 118.8, 118.0 (d, $J = 17.5$ Hz), 115.9 (d, $J = 18.0$ Hz), 110.8, 110.6, 95.2, 62.1, 50.4, 14.4. HRMS (ESI, m/z) calcd for $\text{C}_{21}\text{H}_{16}\text{F}_2\text{N}_2\text{O}_2$ ($[\text{M} + \text{H}]^+$): 367.1258. Found: 367.1272.

Ethyl (E)-2-Cyano-3-(1-(3,5-difluorobenzyl)-1H-indol-3-yl)acrylate (**JXL010**). ^1H NMR (500 MHz, CDCl_3) δ 8.61 (s, 1H), 8.59 (s, 1H), 7.87 (d, $J = 7.1$ Hz, 1H), 7.33 (m, 2H), 7.26 (d, $J = 7.2$ Hz, 1H), 6.75 (app. t, $J = 8.7$ Hz, 1H), 6.64 (app. d, $J = 5.7$ Hz, 2H), 5.41 (s, 2H), 4.38 (q, $J = 7.1$ Hz, 2H), 1.40 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 163.6, 163.3 (dd, $J_{\text{C-F}} = 251.0, 12.5$ Hz), 145.5, 139.2, 136.0, 133.6, 128.4, 124.3, 122.9, 118.7, 118.0, 110.7, 110.6, 109.5 (dd, $J_{\text{C-F}} = 19.9, 6.4$ Hz), 103.8 (t, $J_{\text{C-F}} = 25.2$ Hz), 95.3, 62.0, 50.4, 14.2. HRMS (ESI, m/z) calcd for $\text{C}_{21}\text{H}_{16}\text{F}_2\text{N}_2\text{O}_2$ ($[\text{M} + \text{H}]^+$): 367.1258. Found: 367.1264.

Ethyl (E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylate (**JXL011**). ^1H NMR (500 MHz, CDCl_3) δ 8.62 (s, 1H), 8.61 (s, 1H), 7.90 (d, $J = 7.7$ Hz, 1H), 7.85 (s, 1H), 7.57 (s, 2H), 7.35 (m, 2H), 7.23 (d, $J = 7.8$ Hz, 1H), 5.56 (s, 2H), 4.38 (q, $J = 7.1$ Hz, 2H), 1.40 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ

163.6, 145.6, 138.1, 136.0, 133.4, 132.7 (q, $J = 33.8$ Hz), 128.6, 126.8, 124.7, 123.2, 122.9 (q, $J = 273.4$ Hz), 122.6, 119.0, 118.0, 111.2, 110.4, 95.9, 62.1, 50.5, 14.3. HRMS (ESI, m/z) calcd for $C_{23}H_{16}F_6N_2O_2$ ($[M + H]^+$): 467.1194. Found: 467.1206.

(E)-2-Cyano-3-(1-(4-methoxyphenyl)-1H-indol-3-yl)acrylic Acid (JXL012). 1H NMR (500 MHz, DMSO- d_6) δ 8.54 (s, 1H), 8.52 (s, 1H), 8.05 (d, $J = 7.7$ Hz, 1H), 7.58 (app. d, $J = 8.7$ Hz, 2H), 7.44 (m, 1H), 7.33 (m, 2H), 7.16 (app. d, $J = 8.7$ Hz, 2H), 3.82 (s, 3H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.7, 160.5, 136.7, 133.9, 130.5, 127.8, 126.6, 124.8, 123.2, 120.0, 119.5, 118.6, 115.6, 115.4, 111.9, 110.9, 55.9. HRMS (ESI, m/z) calcd for $C_{19}H_{14}N_2O_3$ ($[M + H]^+$): 319.1083. Found: 319.1077.

(E)-3-(6-Chloro-1-phenyl-1H-indol-3-yl)-2-cyanoacrylic Acid (JXL013). 1H NMR (500 MHz, DMSO- d_6) δ 13.58 (br. s, 1H), 8.59 (s, 1H), 8.54 (s, 1H), 8.11 (d, $J = 7.5$ Hz, 1H), 7.67 (m, 4H), 7.53 (m, 2H), 7.35 (d, $J = 7.5$ Hz, 1H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.3, 145.4, 137.3, 136.7, 134.4, 130.6, 129.6, 129.1, 126.7, 125.1, 123.5, 121.5, 118.2, 111.6, 111.3, 97.9. HRMS (ESI, m/z) calcd for $C_{18}H_{11}ClN_2O_2$ ($[M + H]^+$): 323.0587. Found: 323.0587.

(E)-2-Cyano-3-(1-(2-methoxyphenyl)-1H-indol-3-yl)acrylic Acid (JXL014). 1H NMR (500 MHz, DMSO- d_6) δ 8.55 (s, 1H), 8.47 (s, 1H), 8.02 (d, $J = 7.4$ Hz, 1H), 7.54 (m, 2H), 7.31 (m, 3H), 7.15 (m, 2H), 3.74 (s, 3H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.6, 154.2, 145.8, 137.3, 135.1, 131.1, 128.2, 127.3, 125.7, 124.6, 123.0, 121.5, 119.3, 118.5, 113.6, 112.2, 110.8, 96.1, 56.2. HRMS (ESI, m/z) calcd for $C_{19}H_{14}N_2O_3$ ($[M + H]^+$): 319.1083. Found: 319.1084.

(E)-2-Cyano-3-(1-(4-fluorobenzyl)-1H-indol-3-yl)acrylic Acid (JXL015). 1H NMR (500 MHz, DMSO- d_6) δ 8.64 (s, 1H), 8.46 (s, 1H), 7.93 (d, $J = 7.1$ Hz, 1H), 7.61 (d, $J = 7.3$ Hz, 1H), 7.33 (m, 2H), 7.26 (m, 2H), 7.16 (m, 2H), 5.60 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.8, 162.0 (d, $J_{C-F} = 244.3$ Hz), 145.6, 136.4, 134.6, 133.1, 130.0, 128.0, 124.1, 122.8, 119.2, 118.5, 116.0 (d, $J_{C-F} = 21.7$ Hz), 112.0, 109.8, 95.0, 49.6. HRMS (ESI, m/z) calcd for $C_{19}H_{13}FN_2O_2$ ($[M + H]^+$): 321.1039. Found: 321.1046.

(E)-2-Cyano-3-(1-(3,4-difluorobenzyl)-1H-indol-3-yl)acrylic Acid (JXL016). 1H NMR (500 MHz, DMSO- d_6) δ 13.34 (br. s, 1H), 8.62 (s, 1H), 8.47 (s, 1H), 7.94 (d, $J = 7.3$ Hz, 1H), 7.61 (d, $J = 7.6$ Hz, 1H), 7.40 (m, 2H), 7.27 (m, 2H), 7.10 (br. s, 1H), 5.61 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.8, 149.7 (dd, $J_{C-F} = 253.3$, 13.6 Hz), 149.4 (dd, $J_{C-F} = 246.3$, 11.6 Hz), 145.7, 136.4, 134.7, 134.5, 128.0, 124.7 (dd, $J_{C-F} = 5.9$, 3.0 Hz), 124.2, 122.8, 119.2, 118.4, 118.3 (d, $J_{C-F} = 17.0$ Hz), 117.1 (d, $J_{C-F} = 17.6$ Hz), 112.0, 109.9, 95.0, 49.3. HRMS (ESI, m/z) calcd for $C_{19}H_{12}F_2N_2O_2$ ($[M + H]^+$): 339.0945. Found: 339.0946.

(E)-2-Cyano-3-(1-(3,5-difluorobenzyl)-1H-indol-3-yl)acrylic Acid (JXL017). 1H NMR (500 MHz, DMSO- d_6) δ 8.67 (s, 1H), 8.48 (s, 1H), 7.95 (d, $J = 4.8$ Hz, 1H), 7.59 (d, $J = 4.4$ Hz, 1H), 7.27 (m, 2H), 7.15 (s, 1H), 6.98 (br. s, 2H), 5.65 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.8, 162.9 (dd, $J_{C-F} = 247.0$, 12.8 Hz), 145.6, 141.4, 136.4, 134.8, 128.0, 124.2, 122.9, 119.3, 118.4, 111.9, 111.0 (d, $J_{C-F} = 26.1$ Hz), 110.8, 103.8 (t, $J_{C-F} = 26.5$ Hz), 95.5, 49.5. HRMS (ESI, m/z) calcd for $C_{19}H_{12}F_2N_2O_2$ ($[M + H]^+$): 339.0945. Found: 339.0944.

Ethyl (E)-3-(1-Benzyl-1H-indol-3-yl)-2-cyanoacrylate (JXL018). 1H NMR (500 MHz, $CDCl_3$) δ 8.61 (s, 1H), 8.60 (s, 1H), 7.85 (d, $J = 7.9$ Hz, 1H), 7.33 (m, 6H), 7.17 (m, 2H), 5.42 (s, 2H), 4.37 (q, $J = 7.1$ Hz, 2H), 1.40 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 163.8, 145.7, 136.2, 135.1, 134.0, 129.0, 128.5, 128.2, 126.8, 124.0, 122.6, 118.5, 118.1, 111.0, 110.3, 94.3, 61.8, 51.4, 14.2. HRMS (ESI, m/z) calcd for $C_{21}H_{18}N_2O_2$ ($[M + H]^+$): 331.1447. Found: 331.1442.

(E)-3-(1-Benzyl-1H-indol-3-yl)-2-cyanoacrylic Acid (JXL019). 1H NMR (500 MHz, DMSO- d_6) δ 13.34 (br. s, 1H), 8.65 (s, 1H), 8.48 (s, 1H), 7.93 (d, $J = 6.9$ Hz, 1H), 7.60 (d, $J = 6.8$ Hz, 1H), 7.25 (m, 7H), 5.62 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.8, 145.7, 136.8, 136.5, 134.7, 129.1, 128.2, 127.6, 124.0, 122.7, 120.0, 119.2, 118.5, 112.1, 109.7, 94.7, 50.4. HRMS (ESI, m/z) calcd for $C_{19}H_{14}N_2O_2$ ($[M + H]^+$): 303.1133. Found: 303.1118.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylic Acid (JXL020). 1H NMR (500 MHz, DMSO- d_6) δ 13.37 (br. s, 1H), 8.75 (s, 1H), 8.48 (s, 1H), 7.99 (m, 4H), 7.65 (s, 1H),

7.28 (m, 2H), 5.83 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.7, 145.7, 140.3, 136.3, 134.8, 131.1, 130.8 (q, $J = 31.1$ Hz), 128.9, 128.7, 127.9, 124.8, 124.3, 122.9 (q, $J = 273.4$ Hz), 122.2, 119.3, 118.3, 95.6, 49.2. HRMS (ESI, m/z) calcd for $C_{21}H_{12}F_6N_2O_2$ ($[M + H]^+$): 439.0881. Found: 439.0880.

2-((1-Phenyl-1H-indol-3-yl)methylene)malononitrile (JXL021). 1H NMR (500 MHz, $CDCl_3$) δ 8.62 (s, 1H), 8.13 (s, 1H), 7.80 (d, $J = 7.4$ Hz, 1H), 7.61 (m, 2H), 7.53 (m, 4H), 7.40 (m, 2H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 149.8, 137.3, 136.5, 133.6, 130.1, 128.9, 127.7, 124.9, 124.8, 123.7, 118.2, 115.1, 115.0, 111.9, 111.8, 73.7. HRMS (ESI, m/z) calcd for $C_{18}H_{11}N_3$ ($[M + H]^+$): 321.0698. Found: 321.0695.

Ethyl (E)-2-Cyano-3-(pyridin-4-yl)acrylate (JXL022). 1H NMR (500 MHz, $CDCl_3$) δ 8.81 (d, $J = 5.2$ Hz, 2H), 8.18 (s, 1H), 7.74 (d, $J = 5.2$ Hz, 2H), 4.41 (q, $J = 7.1$ Hz, 2H), 1.41 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 161.2, 152.0, 151.0, 137.9, 123.2, 114.2, 108.2, 63.2, 14.0. HRMS (ESI, m/z) calcd for $C_{11}H_{10}N_2O_2$ ($[M + H]^+$): 203.0820. Found: 203.0815.

(Z)-2-Imino-5-((1-phenyl-1H-indol-3-yl)methylene)thiazolidin-4-one (JXL023). 1H NMR (500 MHz, DMSO- d_6) δ 11.94 (br. s, 1H), 9.33 (br. s, 1H), 8.97 (s, 1H), 7.64 (m, 10H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 180.9, 174.8, 172.5, 138.6, 136.2, 130.5, 129.3, 128.1, 126.3, 124.8, 124.5, 122.2, 120.3, 119.7, 113.5, 111.5. HRMS (ESI, m/z) calcd for $C_{18}H_{13}N_3OS$ ($[M + H]^+$): 320.0858. Found: 320.0862.

(Z)-5-((1-Phenyl-1H-indol-3-yl)methylene)thiazolidine-2,4-dione (JXL024). 1H NMR (500 MHz, DMSO- d_6) δ 7.98 (m, 2H), 7.79 (s, 1H), 7.66 (app. d, $J = 7.7$ Hz, 2H), 7.62 (app. t, $J = 7.7$ Hz, 2H), 7.54 (d, $J = 8.0$ Hz, 1H), 7.49 (t, $J = 7.1$ Hz, 1H), 7.30 (m, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 172.5, 169.5, 138.4, 136.2, 130.5, 129.9, 128.3, 128.2, 125.0, 124.6, 122.4, 121.5, 121.4, 119.6, 113.0, 111.5. HRMS (ESI, m/z) calcd for $C_{18}H_{12}N_2O_2S$ ($[M + H]^+$): 321.0698. Found: 321.0695.

2-(Ethoxycarbonyl)-3-(1-phenyl-1H-indol-3-yl)acrylic Acid (a Mixture of E/Z Isomers, 1:1 Ratio) (JXL025). 1H NMR (500 MHz, DMSO- d_6) δ 13.08 (br. s, 1H), 7.87 (m, 3H), 7.61 (m, 4H), 7.52 (m, 2H), 7.30 (m, 2H), 4.26 (m, 2H), 1.23 (m, 3H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 168.9, 167.8, 166.1, 164.9, 138.3, 136.1, 132.9, 132.1, 131.1, 130.9, 130.8, 128.5, 124.9, 124.4, 122.6, 122.2, 119.5, 111.6, 111.0, 61.7, 61.3, 14.7, 14.4. HRMS (ESI, m/z) calcd for $C_{20}H_{17}NO_4$ ($[M + H]^+$): 336.1236. Found: 336.1239.

(E)-2-Cyano-3-(4-fluoro-1-phenyl-1H-indol-3-yl)acrylic Acid (JXL026). 1H NMR (500 MHz, DMSO- d_6) δ 8.65 (s, 1H), 8.61 (s, 1H), 7.67 (m, 4H), 7.57 (m, 1H), 7.36 (m, 2H), 7.19 (m, 1H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.4, 156.8 (d, $J_{C-F} = 245.6$ Hz), 146.4, 138.7, 137.6, 133.4, 130.7, 129.5, 125.9, 125.4, 118.2, 116.3, 116.2, 109.5 (d, $J_{C-F} = 34.5$ Hz), 109.2 (d, $J_{C-F} = 23.2$ Hz), 98.0. HRMS (ESI, m/z) calcd for $C_{18}H_{11}FN_2O_2$ ($[M + H]^+$): 307.0880. Found: 307.0885.

(E)-2-Cyano-3-(6-fluoro-1-phenyl-1H-indol-3-yl)acrylic Acid (JXL027). 1H NMR (500 MHz, DMSO- d_6) δ 13.59 (br. s, 1H), 8.62 (s, 1H), 8.59 (s, 1H), 8.16 (m, 1H), 7.66 (m, 4H), 7.56 (m, 1H), 7.36 (d, $J = 9.2$ Hz, 1H), 7.25 (m, 1H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.6, 160.8 (d, $J_{C-F} = 240.0$ Hz), 145.7, 137.6, 136.6, 134.4, 130.8, 129.1, 125.0, 124.7, 121.6, 118.4, 111.8 (d, $J_{C-F} = 24.2$ Hz), 111.5, 98.7 (d, $J_{C-F} = 26.2$ Hz), 97.7. HRMS (ESI, m/z) calcd for $C_{18}H_{11}FN_2O_2$ ($[M + H]^+$): 307.0883. Found: 307.0886.

(E)-2-Cyano-3-(7-fluoro-1-phenyl-1H-indol-3-yl)acrylic Acid (JXL028). 1H NMR (500 MHz, DMSO- d_6) δ 13.62 (br. s, 1H), 8.56 (s, 1H), 8.47 (s, 1H), 7.89 (br. s, 1H), 7.61 (m, 5H), 7.30 (br. s, 1H), 7.17 (br. s, 1H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.4, 149.7 (d, $J_{C-F} = 247.5$ Hz), 145.5, 139.0, 135.5, 131.9, 129.9, 129.3, 126.2, 124.3, 124.0, 118.3, 115.9, 111.8, 110.9 (d, $J_{C-F} = 17.4$ Hz), 98.1. HRMS (ESI, m/z) calcd for $C_{18}H_{11}FN_2O_2$ ($[M + H]^+$): 307.0883. Found: 307.0870.

(E)-2-Cyano-3-(5-fluoro-1-phenyl-1H-indol-3-yl)acrylic Acid (JXL029). 1H NMR (500 MHz, DMSO- d_6) δ 13.57 (br. s, 1H), 8.67 (s, 1H), 8.60 (s, 1H), 8.01 (d, $J = 9.0$ Hz, 1H), 7.70 (m, 4H), 7.58 (m, 2H), 7.24 (m, 1H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.6, 159.4 (d, $J_{C-F} = 237.8$ Hz), 145.8, 137.7, 135.1, 133.1, 130.7, 129.2, 129.1, 125.2, 118.5, 113.7, 113.2 (d, $J_{C-F} = 26.5$ Hz), 111.4,

105.7 (d, $J_{c-f} = 24.2$ Hz), 97.2. HRMS (ESI, m/z) calcd for $C_{18}H_{11}FN_2O_2$ ($[M + H]^+$): 307.0883. Found: 307.0872.

(E)-2-Cyano-3-(2-fluorophenyl)acrylic Acid (JXL030). 1H NMR (500 MHz, DMSO- d_6) δ 8.49 (s, 1H), 8.31 (app. t, $J = 7.2$ Hz, 1H), 7.63 (m, 1H), 7.36 (app. t, $J = 7.4$ Hz, 1H), 7.30 (m, 1H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 162.9, 161.5 (d, $J_{c-f} = 256.2$ Hz), 145.4 (d, $J_{c-f} = 7.8$ Hz), 135.0 (d, $J_{c-f} = 9.2$ Hz), 128.7, 124.7, 119.8 (d, $J_{c-f} = 10.9$ Hz), 115.8 (d, $J_{c-f} = 21.9$ Hz), 114.9, 105.9. HRMS (ESI, m/z) calcd for $C_{10}H_6FNO_2$ ($[M - H]^-$): 190.0304. Found: 190.0305.

(E)-2-Cyano-3-(4-fluorophenyl)acrylic Acid (JXL032). 1H NMR (500 MHz, DMSO- d_6) δ 8.30 (s, 1H), 8.10 (m, 2H), 7.29 (m, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 165.2 (d, $J_{c-f} = 255.2$ Hz), 163.5, 153.1, 133.3 (d, $J_{c-f} = 9.3$ Hz), 128.3, 116.0 (d, $J_{c-f} = 22.4$ Hz), 115.3, 103.2. HRMS (ESI, m/z) calcd for $C_{10}H_6FNO_2$ ($[M - H]^-$): 190.0304. Found: 190.0303.

(E)-2-Cyano-3-(4-(trifluoromethyl)phenyl)acrylic Acid (JXL034). 1H NMR (500 MHz, DMSO- d_6) δ 8.39 (s, 1H), 8.17 (d, $J = 7.7$ Hz, 2H), 7.84 (d, $J = 7.7$ Hz, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 162.8, 152.6, 135.4, 133.1 (q, $J_{c-f} = 32.9$ Hz), 131.0, 125.7, 123.7 (q, $J_{c-f} = 272.2$ Hz), 114.8, 106.7. HRMS (ESI, m/z) calcd for $C_{11}H_6F_3NO_2$ ($[M - H]^-$): 240.0272. Found: 240.0271.

(E)-3-(4-Chloro-1-phenyl-1H-indol-3-yl)-2-cyanoacrylic Acid (JXL035). 1H NMR (500 MHz, DMSO- d_6) δ 9.22 (s, 1H), 8.68 (s, 1H), 7.64 (m, 3H), 7.59 (m, 2H), 7.48 (d, $J = 7.6$ Hz, 1H), 7.38 (m, 1H), 7.32 (m, 1H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.5, 146.4, 138.0, 137.4, 134.5, 130.7, 130.5, 129.5, 125.7, 125.3, 124.9, 123.7, 118.4, 111.7, 110.9, 97.5. HRMS (ESI, m/z) calcd for $C_{18}H_{11}ClN_2O_2$ ($[M + H]^+$): 323.0587. Found: 323.0587.

(E)-2-Cyano-3-(1-(3,4-difluorobenzyl)-5-fluoro-1H-indol-3-yl)acrylic Acid (JXL036). 1H NMR (500 MHz, DMSO- d_6) δ 8.69 (s, 1H), 8.47 (s, 1H), 7.84 (d, $J = 9.6$ Hz, 1H), 7.63 (dd, $J = 8.9, 4.3$ Hz, 1H), 7.41 (m, 2H), 7.14 (m, 2H), 5.61 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.9, 159.4 (d, $J_{c-f} = 237.3$ Hz), 149.8 (dd, $J_{c-f} = 247.1, 12.7$ Hz), 149.5 (dd, $J_{c-f} = 246.6, 12.3$ Hz), 146.0, 136.1, 134.5, 133.1, 129.1, 125.0, 118.5, 118.4, 117.4, 113.6, 112.5 (d, $J_{c-f} = 26.2$ Hz), 110.1, 105.2 (d, $J_{c-f} = 25.1$ Hz), 95.5, 49.7. HRMS (ESI, m/z) calcd for $C_{19}H_{11}F_3N_2O_2$ ($[M + H]^+$): 357.0851. Found: 357.0857.

(E)-2-Cyano-3-(1-(3,5-difluorobenzyl)-5-fluoro-1H-indol-3-yl)acrylic Acid (JXL037). 1H NMR (500 MHz, DMSO- d_6) δ 8.71 (s, 1H), 8.48 (s, 1H), 7.85 (dd, $J = 9.6, 2.0$ Hz, 1H), 7.62 (dd, $J = 8.9, 4.3$ Hz, 1H), 7.16 (m, 2H), 7.00 (app. d, $J = 6.2$ Hz, 2H), 5.65 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.9, 163.0 (d, $J_{c-f} = 247.7$ Hz), 159.3 (d, $J_{c-f} = 237.6$ Hz), 145.9, 141.4, 136.3, 133.2, 129.1, 118.5, 113.6, 112.6 (d, $J_{c-f} = 26.3$ Hz), 111.3, 110.2, 105.3 (d, $J_{c-f} = 24.9$ Hz), 104.0 (t, $J_{c-f} = 25.2$ Hz), 95.9, 49.8. HRMS (ESI, m/z) calcd for $C_{19}H_{11}F_3N_2O_2$ ($[M + H]^+$): 357.0851. Found: 357.0851.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-5-fluoro-1H-indol-3-yl)-2-cyanoacrylic Acid (JXL038). 1H NMR (500 MHz, DMSO- d_6) δ 8.80 (s, 1H), 8.49 (s, 1H), 8.06 (s, 1H), 8.04 (s, 2H), 7.86 (dd, $J = 9.6, 2.1$ Hz, 1H), 7.69 (dd, $J = 8.9, 4.3$ Hz, 1H), 7.17 (dt, $J = 9.0, 2.2$ Hz, 1H), 5.83 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.8, 159.3 (d, $J_{c-f} = 237.5$ Hz), 145.9, 140.3, 136.2, 133.1, 131.1 (q, $J_{c-f} = 33.1$ Hz), 129.1, 123.6 (q, $J_{c-f} = 272.2$ Hz), 118.4, 113.5, 112.7, 112.5, 110.4, 105.5, 105.3, 96.1, 49.6. HRMS (ESI, m/z) calcd for $C_{21}H_{11}F_7N_2O_2$ ($[M + H]^+$): 457.0787. Found: 457.0787.

(E)-2-Cyano-3-(1-(3,4-difluorobenzyl)-6-fluoro-1H-indol-3-yl)acrylic Acid (JXL039). 1H NMR (500 MHz, DMSO- d_6) δ 8.64 (s, 1H), 8.46 (s, 1H), 7.99 (dd, $J = 8.7, 5.1$ Hz, 1H), 7.58 (dd, $J = 9.8, 1.8$ Hz, 1H), 7.43 (m, 2H), 7.12 (m, 2H), 5.57 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.8, 160.3 (d, $J_{c-f} = 239.3$ Hz), 149.8 (dd, $J_{c-f} = 239.3, 25.2$ Hz), 149.6 (dd, $J_{c-f} = 246.3, 25.2$ Hz), 145.8, 136.8, 135.3, 134.5, 125.1, 125.0, 124.6, 121.1, 118.5, 117.5, 111.3 (d, $J_{c-f} = 23.9$ Hz), 110.2, 98.7 (d, $J_{c-f} = 26.5$ Hz), 96.2, 49.4. HRMS (ESI, m/z) calcd for $C_{19}H_{11}F_3N_2O_2$ ($[M + H]^+$): 357.0851. Found: 457.0835.

(E)-2-Cyano-3-(1-(3,5-difluorobenzyl)-6-fluoro-1H-indol-3-yl)acrylic Acid (JXL040). 1H NMR (500 MHz, DMSO- d_6) δ 8.65 (s, 1H), 8.46 (s, 1H), 8.00 (dd, $J = 8.6, 5.2$ Hz, 1H), 7.57 (d, $J = 9.7$ Hz, 1H), 7.15 (m, 2H), 7.03 (s, 1H), 7.02 (s, 1H), 5.62 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.8, 163.0 (d, $J_{c-f} = 239.4$ Hz), 162.9 (d, $J_{c-f} = 248.6$ Hz), 160.4 (d, $J_{c-f} = 239.4$ Hz), 145.5, 141.3,

136.8, 135.2, 124.6, 121.1, 118.4, 111.3, 111.1, 110.3, 104.0 (t, $J_{c-f} = 25.2$ Hz), 98.7 (d, $J_{c-f} = 26.5$ Hz), 96.9, 49.6. HRMS (ESI, m/z) calcd for $C_{19}H_{11}F_3N_2O_2$ ($[M + H]^+$): 357.0851. Found: 357.0847.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-6-fluoro-1H-indol-3-yl)-2-cyanoacrylic Acid (JXL041). 1H NMR (500 MHz, DMSO- d_6) δ 8.74 (s, 1H), 8.48 (s, 1H), 8.06 (app. s, 3H), 8.01 (dd, $J = 8.7, 5.1$ Hz, 1H), 7.66 (dd, $J = 9.8, 2.0$ Hz, 1H), 7.13 (dt, $J = 9.3, 2.1$ Hz, 1H), 5.78 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.7, 159.9 (d, $J_{c-f} = 264.6$ Hz), 145.8, 140.2, 136.8, 135.3, 131.1 (q, $J_{c-f} = 33.3$ Hz), 129.1, 124.5, 123.6 (q, $J_{c-f} = 273.7$ Hz), 122.5, 121.2, 118.4, 111.4 (d, $J_{c-f} = 25.2$ Hz), 110.4, 98.6 (d, $J_{c-f} = 27.2$ Hz), 96.7, 49.4. HRMS (ESI, m/z) calcd for $C_{21}H_{11}F_7N_2O_2$ ($[M + H]^+$): 457.0787. Found: 457.0783.

(E)-2-Cyano-3-(3-fluoro-4-methylphenyl)acrylic Acid (JXL042). 1H NMR (500 MHz, DMSO- d_6) δ 8.28 (s, 1H), 7.78 (m, 2H), 7.49 (t, $J = 8.0$ Hz, 1H), 2.30 (s, 3H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 163.6, 160.9 (d, $J_{c-f} = 244.6$ Hz), 153.5, 133.0, 131.6 (d, $J_{c-f} = 7.5$ Hz), 130.8 (d, $J_{c-f} = 17.6$ Hz), 127.2, 117.9 (d, $J_{c-f} = 23.9$ Hz), 116.4, 104.5, 15.0. HRMS (ESI, m/z) calcd for $C_{11}H_8FNO_2$ ($[M - H]^-$): 204.0461. Found: 204.0460.

(E)-2-Cyano-3-(3,4-difluorophenyl)acrylic Acid (JXL043). 1H NMR (500 MHz, DMSO- d_6) δ 8.32 (s, 1H), 8.09 (d, $J = 8.0$ Hz, 1H), 7.94 (br. s, 1H), 7.67 (d, $J = 8.8$ Hz, 1H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 163.3, 152.3, 152.2 (dd, $J_{c-f} = 255.4, 12.6$ Hz), 149.9 (dd, $J_{c-f} = 248.2, 12.6$ Hz), 129.8, 128.8, 120.1 (d, $J_{c-f} = 17.6$ Hz), 119.1 (d, $J_{c-f} = 17.6$ Hz), 116.3, 105.8. HRMS (ESI, m/z) calcd for $C_{10}H_5F_2NO_2$ ($[M - H]^-$): 208.0210. Found: 208.0212.

(E)-2-Cyano-3-(2,4-difluorophenyl)acrylic Acid (JXL044). 1H NMR (500 MHz, DMSO- d_6) δ 14.2 (br. s, 1H), 8.27 (s, 1H), 8.24 (m, 1H), 7.52 (m, 1H), 7.35 (dt, $J = 8.6, 2.2$ Hz, 1H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 165.5 (dd, $J_{c-f} = 255.8, 12.6$ Hz), 163.1, 161.9 (dd, $J_{c-f} = 270.5, 12.6$ Hz), 145.1, 131.1, 117.0, 115.9, 113.5 (d, $J_{c-f} = 22.7$ Hz), 106.8, 105.6 (t, $J_{c-f} = 26.5$ Hz). HRMS (ESI, m/z) calcd for $C_{10}H_5F_2NO_2$ ($[M - H]^-$): 208.0210. Found: 208.0208.

(E)-3-(3,5-Bis(trifluoromethyl)phenyl)-2-cyanoacrylic Acid (JXL045). 1H NMR (500 MHz, CD $_3$ OD) δ 8.59 (s, 2H), 8.48 (s, 1H), 8.19 (s, 1H). HRMS (ESI, m/z) calcd for $C_{12}H_5F_6NO_2$ ($[M + H]^+$): 308.0146. Found: 308.0143.

(E)-2-Cyano-3-(4-fluoro-3-(trifluoromethyl)phenyl)acrylic Acid (JXL047). 1H NMR (500 MHz, DMSO- d_6) δ 8.46 (dd, $J = 7.1, 1.8$ Hz, 1H), 8.44 (s, 1H), 8.40 (m, 1H), 7.75 (m, 1H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 163.3, 161.1 (d, $J_{c-f} = 262.1$ Hz), 152.2, 137.6 (d, $J_{c-f} = 10.1$ Hz), 130.5, 129.3, 122.6 (q, $J_{c-f} = 272.8$ Hz), 118.9 (d, $J_{c-f} = 21.2$ Hz), 118.2 (qd, $J_{c-f} = 32.9, 12.6$ Hz), 116.2, 106.3. HRMS (ESI, m/z) calcd for $C_{11}H_5F_4NO_2$ ($[M + H]^+$): 258.0178. Found: 258.0176.

(E)-2-Cyano-3-phenylacrylic Acid (JXL048). 1H NMR (500 MHz, DMSO- d_6) δ 8.33 (s, 1H), 8.02 (m, 2H), 7.59 (m, 3H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 163.7, 154.9, 133.6, 132.0, 131.1, 129.8, 116.5, 104.3. HRMS (ESI, m/z) calcd for $C_{10}H_7NO_2$ ($[M - H]^-$): 172.0399. Found: 172.0396.

(E)-2-Cyano-3-(4-hydroxyphenyl)acrylic Acid (JXL049). 1H NMR (500 MHz, DMSO- d_6) δ 8.16 (s, 1H), 7.95 (d, $J = 8.8$ Hz, 2H), 6.92 (d, $J = 8.8$ Hz, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.2, 163.0, 153.7, 133.6, 122.8, 117.2, 116.3, 99.2. HRMS (ESI, m/z) calcd for $C_{10}H_7NO_3$ ($[M - H]^-$): 188.0348. Found: 188.0346.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-chloro-1H-indol-3-yl)-2-cyanoacrylic Acid (JXL050). 1H NMR (500 MHz, DMSO- d_6) δ 9.20 (s, 1H), 8.90 (s, 1H), 8.06 (m, 3H), 7.72 (d, $J = 7.5$ Hz, 1H), 7.32 (m, 2H), 5.86 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.7, 146.7, 140.0, 138.0, 135.6, 131.1 (q, $J_{c-f} = 33.1$ Hz), 129.1, 125.6, 125.1, 124.7, 124.5, 123.7, 123.6 (q, $J_{c-f} = 273.7$ Hz), 122.5, 111.6, 109.9, 96.7, 49.6. HRMS (ESI, m/z) calcd for $C_{21}H_{11}ClF_6N_2O_2$ ($[M + H]^+$): 473.0492. Found: 473.0494.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-bromo-1H-indol-3-yl)-2-cyanoacrylic Acid (JXL051). 1H NMR (500 MHz, DMSO- d_6) δ 9.37 (s, 1H), 8.91 (s, 1H), 8.07 (app. s, 3H), 7.77 (d, $J = 8.2$ Hz, 1H), 7.51 (d, $J = 7.6$ Hz, 1H), 7.23 (t, $J = 8.0$ Hz, 1H), 5.85 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.7, 146.3, 140.0, 138.0, 135.8, 131.0 (q, $J_{c-f} = 33.1$ Hz), 129.1, 127.9, 125.4, 124.9, 123.7 (q, $J_{c-f} =$

273.2 Hz), 122.5, 118.2, 113.6, 122.1, 110.2, 96.6, 49.5. HRMS (ESI, m/z) calcd for $C_{21}H_{11}BrF_6N_2O_2$ ($[M + H]^+$): 518.9968. Found: 518.9948.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-fluoro-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL052**). 1H NMR (500 MHz, DMSO- d_6) δ 8.80 (s, 1H), 8.58 (s, 1H), 8.05 (app. s, 3H), 7.54 (d, $J = 8.2$ Hz, 1H), 7.29 (d, $J = 12.9$ Hz, 1H), 7.09 (dd, $J = 11.1, 8.2$ Hz, 1H), 5.85 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.5, 155.7 (d, $J_{C-F} = 239.4$ Hz), 146.5, 140.1, 138.8, 134.5, 131.1 (q, $J_{C-F} = 33.0$ Hz), 129.0, 125.1 (d, $J_{C-F} = 7.6$ Hz), 124.7, 123.6 (q, $J_{C-F} = 273.5$ Hz), 122.5, 118.1, 116.1 (d, $J_{C-F} = 18.5$ Hz), 108.8, 108.5, 97.2, 49.7. HRMS (ESI, m/z) calcd for $C_{21}H_{11}F_7N_2O_2$ ($[M + H]^+$): 457.0787. Found: 457.0786.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-7-fluoro-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL053**). 1H NMR (500 MHz, DMSO- d_6) δ 8.73 (br. s, 1H), 8.48 (br. s, 1H), 8.06 (br. s, 1H), 7.89 (br. s, 2H), 7.78 (br. s, $J = 7.4$ Hz, 1H), 7.22 (br. s, 1H), 7.10 (br. s, 1H), 5.89 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.6, 159.7 (d, $J_{C-F} = 245.7$ Hz), 145.6, 141.0, 136.2, 132.1, 130.0 (q, $J_{C-F} = 33.0$ Hz), 128.4, 123.9, 123.8, 123.6 (q, $J_{C-F} = 273.3$ Hz), 122.4 (d, $J_{C-F} = 18.9$ Hz), 118.1, 115.8, 110.6, 110.2 (d, $J_{C-F} = 18.9$ Hz), 97.3, 52.0. HRMS (ESI, m/z) calcd for $C_{21}H_{11}F_7N_2O_2$ ($[M + H]^+$): 457.0787. Found: 457.0792.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-5-chloro-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL054**). 1H NMR (500 MHz, DMSO- d_6) δ 8.78 (s, 1H), 8.51 (s, 1H), 8.12 (s, 1H), 8.06 (s, 1H), 8.03 (s, 2H), 7.70 (d, $J = 7.4$ Hz, 1H), 7.33 (d, $J = 7.0$ Hz, 1H), 5.83 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.7, 145.7, 140.2, 135.9, 135.1, 131.1 (q, $J_{C-F} = 33.0$ Hz), 129.3, 129.0, 127.9, 124.4, 123.6 (q, $J_{C-F} = 274.0$ Hz), 122.5, 119.5, 118.4, 113.6, 109.9, 96.9, 49.5. HRMS (ESI, m/z) calcd for $C_{21}H_{11}ClF_6N_2O_2$ ($[M + H]^+$): 473.0492. Found: 473.0498.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-cyano-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL055**). 1H NMR (500 MHz, DMSO- d_6) δ 9.03 (s, 1H), 9.00 (s, 1H), 8.09 (m, 4H), 7.80 (d, $J = 7.3$ Hz, 1H), 7.47 (t, $J = 7.8$ Hz, 1H), 5.90 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.4, 144.4, 139.9, 136.7, 131.1 (q, $J_{C-F} = 33.0$ Hz), 129.7, 129.2, 126.6, 124.3, 123.6 (q, $J_{C-F} = 273.7$ Hz), 122.7, 118.6, 117.9, 117.8, 109.3, 101.7, 98.1, 49.5. HRMS (ESI, m/z) calcd for $C_{22}H_{11}F_6N_3O_2$ ($[M + H]^+$): 464.0834. Found: 464.0832.

(*E*)-1-(3,5-Bis(trifluoromethyl)benzyl)-3-(2-carboxy-2-cyanovinyl)-1*H*-indole-4-carboxylic Acid (**JXL056**). 1H NMR (500 MHz, DMSO- d_6) δ 9.26 (br. s, 1H), 8.88 (br. s, 1H), 8.05 (app. s, 3H), 7.94 (br. s, 1H), 7.76 (br. s, 1H), 7.37 (br. s, 1H), 5.85 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 169.3, 165.0, 150.6, 140.2, 137.7, 136.0, 131.1 (q, $J_{C-F} = 33.0$ Hz), 129.0, 125.9, 125.2, 124.7, 123.6 (q, $J_{C-F} = 273.7$ Hz), 123.5, 122.5, 118.3, 116.1, 110.1, 96.4, 49.3. HRMS (ESI, m/z) calcd for $C_{22}H_{12}F_6N_2O_4$ ($[M + H]^+$): 483.0779. Found: 483.0778.

(*E*)-3-(4-(Benzyloxy)-1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL057**). 1H NMR (500 MHz, DMSO- d_6) δ 9.15 (s, 1H), 8.72 (s, 1H), 8.04 (s, 1H), 7.99 (s, 2H), 7.54 (br. s, 3H), 7.37 (br. s, 2H), 7.28 (m, 2H), 6.95 (s, 1H), 5.81 (s, 2H), 5.28 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 165.0, 153.7, 148.6, 140.5, 137.9, 133.4, 131.2 (q, $J_{C-F} = 32.8$ Hz), 128.9, 128.8, 128.1, 127.5, 125.4, 124.7, 123.6 (q, $J_{C-F} = 273.7$ Hz), 122.3, 118.5, 117.0, 110.5, 105.6, 105.2, 95.5, 70.0, 49.5. HRMS (ESI, m/z) calcd for $C_{28}H_{18}F_6N_2O_3$ ($[M + H]^+$): 545.1300. Found: 545.1299.

(*E*)-3-(6-(Benzyloxy)-1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL058**). 1H NMR (500 MHz, DMSO- d_6) δ 8.63 (s, 1H), 8.43 (s, 1H), 8.05 (s, 1H), 8.03 (s, 2H), 7.86 (d, $J = 8.5$ Hz, 1H), 7.32 (m, 6H), 6.97 (d, $J = 8.3$ Hz, 1H), 5.77 (s, 2H), 5.09 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 165.0, 156.7, 146.1, 140.5, 137.5, 137.3, 134.3, 131.0 (q, $J_{C-F} = 32.8$ Hz), 128.9, 128.8, 128.3, 128.2, 126.9, 123.6 (q, $J_{C-F} = 273.4$ Hz), 122.4, 121.9, 118.5, 113.3, 110.5, 96.8, 95.6, 70.2, 49.2. HRMS (ESI, m/z) calcd for $C_{28}H_{18}F_6N_2O_3$ ($[M + H]^+$): 545.1300. Found: 545.1292.

(*E*)-3-(7-(Benzyloxy)-1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL059**). 1H NMR (500 MHz, DMSO- d_6) δ 8.61 (br. s, 1H), 8.45 (br. s, 1H), 7.97 (br. s, 1H), 7.60 (br. s, 2H), 7.51 (br. s, 1H), 7.25 (br. s, 2H), 7.16 (br. s, 2H), 6.91 (br. s, 1H), 5.94 (s, 2H), 5.13 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.8,

146.6, 145.7, 142.3, 136.6, 135.4, 130.7 (q, $J_{C-F} = 32.8$ Hz), 130.5, 128.8, 128.4, 128.0, 127.5, 125.7, 124.1, 123.6 (q, $J_{C-F} = 273.7$ Hz), 121.8, 118.3, 111.8, 110.3, 107.1, 96.2, 70.3, 52.4. HRMS (ESI, m/z) calcd for $C_{28}H_{18}F_6N_2O_3$ ($[M + H]^+$): 545.1300. Found: 545.1298.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-methoxy-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL060**). 1H NMR (500 MHz, DMSO- d_6) δ 8.99 (s, 1H), 8.71 (s, 1H), 8.05 (s, 1H), 8.00 (s, 2H), 7.24 (app. s, 2H), 6.82 (d, $J = 6.0$ Hz, 1H), 5.81 (s, 2H), 3.92 (s, 3H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 165.0, 154.8, 148.6, 140.5, 137.8, 133.3, 131.0 (q, $J_{C-F} = 32.8$ Hz), 128.9, 125.4, 123.6 (q, $J_{C-F} = 273.7$ Hz), 122.5, 118.5, 116.8, 110.5, 105.0, 104.3, 95.3, 56.2, 49.5. HRMS (ESI, m/z) calcd for $C_{22}H_{14}F_6N_2O_3$ ($[M + H]^+$): 469.0987. Found: 469.0987.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-5-bromo-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL061**). 1H NMR (500 MHz, DMSO- d_6) δ 8.75 (s, 1H), 8.49 (s, 1H), 8.25 (s, 1H), 8.06 (s, 1H), 8.02 (m, 3H), 7.64 (app. s, 1H), 7.44 (app. s, 1H), 5.82 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.7, 145.3, 140.3, 135.5, 135.3, 131.1 (q, $J_{C-F} = 32.8$ Hz), 129.8, 128.9, 127.0, 124.7, 123.6 (q, $J_{C-F} = 274.0$ Hz), 122.5, 122.3, 118.5, 113.9, 109.9, 97.7, 49.5. HRMS (ESI, m/z) calcd for $C_{21}H_{11}BrF_6N_2O_2$ ($[M + H]^+$): 516.9987. Found: 516.9983.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-6-bromo-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL062**). 1H NMR (500 MHz, DMSO- d_6) δ 8.72 (s, 1H), 8.47 (s, 1H), 8.06 (s, 2H), 8.03 (s, 1H), 7.94 (d, $J = 8.2$ Hz, 1H), 7.39 (d, $J = 7.9$ Hz, 1H), 5.81 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.7, 145.6, 140.2, 137.4, 135.3, 131.1 (q, $J_{C-F} = 32.8$ Hz), 129.1, 127.0, 125.9, 123.6 (q, $J_{C-F} = 274.0$ Hz), 122.5, 121.5, 118.2, 117.2, 114.8, 110.3, 96.9, 49.3. HRMS (ESI, m/z) calcd for $C_{21}H_{11}BrF_6N_2O_2$ ($[M + H]^+$): 516.9987. Found: 516.9965.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-6-chloro-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL063**). 1H NMR (500 MHz, DMSO- d_6) δ 8.75 (s, 1H), 8.47 (s, 1H), 8.05 (m, 3H), 8.00 (d, $J = 7.4$ Hz, 1H), 7.89 (s, 1H), 7.28 (d, $J = 6.5$ Hz, 1H), 5.81 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.7, 145.6, 140.2, 137.0, 135.4, 131.1 (q, $J_{C-F} = 32.8$ Hz), 129.2, 129.0, 126.7, 123.6 (q, $J_{C-F} = 274.0$ Hz), 123.3, 122.5, 121.2, 118.2, 111.9, 110.3, 97.0, 49.3. HRMS (ESI, m/z) calcd for $C_{21}H_{11}ClF_6N_2O_2$ ($[M + H]^+$): 473.0492. Found: 473.0493.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-7-chloro-1*H*-indol-3-yl)-2-cyanoacrylic Acid (**JXL064**). 1H NMR (500 MHz, DMSO- d_6) δ 8.69 (s, 1H), 8.50 (s, 1H), 8.03 (s, 1H), 7.97 (d, $J = 7.5$ Hz, 1H), 7.73 (s, 2H), 7.32 (d, $J = 7.1$ Hz, 1H), 7.25 (m, 1H), 6.14 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 164.5, 145.1, 142.1, 136.9, 131.5, 131.2 (q, $J_{C-F} = 32.8$ Hz), 127.6, 126.1, 124.1, 123.6 (q, $J_{C-F} = 274.0$ Hz), 122.5, 122.0, 118.8, 118.0, 116.9, 110.3, 97.9, 51.8. HRMS (ESI, m/z) calcd for $C_{21}H_{11}ClF_6N_2O_2$ ($[M + H]^+$): 473.0492. Found: 473.0496.

Ethyl (*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-bromo-1*H*-indol-3-yl)-2-cyanoacrylate (**JXL065**). 1H NMR (500 MHz, DMSO- d_6) δ 9.67 (s, 1H), 8.71 (s, 1H), 7.86 (s, 1H), 7.56 (s, 2H), 7.52 (d, $J = 7.4$ Hz, 1H), 7.17 (m, 2H), 5.55 (s, 2H), 4.37 (q, $J = 7.1$ Hz, 2H), 1.40 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 163.4, 146.9, 137.6, 137.4, 134.3, 132.7 (q, $J_{C-F} = 33.9$ Hz), 128.2, 126.7, 125.6, 125.1, 122.8 (q, $J_{C-F} = 273.4$ Hz), 122.7, 118.0, 114.9, 111.8, 109.9, 96.3, 62.2, 50.6, 14.3. HRMS (ESI, m/z) calcd for $C_{23}H_{15}BrF_6N_2O_2$ ($[M + H]^+$): 545.0299. Found: 545.0300.

Ethyl (*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-fluoro-1*H*-indol-3-yl)-2-cyanoacrylate (**JXL066**). 1H NMR (500 MHz, DMSO- d_6) δ 8.86 (s, 1H), 8.61 (s, 1H), 7.86 (s, 1H), 7.56 (s, 2H), 7.23 (m, 1H), 7.01 (m, 2H), 5.54 (s, 2H), 4.37 (q, $J = 7.1$ Hz, 2H), 1.39 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 163.1, 157.4 (d, $J = 243.9$ Hz), 147.4, 138.2 (d, $J = 10.1$ Hz), 137.8, 133.2, 132.7 (q, $J_{C-F} = 33.9$ Hz), 126.7, 125.3, 122.8 (q, $J_{C-F} = 273.4$ Hz), 122.7, 117.9, 116.9 (d, $J_{C-F} = 17.6$ Hz), 110.0, 109.0 (d, $J_{C-F} = 19.5$ Hz), 106.6, 97.1, 62.2, 50.7, 14.3. HRMS (ESI, m/z) calcd for $C_{23}H_{15}F_7N_2O_2$ ($[M + H]^+$): 485.1100. Found: 485.1100.

(*Z*)-5-((1-(3,5-Bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-methylene)-2-iminothiazolidin-4-one (**JXL067**). 1H NMR (500 MHz, DMSO- d_6) δ 9.21 (s, 1H), 9.01 (s, 1H), 8.03 (s, 1H), 7.94 (br. s, 3H), 7.86 (d, $J = 7.7$ Hz, 1H), 7.81 (s, 1H), 7.58 (d, $J = 7.9$ Hz, 1H), 7.24 (m, 1H), 7.19 (m, 1H), 5.76 (s, 2H). ^{13}C NMR (126 MHz, DMSO- d_6) δ 180.0, 174.9, 141.4, 136.5, 131.0 (q, $J_{C-F} = 32.8$ Hz),

130.4, 128.5, 127.9, 125.1, 123.8, 123.6 (q, $J_{c-f} = 274.0$ Hz), 122.1, 121.7, 120.8, 119.3, 111.8, 111.3, 48.9. HRMS (ESI, m/z) calcd for $C_{21}H_{13}F_6N_3OS$ ($[M + H]^+$): 470.0762. Found: 470.0761.

tert-Butyl (E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylate (JXL068). 1H NMR (500 MHz, $CDCl_3$) δ 8.57 (s, 1H), 8.54 (s, 1H), 7.86 (d, $J = 7.6$ Hz, 1H), 7.84 (s, 1H), 7.56 (s, 2H), 7.33 (m, 2H), 7.22 (d, $J = 7.4$ Hz, 1H), 5.55 (s, 2H), 1.59 (s, 9H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 162.4, 144.7, 138.2, 135.9, 133.0, 132.7 (q, $J_{c-f} = 32.8$ Hz), 128.6, 126.7, 124.5, 123.0, 122.9 (q, $J_{c-f} = 277.2$ Hz), 122.6, 119.0, 118.2, 111.1, 110.4, 97.7, 82.9, 50.4, 28.1. HRMS (ESI, m/z) calcd for $C_{25}H_{20}F_6N_2O_2$ ($[M + H]^+$): 495.1507. Found: 495.1503.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-pyrrolo[2,3-b]pyridin-3-yl)-2-cyanoacrylic Acid (JXL069). 1H NMR (500 MHz, $DMSO-d_6$) δ 8.85 (s, 1H), 8.47 (m, 3H), 8.09 (s, 2H), 8.04 (s, 1H), 7.35 (dd, $J = 7.1, 4.6$ Hz, 1H), 5.84 (s, 2H). ^{13}C NMR (126 MHz, $DMSO-d_6$) δ 164.6, 147.8, 146.1, 145.5, 141.0, 135.2, 131.0 (q, $J_{c-f} = 32.8$ Hz), 129.4, 129.2, 123.6 (q, $J_{c-f} = 274.0$ Hz), 122.3, 120.1, 119.1, 118.2, 108.7, 97.1, 47.8. HRMS (ESI, m/z) calcd for $C_{20}H_{11}F_6N_3O_2$ ($[M + H]^+$): 440.0834. Found: 440.0834.

(Z)-5-((4-Fluoro-1-phenyl-1H-indol-3-yl)methylene)thiazolidine-2,4-dione (JXL070). 1H NMR (500 MHz, $DMSO-d_6$) δ 12.45 (br. s, 1H), 8.11 (s, 1H), 7.83 (s, 1H), 7.65 (m, 4H), 7.52 (s, 1H), 7.32 (m, 2H), 7.12 (m, 1H). ^{13}C NMR (126 MHz, $DMSO-d_6$) δ 168.0, 167.7, 156.9 (d, $J_{c-f} = 245.7$ Hz), 138.7 (d, $J_{c-f} = 10.1$ Hz), 137.9, 130.8, 130.5, 128.8, 125.5, 125.4, 124.3, 120.0, 116.2 (d, $J_{c-f} = 18.9$ Hz), 110.7, 108.5, 108.1 (d, $J_{c-f} = 18.9$ Hz). HRMS (ESI, m/z) calcd for $C_{18}H_{11}FN_2O_2S$ ($[M + H]^+$): 339.0640. Found: 339.0610.

(Z)-5-((6-Fluoro-1-phenyl-1H-indol-3-yl)methylene)thiazolidine-2,4-dione (JXL071). 1H NMR (500 MHz, $DMSO-d_6$) δ 12.42 (s, 1H), 8.06 (app. s, 2H), 7.85 (s, 1H), 7.67 (m, 2H), 7.62 (m, 2H), 7.50 (m, 1H), 7.31 (d, $J = 9.6$ Hz, 1H), 7.16 (t, $J = 8.5$ Hz, 1H). ^{13}C NMR (126 MHz, $DMSO-d_6$) δ 168.0, 167.7, 160.6 (d, $J_{c-f} = 245.7$ Hz), 138.0, 136.3 (d, $J_{c-f} = 12.6$ Hz), 131.0, 130.6, 128.5, 125.0, 124.8, 123.5, 121.3 (d, $J_{c-f} = 10.0$ Hz), 120.0, 112.6, 111.1 (d, $J_{c-f} = 18.9$ Hz), 98.2 (d, $J_{c-f} = 18.9$ Hz). HRMS (ESI, m/z) calcd for $C_{18}H_{11}FN_2O_2S$ ($[M + H]^+$): 339.0604. Found: 339.0601.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-pyrrolo[3,2-b]pyridin-3-yl)-2-cyanoacrylic Acid (JXL072). 1H NMR (500 MHz, $DMSO-d_6$) δ 8.94 (s, 1H), 8.68 (s, 1H), 8.54 (s, 1H), 8.11 (m, 4H), 7.35 (s, 1H), 5.87 (s, 2H). ^{13}C NMR (126 MHz, $DMSO-d_6$) δ 164.5, 145.9, 144.2, 140.1, 135.9, 131.1 (q, $J_{c-f} = 32.8$ Hz), 129.6, 129.2, 124.7, 123.6 (q, $J_{c-f} = 274.0$ Hz), 122.5, 120.0, 119.5, 117.9, 110.1, 97.2, 49.8. HRMS (ESI, m/z) calcd for $C_{20}H_{11}F_6N_3O_2$ ($[M + H]^+$): 440.0834. Found: 440.0834.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-pyrrolo[2,3-c]pyridin-3-yl)-2-cyanoacrylic Acid (JXL073). 1H NMR (500 MHz, $DMSO-d_6$) δ 9.14 (s, 1H), 8.96 (s, 1H), 8.55 (s, 1H), 8.40 (d, $J = 5.1$ Hz, 1H), 8.19 (s, 2H), 8.11 (app. s, 2H), 5.95 (s, 2H). ^{13}C NMR (126 MHz, $DMSO-d_6$) δ 164.5, 145.5, 140.7, 139.9, 138.1, 134.5, 133.6, 131.1 (q, $J_{c-f} = 32.8$ Hz), 129.4, 127.2, 123.6 (q, $J_{c-f} = 274.0$ Hz), 122.7, 118.0, 114.6, 109.7, 97.9, 49.8. HRMS (ESI, m/z) calcd for $C_{30}H_{11}F_6N_3O_2$ ($[M + H]^+$): 440.0834. Found: 440.0837.

(Z)-5-Benzylidenethiazolidine-2,4-dione (JXL074). 1H NMR (500 MHz, $DMSO-d_6$) δ 12.60 (br. s, 1H), 7.77 (s, 1H), 7.58 (app. d, $J = 7.3$ Hz, 2H), 7.51 (app. t, $J = 7.4$ Hz, 2H), 7.46 (m, 1H). ^{13}C NMR (126 MHz, $DMSO-d_6$) δ 168.4, 167.8, 133.5, 132.3, 130.9, 130.5, 129.8, 124.0. HRMS (ESI, m/z) calcd for $C_{10}H_7NO_2S$ ($[M - H]^-$): 204.0119. Found: 204.0122.

(Z)-5-((1H-Indol-3-yl)methylene)thiazolidine-2,4-dione (JXL075). 1H NMR (500 MHz, $DMSO-d_6$) δ 12.28 (s, 1H), 12.11 (s, 1H), 8.03 (s, 1H), 7.87 (d, $J = 7.3$ Hz, 1H), 7.72 (s, 1H), 7.48 (d, $J = 7.6$ Hz, 1H), 7.23 (m, 1H), 7.18 (m, 1H). ^{13}C NMR (126 MHz, $DMSO-d_6$) δ 168.2, 167.8, 136.7, 129.1, 127.3, 125.0, 123.5, 121.5, 118.8, 116.7, 112.9, 110.9. HRMS (ESI, m/z) calcd for $C_{12}H_8N_2O_2S$ ($[M + H]^+$): 243.0228. Found: 243.0230.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-chloro-1H-pyrrolo[2,3-b]pyridin-3-yl)-2-cyanoacrylic Acid (JXL076). 1H NMR (500 MHz, $DMSO-d_6$) δ 9.00 (s, 1H), 8.98 (s, 1H), 8.36 (s, 1H), 8.12 (s, 2H), 8.04 (br. s, 1H), 7.47 (br. s, 1H), 5.85 (s, 2H). ^{13}C NMR (126 MHz,

$DMSO-d_6$) δ 164.4, 148.7, 145.8, 145.7, 140.2, 136.1, 135.3, 131.0 (q, $J_{c-f} = 32.8$ Hz), 129.5, 123.6 (q, $J_{c-f} = 274.0$ Hz), 122.4, 120.4, 117.7, 116.8, 108.2, 98.1, 48.2. HRMS (ESI, m/z) calcd for $C_{20}H_{10}ClF_6N_3O_2$ ($[M + H]^+$): 474.0444. Found: 474.0448.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-bromo-1H-pyrrolo[2,3-b]pyridin-3-yl)-2-cyanoacrylic Acid (JXL077). 1H NMR (500 MHz, $DMSO-d_6$) δ 9.17 (s, 1H), 9.02 (s, 1H), 8.27 (d, $J = 5.0$ Hz, 1H), 8.12 (s, 2H), 8.05 (s, 1H), 7.63 (d, $J = 5.0$ Hz, 1H), 5.85 (s, 2H). ^{13}C NMR (126 MHz, $DMSO-d_6$) δ 164.4, 148.2, 145.5, 145.3, 140.2, 135.4, 131.0 (q, $J_{c-f} = 32.8$ Hz), 129.6, 124.9, 123.7, 123.6 (q, $J_{c-f} = 274.0$ Hz), 122.4, 118.4, 117.7, 108.6, 97.9, 48.2. HRMS (ESI, m/z) calcd for $C_{20}H_{10}BrF_6N_3O_2$ ($[M + H]^+$): 517.9939. Found: 517.9935.

Methyl (E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylate (JXL078). 1H NMR (500 MHz, $CDCl_3$) δ 8.62 (s, 1H), 8.61 (s, 1H), 7.89 (d, $J = 7.7$ Hz, 1H), 7.84 (s, 1H), 7.57 (s, 2H), 7.35 (m, 2H), 7.24 (m, 1H), 5.55 (s, 2H), 3.92 (s, 3H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 164.5, 145.8, 138.1, 136.0, 133.5, 132.7 (q, $J_{c-f} = 32.8$ Hz), 128.6, 126.8, 124.7, 123.9, 123.6 (q, $J_{c-f} = 274.0$ Hz), 123.2, 119.1, 118.0, 111.2, 110.5, 95.4, 53.0, 50.5. HRMS (ESI, m/z) calcd for $C_{22}H_{14}F_6N_2O_2$ ($[M + H]^+$): 453.1038. Found: 453.1045.

(E)-3-(1-(3,5-Bis(trifluoromethyl)phenethyl)-1H-indol-3-yl)-2-cyanoacrylic Acid (JXL079). 1H NMR (500 MHz, CD_3OD) δ 8.51 (s, 1H), 8.14 (s, 1H), 7.83 (d, $J = 7.3$ Hz, 1H), 7.74 (s, 1H), 7.52 (m, 3H), 7.31 (m, 2H), 4.64 (t, $J = 6.4$ Hz, 2H), 3.35 (t, $J = 6.3$ Hz, 2H). ^{13}C NMR (126 MHz, CD_3OD) δ 165.3, 145.3, 141.1, 136.1, 133.6, 131.3 (q, $J_{c-f} = 32.8$ Hz), 129.3, 128.3, 123.7, 123.3 (q, $J_{c-f} = 272.5$ Hz), 122.3, 120.3, 118.1, 117.6, 110.6, 109.6, 94.1, 47.8, 35.0. HRMS (ESI, m/z) calcd for $C_{22}H_{14}F_6N_2O_2$ ($[M + H]^+$): 453.1038. Found: 453.1031.

Ethyl (E)-3-(1-(2-(3,5-Bis(trifluoromethyl)phenyl)acetyl)-1H-indol-3-yl)-2-cyanoacrylate (JXL080). 1H NMR (500 MHz, $CDCl_3$) δ 8.90 (s, 1H), 8.52 (s, 1H), 8.49 (d, $J = 8.1$ Hz, 1H), 7.88 (s, 1H), 7.86 (s, 2H), 7.78 (d, $J = 7.5$ Hz, 1H), 7.47 (m, 2H), 4.50 (s, 2H), 4.41 (q, $J = 7.1$ Hz, 2H), 1.43 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 168.0, 162.2, 144.2, 135.6, 134.4, 132.3 (q, $J_{c-f} = 33.6$ Hz), 130.1, 129.8, 128.8, 127.8, 127.3, 125.5, 123.1 (q, $J_{c-f} = 273.3$ Hz), 118.3, 117.1, 117.0, 115.9, 101.8, 62.8, 41.9, 14.3. HRMS (ESI, m/z) calcd for $C_{24}H_{16}F_6N_2O_3$ ($[M + H]^+$): 495.1143. Found: 495.1136.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-(morpholine-4-carbonyl)acrylonitrile (JXL081). 1H NMR (500 MHz, $CDCl_3$) δ 8.48 (s, 1H), 8.34 (s, 1H), 7.84 (s, 2H), 7.56 (app. s, 2H), 7.32 (m, 2H), 7.22 (m, 1H), 5.54 (s, 2H), 3.77 (br. s, 8H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 164.0, 145.2, 138.4, 135.8, 132.7 (q, $J_{c-f} = 34.0$ Hz), 131.8, 128.4, 126.8, 125.0, 124.5, 123.1 (q, $J_{c-f} = 273.3$ Hz), 122.8, 121.8, 119.0, 118.6, 111.4, 110.3, 98.1, 66.7, 50.3, 50.0. HRMS (ESI, m/z) calcd for $C_{25}H_{19}F_6N_3O_2$ ($[M + H]^+$): 508.1460. Found: 508.1459.

Ethyl (E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-pyrrolo[2,3-b]pyridin-3-yl)-2-cyanoacrylate (JXL082). 1H NMR (500 MHz, $CDCl_3$) δ 8.64 (s, 1H), 8.51 (s, 1H), 8.48 (dd, $J = 4.6, 1.2$ Hz, 1H), 8.21 (dd, $J = 7.9, 1.2$ Hz, 1H), 7.83 (s, 1H), 7.77 (s, 2H), 7.34 (dd, $J = 7.9, 4.6$ Hz, 1H), 5.69 (s, 2H), 4.38 (q, $J = 7.1$ Hz, 2H), 1.40 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 163.3, 147.6, 145.7, 145.0, 138.5, 132.7, 132.3 (q, $J_{c-f} = 33.6$ Hz), 127.9, 127.7, 123.1 (q, $J_{c-f} = 273.3$ Hz), 122.5, 120.3, 119.0, 117.6, 109.4, 97.0, 62.3, 48.3, 14.3. HRMS (ESI, m/z) calcd for $C_{22}H_{15}F_6N_3O_2$ ($[M + H]^+$): 468.1147. Found: 468.1146.

Ethyl (Z)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyano-3-hydroxyacrylate (JXL083). 1H NMR (500 MHz, $CDCl_3$) δ 14.61 (s, 1H), 8.67 (s, 1H), 8.31 (dd, $J = 7.0, 1.3$ Hz, 1H), 7.84 (s, 1H), 7.57 (s, 2H), 7.33 (m, 2H), 7.21 (d, $J = 7.4$ Hz, 1H), 5.53 (s, 2H), 4.40 (q, $J = 7.1$ Hz, 2H), 1.41 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 179.1, 172.2, 138.1, 136.0, 135.7, 132.7 (q, $J_{c-f} = 33.6$ Hz), 126.9, 124.5, 123.7, 123.6, 122.6, 122.9 (q, $J_{c-f} = 273.4$ Hz), 121.8, 118.3, 110.1, 109.6, 73.1, 62.3, 50.4, 14.3. HRMS (ESI, m/z) calcd for $C_{23}H_{16}F_6N_2O_3$ ($[M + H]^+$): 483.1143. Found: 483.1145.

Ethyl (Z)-3-Acetoxy-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylate (JXL084). ¹H NMR (500 MHz, CDCl₃) δ 8.56 (s, 1H), 7.90 (m, 1H), 7.85 (s, 1H), 7.61 (s, 2H), 7.32 (m, 2H), 7.24 (m, 1H), 5.51 (s, 2H), 4.29 (q, J = 7.1 Hz, 2H), 2.48 (s, 3H), 1.36 (t, J = 7.1 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 166.9, 161.4, 137.7, 136.2, 135.5, 132.8 (q, J_{c-f} = 33.9 Hz), 127.0, 126.7, 124.6, 123.6, 122.8 (q, J_{c-f} = 273.4 Hz), 122.7, 121.8, 117.7, 110.7, 109.8, 89.3, 61.8, 50.5, 29.7, 21.3, 14.2. HRMS (ESI, m/z) calcd for C₂₅H₁₈F₆N₂O₄ ([M + H]⁺): 525.1249. Found: 525.1233.

Ethyl (Z)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-3-chloro-2-cyanoacrylate (JXL085). ¹H NMR (500 MHz, CDCl₃) δ 7.91 (s, 1H), 7.85 (s, 1H), 7.77 (m, 1H), 7.65 (s, 2H), 7.63 (s, 1H), 7.30 (m, 2H), 5.48 (s, 2H), 4.18 (q, J = 7.1 Hz, 2H), 1.18 (t, J = 7.1 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 161.0, 155.6, 138.1, 136.2, 135.4, 133.4, 132.7 (q, J_{c-f} = 33.7 Hz), 127.1, 126.6, 124.5, 124.3, 122.8 (q, J_{c-f} = 273.4 Hz), 121.5, 115.8, 122.1, 110.4, 102.5, 62.4, 50.0, 13.9. HRMS (ESI, m/z) calcd for C₂₃H₁₅ClF₆N₂O₂ ([M + H]⁺): 501.0804. Found: 501.0803.

Diethyl (E)-(2-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-1-cyanovinyl)phosphonate (JXL086). ¹H NMR (500 MHz, CDCl₃) δ 8.55 (s, 1H), 8.33 (d, J = 19.7 Hz, 1H), 7.86 (d, J = 7.9 Hz, 1H), 7.83 (s, 1H), 7.58 (s, 2H), 7.31 (m, 2H), 7.22 (m, 1H), 5.54 (s, 2H), 4.21 (m, 4H), 1.39 (t, J = 7.0 Hz, 6H). ¹³C NMR (126 MHz, CDCl₃) δ 149.8, 138.3, 135.8, 132.7 (q, J_{c-f} = 33.7 Hz), 132.6, 128.2, 126.8, 124.5, 123.0, 122.9 (q, J_{c-f} = 273.4 Hz), 122.5, 119.0, 117.8 (d, J_{c-p} = 11.3 Hz), 112.2 (d, J_{c-p} = 18.9 Hz), 110.3, 91.6 (d, J_{c-p} = 207.9 Hz), 63.2, 50.4, 16.3. HRMS (ESI, m/z) calcd for C₂₄H₂₁F₆N₂O₃P ([M + H]⁺): 531.1272. Found: 531.1274.

Ethyl (E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-chloro-1H-pyrrolo[2,3-b]pyridin-3-yl)-2-cyanoacrylate (JXL087). ¹H NMR (500 MHz, CDCl₃) δ 9.22 (s, 1H), 8.77 (s, 1H), 8.32 (d, J = 5.2 Hz, 1H), 7.83 (s, 1H), 7.77 (s, 2H), 7.31 (d, J = 5.2 Hz, 1H), 5.67 (s, 2H), 4.37 (q, J = 7.1 Hz, 2H), 1.39 (t, J = 7.1 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 163.0, 148.5, 146.2, 145.4, 138.1, 137.4, 133.1, 132.4 (q, J_{c-f} = 33.7 Hz), 128.0, 122.9 (q, J_{c-f} = 273.4 Hz), 122.6, 121.8, 117.6, 117.2, 109.7, 97.8, 62.4, 48.6, 14.3. HRMS (ESI, m/z) calcd for C₂₂H₁₄ClF₆N₃O₂ ([M + H]⁺): 502.0757. Found: 502.0756.

Ethyl (E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-bromo-1H-pyrrolo[2,3-b]pyridin-3-yl)-2-cyanoacrylate (JXL088). ¹H NMR (500 MHz, CDCl₃) δ 9.40 (s, 1H), 8.78 (s, 1H), 8.22 (d, J = 5.1 Hz, 1H), 7.83 (s, 1H), 7.77 (s, 2H), 7.49 (d, J = 5.1 Hz, 1H), 5.67 (s, 2H), 4.37 (q, J = 7.1 Hz, 2H), 1.39 (t, J = 7.1 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 163.1, 148.1, 145.7, 145.1, 138.1, 137.4, 133.4, 132.4 (q, J_{c-f} = 33.7 Hz), 123.7, 122.9 (q, J_{c-f} = 273.4 Hz), 122.6, 119.7, 117.6, 110.0, 101.4, 97.4, 62.4, 48.6, 14.3. HRMS (ESI, m/z) calcd for C₂₂H₁₄BrF₆N₃O₂ ([M + H]⁺): 546.0252. Found: 546.0238.

Ethyl (E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-5-chloro-1H-indol-3-yl)-2-cyanoacrylate (JXL089). ¹H NMR (500 MHz, CDCl₃) δ 8.60 (s, 1H), 8.50 (s, 1H), 7.85 (s, 1H), 7.83 (s, 1H), 7.55 (s, 2H), 7.28 (dd, J = 8.7, 1.7 Hz, 1H), 7.14 (d, J = 8.7 Hz, 1H), 5.54 (s, 2H), 4.38 (q, J = 7.1 Hz, 2H), 1.40 (t, J = 7.1 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 163.2, 144.8, 137.7, 134.3, 134.1, 132.8 (q, J_{c-f} = 33.7 Hz), 129.7, 129.3, 126.7, 125.1, 122.8 (q, J_{c-f} = 273.4 Hz), 122.7, 118.9, 117.7, 111.6, 110.6, 96.9, 62.3, 50.6, 14.3. HRMS (ESI, m/z) calcd for C₂₃H₁₅ClF₆N₂O₂ ([M + H]⁺): 501.0804. Found: 501.0801.

Ethyl (E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-cyano-1H-indol-3-yl)-2-cyanoacrylate (JXL090). ¹H NMR (500 MHz, CDCl₃) δ 9.29 (s, 1H), 8.78 (s, 1H), 7.87 (s, 1H), 7.68 (dd, J = 7.4, 0.8 Hz, 1H), 7.56 (s, 2H), 7.49 (dd, J = 8.4, 0.8 Hz, 1H), 7.38 (dd, J = 8.4, 7.4 Hz, 1H), 5.62 (s, 2H), 4.38 (q, J = 7.1 Hz, 2H), 1.40 (t, J = 7.1 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 162.6, 144.3, 137.3, 136.2, 135.1, 132.9 (q, J_{c-f} = 33.7 Hz), 129.4, 127.3, 126.7, 124.1, 123.0, 122.8 (q, J_{c-f} = 273.4 Hz), 117.9, 117.5, 115.4, 110.7, 103.5, 98.6, 62.4, 50.6, 14.3. HRMS (ESI, m/z) calcd for C₂₄H₁₅F₆N₃O₂ ([M + H]⁺): 492.1147. Found: 492.1149.

Methyl (E)-1-(3,5-Bis(trifluoromethyl)benzyl)-3-(2-cyano-3-ethoxy-3-oxoprop-1-en-1-yl)-1H-indole-4-carboxylate (JXL091). ¹H NMR (500 MHz, CDCl₃) δ 9.36 (s, 1H), 8.71 (s, 1H), 7.93 (dd, 1H, J = 7.4, 1.1 Hz, 1H), 7.85 (s, 1H), 7.55 (s, 2H), 7.40 (dd, J = 8.3, 1.1 Hz, 1H), 7.34 (dd, J = 8.3, 7.4 Hz, 1H), 5.58 (s, 2H), 4.37 (q,

J = 7.1 Hz, 2H), 4.04 (s, 3H), 1.40 (t, J = 7.1 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 167.8, 163.5, 150.1, 137.8, 137.2, 134.9, 132.8 (q, J_{c-f} = 33.7 Hz), 126.7, 125.8, 125.1, 123.6, 122.8 (q, J_{c-f} = 273.4 Hz), 122.7, 121.7, 118.0, 114.6, 111.3, 96.4, 62.1, 52.6, 50.5, 14.3. HRMS (ESI, m/z) calcd for C₂₅H₁₈F₆N₂O₄ ([M + H]⁺): 525.1249. Found: 525.1238.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-(methoxycarbonyl)-1H-indol-3-yl)-2-cyanoacrylic Acid (JXL092). ¹H NMR (500 MHz, CDCl₃) δ 9.09 (s, 1H), 8.58 (s, 1H), 7.75 (d, J = 7.5 Hz, 1H), 7.68 (s, 1H), 7.47 (s, 2H), 7.34 (d, J = 8.2 Hz, 1H), 7.20 (app. t, J = 7.9 Hz, 1H), 5.51 (s, 2H), 3.87 (s, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 167.8, 165.1, 149.5, 138.2, 137.2, 134.8, 132.3 (q, J_{c-f} = 33.7 Hz), 126.8, 126.2, 125.5, 124.8, 123.3, 122.8 (q, J_{c-f} = 273.4 Hz), 122.3, 118.2, 114.7, 110.9, 97.2, 52.4, 50.2. HRMS (ESI, m/z) calcd for C₂₃H₁₄F₆N₂O₄ ([M + H]⁺): 497.0936. Found: 497.0939.

Ethyl (E)-3-(4-Chloro-1-phenyl-1H-indol-3-yl)-2-cyanoacrylate (JXL093). ¹H NMR (500 MHz, CDCl₃) δ 9.51 (s, 1H), 8.78 (s, 1H), 7.59 (m, 2H), 7.51 (m, 3H), 7.40 (dd, J = 8.3, 0.8 Hz, 1H), 7.32 (dd, J = 7.7, 0.8 Hz, 1H), 7.22 (app. t, J = 8.0 Hz, 1H), 4.38 (q, J = 7.1 Hz, 2H), 1.40 (t, J = 7.1 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 163.6, 147.6, 138.1, 137.4, 134.2, 130.1, 129.0, 126.9, 125.3, 124.6, 124.5, 124.3, 118.0, 111.8, 110.6, 96.0, 62.1, 14.3. HRMS (ESI, m/z) calcd for C₂₀H₁₅ClN₂O₂ ([M + H]⁺): 351.0900. Found: 351.0900.

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylamide (JXL094). ¹H NMR (500 MHz, CDCl₃) δ 8.70 (s, 1H), 8.47 (s, 1H), 7.92 (d, J = 7.1 Hz, 1H), 7.85 (s, 1H), 7.58 (s, 2H), 7.34 (m, 2H), 7.23 (m, 1H), 5.55 (s, 2H). ¹³C NMR (126 MHz, CDCl₃) δ 163.0, 144.8, 138.1, 136.0, 132.7 (q, J_{c-f} = 33.7 Hz), 132.5, 128.6, 126.8, 126.1, 124.6, 123.1, 122.9 (q, J_{c-f} = 273.4 Hz), 122.6, 119.3, 111.3, 110.3, 96.1, 50.4. HRMS (ESI, m/z) calcd for C₂₁H₁₃F₆N₃O ([M + H]⁺): 438.1041. Found: 438.1046.

Diethyl (E)-(2-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-pyrrolo[2,3-b]pyridin-3-yl)-1-cyanovinyl)phosphonate (JXL095). ¹H NMR (500 MHz, CDCl₃) δ 8.58 (s, 1H), 8.46 (dd, J = 4.7, 1.4 Hz, 1H), 8.24 (d, J = 19.5 Hz, 1H), 8.20 (dd, J = 8.0, 1.4 Hz, 1H), 7.82 (s, 1H), 7.77 (s, 2H), 7.32 (dd, J = 8.0, 4.7 Hz, 1H), 5.67 (s, 2H), 4.22 (m, 4H), 1.40 (t, J = 7.1 Hz, 6H). ¹³C NMR (126 MHz, CDCl₃) δ 149.2 (d, J_{c-p} = 8.2 Hz), 147.4, 145.6, 138.6, 132.4 (q, J_{c-f} = 33.7 Hz), 132.0, 127.9, 127.8, 123.0 (q, J_{c-f} = 273.4 Hz), 122.4, 119.9, 118.9, 117.4 (d, J_{c-p} = 11.3 Hz), 110.5 (d, J_{c-p} = 19.5 Hz), 92.8 (d, J_{c-p} = 205.1 Hz), 63.4, 48.3, 16.3. HRMS (ESI, m/z) calcd for C₂₃H₂₀F₆N₃O₃P ([M + H]⁺): 532.1225. Found: 532.1238.

(E)-(2-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-1-cyanovinyl)phosphonic Acid (JXL096). ¹H NMR (500 MHz, CD₃OD) δ 8.58 (s, 1H), 8.25 (d, J = 19.6 Hz, 1H), 7.90 (s, 1H), 7.87 (m, 1H), 7.76 (s, 2H), 7.45 (m, 1H), 7.31 (m, 2H), 5.75 (s, 2H). ¹³C NMR (126 MHz, CD₃OD) δ 146.7 (J_{c-p} = 7.2 Hz), 140.2, 136.1, 132.1, 131.9 (q, J_{c-f} = 33.7 Hz), 128.0, 127.2, 123.8, 122.3, 123.2 (q, J_{c-f} = 273.4 Hz), 121.4, 118.2, 117.4 (d, J_{c-p} = 11.3 Hz), 111.5 (J_{c-p} = 18.4 Hz), 110.5, 94.6 (d, J_{c-p} = 201.2 Hz), 49.1. HRMS (ESI, m/z) calcd for C₂₀H₁₃F₆N₂O₃P ([M + H]⁺): 475.0646. Found: 475.0642.

In Vitro Evaluation of the Analogues' Activity in Promoting Lactate Production Rate. Human breast epithelial (MCF10A) cells were obtained from the American Type Culture Collection and cultured in Dulbecco's modified Eagle medium (DMEM):F12 media supplemented with 5% of horse serum, 50 U/mL of penicillin-streptomycin, 10 μg/mL of insulin, 0.5 μg/mL of hydrocortisone, 20 ng/mL of EGF, and 10 μg/mL of cholera toxin. To determine whether the analogues could promote cellular lactate production, MCF10A cells were treated with analogues and lactate concentrations were measured using a BioProfile Basic Analyzer (Nova Biomedical). Briefly, MCF10A cells were seeded in 6-well plates in triplicate, treated the following day with 10 μM analogue, or an equivalent volume of DMSO, for 24–30 h. Culture media lactate concentrations were measured and normalized to cell number and duration of the experiment to calculate a cellular lactate production rate (nmol lactate per million cells per hour). Ordinary one-way ANOVA followed by Dunnett's post-test was performed using GraphPad Prism 8.3.0, comparing each analogue to DMSO.

A Seahorse XF Analyzer (Agilent) was used to measure the oxygen consumption rate (OCR) to calculate IC_{50} values for several of the analogues. MCF10A cells were seeded in 96-well plates, treated with half-log doses of analogue, ranging from 3 nM to 10 μ M, in the presence of 5 mM of pyruvate and 0.5 mM of malate. OCR (fmol O_2 per cell per min) was measured in duplicate and normalized to cell number. IC_{50} values were calculated using Prism (GraphPad).

In Vivo Evaluation of the Selected Analogues' Activity in Accelerating Hair Growth on Mice. To determine the efficacy of the compounds on the hair cycle, wild-type C57BL/6J mice were shaved at postnatal day 50 and topically treated with the compounds disclosed herein suspended in lotion (20 μ M) every other day for up to 3 weeks. The lotion used was a Transderma Plo Ultramax gel. The mice were treated topically every other day with \sim 300 μ L at 20 μ M. Photographs were taken every week to monitor pigmentation and hair growth.

All mice were kept under defined pathogen-free conditions at the AAALAC-approved animal facility of the Division of Laboratory Animals (DLAM) at UCLA. All animal experiments were performed with the approval of the UCLA Office of Animal Resource Oversight (OARO); the approval number is ARC no. 2016-055-11A.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jmedchem.0c01570>.

High field proton and carbon NMR spectra for compounds JXL001–JXL096 (PDF)

Molecular formula strings (CSV)

■ AUTHOR INFORMATION

Corresponding Author

Michael E. Jung – Department of Chemistry and Biochemistry and Department of Molecular and Medical Pharmacology, UCLA, Los Angeles, California 90095, United States; orcid.org/0000-0003-1290-8588; Phone: (310) 825-7954; Email: jung@chem.ucla.edu

Authors

Xiaoguang Liu – Department of Chemistry and Biochemistry, UCLA, Los Angeles, California 90095, United States

Aimee A. Flores – Department of Molecular Cell and Developmental Biology, Eli and Edythe Broad Center for Regenerative Medicine, and Molecular Biology Institute, UCLA, Los Angeles, California 90095, United States

Lisa Situ – Department of Molecular Cell and Developmental Biology and Department of Biological Chemistry, UCLA, Los Angeles, California 90095, United States

Wen Gu – Department of Molecular and Medical Pharmacology, UCLA, Los Angeles, California 90095, United States

Hui Ding – Department of Chemistry and Biochemistry, UCLA, Los Angeles, California 90095, United States

Heather R. Christofk – Department of Molecular and Medical Pharmacology, Jonsson Comprehensive Cancer Center, Eli and Edythe Broad Center for Regenerative Medicine, and Department of Biological Chemistry, UCLA, Los Angeles, California 90095, United States

William E. Lowry – Department of Molecular Cell and Developmental Biology, Jonsson Comprehensive Cancer Center, Eli and Edythe Broad Center for Regenerative Medicine, and Molecular Biology Institute, UCLA, Los Angeles, California 90095, United States

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.jmedchem.0c01570>

Author Contributions

The manuscript was written and edited by Drs. X.L. and M.E.J. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Dr. X.L. is a postdoctoral fellow supported by the UCLA Tumor Cell Biology Training Program (USHHS Ruth L. Kirschstein Institutional National Research Service Award no. T32 CA009056). We thank Linsey Stiles at the UCLA Mitochondria Core for her help with the Seahorse assays.

■ ABBREVIATIONS

MPC, mitochondrial pyruvate carrier; LDH, lactate dehydrogenase; TZD, thiazolidinedione; OCR, oxygen consumption rate; DMSO, dimethyl sulfoxide

■ REFERENCES

- (1) Vary, J. C., Jr. Selected Disorders of Skin Appendages-Acne, Alopecia, Hyperhidrosis. *Med. Clin. North Am.* **2015**, *99*, 1195–1211.
- (2) Rogers, N. E.; Avram, M. R. Medical Treatments for Male and Female Pattern Hair Loss. *J. Am. Acad. Dermatol.* **2008**, *59*, 547–566.
- (3) Stenn, K. S.; Paus, R. Controls of Hair Follicle Cycling. *Physiol. Rev.* **2001**, *81*, 449–494.
- (4) Hsu, Y. C.; Pasolli, H. A.; Fuchs, E. Dynamics Between Stem Cells, Niche, and Progeny in The Hair Follicle. *Cell* **2011**, *144*, 92–105.
- (5) Fuchs, E. The Tortoise and The Hair: Slow-Cycling Cells in The Stem Cell Race. *Cell* **2009**, *137*, 811–819.
- (6) Fuchs, E.; Merrill, B. J.; Jamora, C.; DasGupta, R. At The Roots of A Never-Ending Cycle. *Dev. Cell* **2001**, *1*, 13–25.
- (7) Flores, A.; Schell, J.; Krall, A. S.; Jelinek, D.; Miranda, M.; Grigorian, M.; Braas, D.; White, A. C.; Zhou, J. L.; Graham, N. A.; Graeber, T.; Seth, P.; Evseenko, D.; Collier, H. A.; Rutter, J.; Christofk, H. R.; Lowry, W. E. Lactate Dehydrogenase Activity Drives Hair Follicle Stem Cell Activation. *Nat. Cell Biol.* **2017**, *19*, 1017–1026.
- (8) Herzig, S.; Raemy, E.; Montessuit, S.; Veuthey, J. L.; Zamboni, N.; Westermann, B.; Kunji, E. R. S.; Martinou, J. C. Identification and Functional Expression of The Mitochondrial Pyruvate Carrier. *Science* **2012**, *337*, 93–96.
- (9) Bricker, D. K.; Taylor, E. B.; Schell, J. C.; Orsak, T.; Boutron, A.; Chen, Y. C.; Cox, J. E.; Cardon, C. M.; Van Vranken, J. G.; Dephoure, N.; Redin, C.; Boudina, S.; Gygi, S. P.; Brivet, M.; Thummel, C. S.; Rutter, J. A Mitochondrial Pyruvate Carrier Required for Pyruvate Uptake in Yeast, Drosophila, and Humans. *Science* **2012**, *337*, 96–100.
- (10) Halestrap, A. P. The Mechanism of the Inhibition of The Mitochondrial Pyruvate Transporter by α -Cyanocinnamate Derivatives. *Biochem. J.* **1976**, *156*, 181–183.
- (11) Shearman, M. S.; Halestrap, A. P. The Concentration of the Mitochondrial Pyruvate Carrier in Rat Liver and Heart Mitochondria Determined with α -cyano- β -(1-phenylindol-3-yl)acrylate. *Biochem. J.* **1984**, *223*, 673–676.
- (12) Zhong, Y.; Li, X.; Yu, D.; Li, X.; Li, Y.; Long, Y.; Yuan, Y.; Ji, Z.; Zhang, M.; Wen, J. G.; Nesland, J. M.; Suo, Z. Application of Mitochondrial Pyruvate Carrier Blocker UK-5099 Creates Metabolic Reprogram and Greater Stem-Like Properties in LnCap Prostate Cancer Cells in vitro. *Oncotarget* **2015**, *6*, 37758–37769.
- (13) Divakaruni, A. S.; Wiley, S. E.; Rogers, G. W.; Andreyev, A. Y.; Petrosyan, S.; Loviscach, M.; Wall, E. A.; Yadava, N.; Heuck, A. P.; Ferrick, D. A.; Henry, R. R.; McDonald, W. G.; Colca, J. R.; Simon,

M. I.; Ciaraldi, T. P.; Murphy, A. N. Thiazolidinediones Are Acute, Specific Inhibitors of The Mitochondrial Pyruvate Carrier. *Proc. Natl. Acad. Sci. U. S. A.* **2013**, *110*, 5422–5427.