

CHEMISTRY  
XL-14A

MOLECULAR  
SHAPE AND  
STRUCTURE  
PART II




July 30, 2011

Robert lafe

# Midterm Results

2

## Histogram of Midterm #1



Average: 55.8

StDev: 20.3

High: 91.5

Low: 8

# Midterm 1

3

1. How many seconds are in 65 years?

- a)  $3.3 \times 10^7$  sec
- b)  $3.4 \times 10^7$  sec
- c)  $2.0 \times 10^9$  sec
- d)  $2.1 \times 10^9$  sec

2. The molecular formula of hydroquinone is  $C_6H_6O_2$ .  
What is the empirical formula of hydroquinone?

- a)  $C_9H_9O_3$
- b)  $C_6H_6O_2$
- c)  $C_3H_3O$
- d) CHO

# Midterm 1

4

3. For the following set of four quantum numbers ( $n, l, m_l, m_s$ ) identify the set that is not valid:

- a)  $\{9, 5, -3, 1/2\}$
- b)  $\{3, 3, 3, 1/2\}$
- c)  $\{2, 0, 0, 1/2\}$
- d)  $\{4, 2, -2, 1/2\}$

4. How many electrons are in principle quantum number  $n = 3$ ?

- a) 3
- b) 6
- c) 9
- d) 18



# Midterm 1

5

5. How many radial nodes are in a 3d orbital?

- a) 0
- b) 1
- c) 2
- d) 3

6. Which of the following is isoelectronic with  $\text{N}^{3-}$ ?

- a) F
- b)  $\text{Cl}^{-1}$
- c)  $\text{Mg}^{2+}$
- d) Ar

# Midterm 1

6

7. Which of the compounds below has bonds with more covalent character?

- a) CaO
- b) Li<sub>2</sub>O
- c) MgO
- d) MgS

8. Which of the following is the most electronegative?

- a) B
- b) In
- c) Te
- d) O

# Midterm 1

7

9. Which of the following has the largest atomic radius?

- a) Be
- b) S
- c) Te
- d) Sr

# Midterm 1

8

10. 1,3-Benzodioxol-5-ol, otherwise known as sesamol, is a natural organic compound found in sesame seed oil. The combustion analysis of a 100 mg sample of 1,3-benzodioxol-5-ol shows its composition is: 60.87% C, 4.38% H, and 34.75% O.

- a) What is the empirical formula of 1,3-benzodioxol-5-ol?
- b) If the molecular weight of 1,3-benzodioxol-5-ol is 138.12 g/mol, what is the molecular formula?

# Midterm 1

11. Sodium metal reacts readily with water to produce hydrogen gas and sodium hydroxide in solution.

- a) Write a balanced reaction.
- b) If  $3.00 \times 10^{-1}$  grams of sodium react with 0.500 L of water (density = 1 g/mL), what is the theoretical yield of hydrogen gas?
- c) If exactly  $1.20 \times 10^{-2}$  grams of hydrogen gas is collected, what is the percent yield?
- d) Knowing the percent yield of the reaction, what is the predicted concentration of NaOH in the solution assuming the volume of water does not change?

# Midterm 1

10

12. Write the noble gas electron configuration for the following atoms/ions. How many *unpaired* electrons does each have?

- a) P
- b)  $\text{Ag}^{+1}$

13. Identify the  $M^{3+}$  ion for each ground state electron configuration

- a)  $[\text{Ar}]3d^7$
- b)  $[\text{Ar}]3d^{10}$

14. Briefly explain why oxygen has a lower first ionization energy than nitrogen.

# Midterm 1

11

15. Calculate the wavelength of the radiation emitted by a hydrogen atom (one electron atom) when an electron makes a transition from  $n = 5$  to  $n = 2$ .
16. The work function for chromium metal is  $4.37 \text{ eV}$  or  $7.00 \times 10^{-19} \text{ J}$ . Light with a wavelength of  $11.0 \text{ nm}$  is shined on the metal. What is the velocity of the ejected electron?

# Midterm 1

12

17. Draw the boundary surface for the three 3p orbitals, making sure to name and label each axis correctly. How many radial nodes does a 3p orbital have?



# Midterm 1

13

- Provide the Lewis dot structure for the following compounds. Make sure every electron is drawn. Indicate atoms with formal charge. If resonance structures can be drawn, draw all valid structures. Make sure to indicate formal charge when necessary.
- a)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
  - b)  $\text{CN}^-$
  - c)  $\text{HNO}_3$
  - d)  $\text{POCl}_3$

# Valence shell electron pair repulsion theory (VSEPR)

14

- Lewis Theory
  - Connectivity, electron tracking
- VSEPR Theory
  - 3-D Structure around an atom
- Valence Bond Theory
- Molecular Orbital Theory

# Review – VSEPR Theory

15


## Valence-Shell Electron-Pair Repulsion (VSEPR)

- Accounts for 3D shapes of molecules
- Based on electron-electron repulsion
- Determine bond angles → shape

Rules are based on experimental observation:


1. Areas of electron concentration (bonds and lone pairs) around the central atom repel each other.
2. Bonds and lone pairs stay as far away from each other as possible (without changing distance)

2




Linear

3




Trigonal planar

4




Tetrahedral

5




Trigonal bipyramidal


6




Octahedral

AX<sub>2</sub>E

Bent (V shaped)

Examples: SO<sub>2</sub>, O<sub>3</sub>, PbCl<sub>2</sub>, SnBr<sub>2</sub>AX<sub>3</sub>E


Trigonal pyramidal

Examples: NH<sub>3</sub>, PF<sub>3</sub>, ClO<sub>3</sub>, H<sub>3</sub>O<sup>+</sup>AX<sub>2</sub>E<sub>2</sub>


Bent (V shaped)

Examples: H<sub>2</sub>O, OF<sub>2</sub>, SCl<sub>2</sub>AX<sub>4</sub>E


Seesaw

Examples: SF<sub>4</sub>, XeO<sub>2</sub>F<sub>2</sub>, IF<sub>4</sub><sup>+</sup>, IO<sub>2</sub>F<sub>2</sub><sup>-</sup>AX<sub>3</sub>E<sub>2</sub>


T-shaped

Examples: ClF<sub>3</sub>, BrF<sub>3</sub>AX<sub>2</sub>E<sub>3</sub>

Linear

Examples: XeF<sub>2</sub>, I<sub>3</sub><sup>-</sup>, IF<sub>2</sub><sup>-</sup>AX<sub>5</sub>E

Square pyramidal

Examples: BrF<sub>5</sub>, TeF<sub>5</sub><sup>-</sup>, XeOF<sub>4</sub>AX<sub>4</sub>E<sub>2</sub>


Square planar

Examples: XeF<sub>4</sub>, ICl<sub>4</sub><sup>-</sup>

# Review - Polar Molecules

17

A **polar covalent bond** has a non-zero dipole moment



**Polar molecules** – molecules with non-zero dipole moment:  $m \neq 0$  D

Examples:                      HCl    HF    HBr


A non-polar molecule has no dipole moment:  $m = 0$  D

Examples:                      O<sub>2</sub>    CH<sub>4</sub>    SF<sub>6</sub>

# Polar Molecules

18

Polar bonds, Non-polar molecule?




**CO<sub>2</sub>**

If the dipole moments cancel out (Vector sum = 0):


Each C-O bond is polar

Two equally polar bonds, with opposing dipole moments




# Polar Molecules

19








Each O-H bond is polar

Overall molecule is ???????



# Polar Molecules


20






# Polar Molecules

21



**non-polar**



**polar**


# Valence Bond (VB) Theory

22

- Lewis Theory
  - ▣ Connectivity, electron tracking
- VSEPR Theory
  - ▣ 3-D Structure around an atom
- Valence Bond Theory
  - ▣ Extended 3-D Structure Information
  - ▣ Delocalization in Molecules
  - ▣ Illustrates Multiple Bonding
  - ▣ Prediction of Reactivity
- Molecular Orbital Theory






# Answers to $H \Psi = E \Psi$

23





# Answers to $H \Psi = E \Psi$

24




# Bonding With Orbitals

25



# Bonding with Orbitals

26




# Valence-Bond Theory

27

So far, we've been thinking of molecules using Lewis' Theory:

Bonding electrons are located in between bonded atoms – electrons are localized



But....

We learned in Ch. 1 that we learned to think of electrons as wavefunctions, which are described by atomic orbitals

Valence Bond Theory – quantum mechanical view of bonding

# 'Types' of Bonds – $\sigma$ bond

28

Lets start with  $H_2$ , the simplest molecule:

Ground state H has one 1s electron


When the 2 H atoms bond, the atomic orbitals merge, forming a  $\sigma$ -bond

$\sigma$ -bond (sigma bond) – along bond axis.

We say the atomic orbitals overlap

More overlap = Stronger bond

All single covalent bonds are  $\sigma$ -bonds





# 'Types' of Bonds – $\sigma$ bond

29

All single covalent bonds are  $\sigma$ -bonds

Can have  $\sigma$ -bonds between any types of orbitals:



Two s orbitals

Two p orbitals

An s and a p orbital

Etc....

Example:  $\sigma$ -bond between 1s orbital of H and  $2p_z$  orbital of F



# 'Types' of Bonds – Multiple bonds



30

What happens in  $N_2$ ?


Remember the bond in  $N_2$  is a triple bond...

Each N atom has 3 unpaired 2p electrons


Lets look at the atomic orbitals of N:



$p_x$



$p_y$



$p_z$

# 'Types' of Bonds – Multiple bonds



31

What happens in  $N_2$ ?

Remember the bond in  $N_2$  is a triple bond...

Each N atom has 3 unpaired 2p electrons

How do the atomic orbitals bond?



Between the 2p<sub>z</sub> and 2p<sub>z</sub> orbitals, we have bonding on the bond axis

This is a  $\sigma$ -bond!

# 'Types' of Bonds – Multiple bonds



32

What happens in  $N_2$ ?

Remember the bond in  $N_2$  is a triple bond...

Each N atom has 3 unpaired 2p electrons

How do the atomic orbitals bond?



Between  $2p_x$  and  $2p_x$  orbitals, the bonding is not on the bond axis



A different type of bond!

$\pi$ -bond – nodal plane along bond axis

Bonding occurs above and below the bond axis


# Bonding of N<sub>2</sub>

33



# Bonding of N<sub>2</sub>

34



# Hybridization of Orbitals

35

What about methane ( $\text{CH}_4$ )?


According to Valence Bond Theory:

C should only make 2 bonds!




But we know that C can make 4 bonds

And  $\text{CH}_4$  has a tetrahedral shape according to VSEPR:

How do we get a tetrahedral shape from the 2s and 2p orbitals?




Carbon,  $[\text{He}]2s^22p_x^12p_y^1$



# Hybridization of Orbitals

36

We solve the 4 bond problem by promoting an electron:



Carbon,  $[\text{He}]2s^1 2p_x^1 2p_y^1 2p_z^1$

It takes energy to promote an electron to a higher E orbital

But, overall the energy is lower if C can make 4 bonds instead of 2.

Ok, what about the geometry problem?



# Hybridization of Orbitals


37

By promoting an electron, we can now make 4 bonds

But the geometry of the  $p_x$ ,  $p_y$ , and  $p_z$  orbitals don't match the tetrahedral shape of  $\text{CH}_4$

$s$  and  $p$  orbitals are described by a wave-like model of the  $e$

If we think of the orbitals as interfering with each other, we can define new hybrid orbitals:




$$h_1 = s + p_x + p_y + p_z$$

$$h_2 = s - p_x - p_y + p_z$$

$$h_3 = s - p_x + p_y - p_z$$


$$h_4 = s + p_x - p_y - p_z$$



# $sp^3$ Hybridization

38


C in  $\text{CH}_4$  uses 4 hybrid orbitals:



We took 1 s orbital and 3 p orbitals to make 4  $sp^3$  orbitals.

The sum of the atomic orbitals = the sum of the hybrid orbitals

The 4  $sp^3$  orbitals point in the 4 directions  
of the tetrahedral bonds



# $sp^3$ Hybridization

39

Back to  $CH_4$ :


The 4  $2sp^3$  hybrid orbitals on carbon make  $\sigma$ -bonds with the  $1s$  orbitals of the 4 H atoms

$NH_3$  also uses  $sp^3$  hybrid orbitals:



**Ammonia,  $NH_3$**

Whenever an atom has a tetrahedral structure, it is  $sp^3$  hybridized




# $sp^3$ Hybridization

40

Whenever an atom has a tetrahedral structure, we say it is  $sp^3$  hybridized

This includes molecules with multiple central atoms:



# Atomic Orbitals → Hybrid Atomic Orbitals

41

$sp^3$  bonding works for tetrahedral-shaped molecules

What about the other VSEPR shapes?


Linear bonding can be described by  $sp$  hybridization:

$$h_1 = s + p$$

$$h_2 = s - p$$

2 of the  $p$  orbitals remain as they were

Unused  $p$  orbitals are available  
for  $p$  bonding



# Other Types of Hybridization

42

$sp^3$  bonding works for tetrahedrally shaped molecules

What about the other VSEPR shapes?


Trigonal planar can be described by  **$sp^2$  hybridization**:

$$h_1 = s + \sqrt{2}p_y$$

$$h_2 = s + \sqrt{\frac{3}{2}}p_x - \sqrt{\frac{1}{2}}p_y$$


$$h_3 = s - \sqrt{\frac{3}{2}}p_x - \sqrt{\frac{1}{2}}p_y$$

The  $p_z$  orbital is not used and remains as it was




# Hybrid Orbitals

43



# Hybrid Orbital Shapes


44






# Look Familiar?

45



sp


2



Linear

sp<sup>2</sup>


3



Trigonal planar

sp<sup>3</sup>


4



Tetrahedral

sp<sup>3</sup>d


5



Trigonal bipyramidal


sp<sup>3</sup>d<sup>2</sup>

6




Octahedral

AX<sub>2</sub>E



Examples: SO<sub>2</sub>, O<sub>3</sub>, PbCl<sub>2</sub>, SnBr<sub>2</sub>

AX<sub>3</sub>E




Examples: NH<sub>3</sub>, PF<sub>3</sub>, ClO<sub>3</sub>, H<sub>3</sub>O<sup>+</sup>

AX<sub>4</sub>E



Examples: SF<sub>4</sub>, XeO<sub>2</sub>F<sub>2</sub>, IF<sub>4</sub><sup>+</sup>, IO<sub>2</sub>F<sub>2</sub><sup>-</sup>

AX<sub>5</sub>E




Examples: BrF<sub>5</sub>, TeF<sub>5</sub><sup>-</sup>, XeOF<sub>4</sub>

AX<sub>2</sub>E<sub>2</sub>




Examples: H<sub>2</sub>O, OF<sub>2</sub>, SCl<sub>2</sub>

AX<sub>3</sub>E<sub>2</sub>




Examples: ClF<sub>3</sub>, BrF<sub>3</sub>

AX<sub>2</sub>E<sub>3</sub>



Examples: XeF<sub>2</sub>, I<sub>3</sub><sup>-</sup>, IF<sub>2</sub><sup>-</sup>

AX<sub>4</sub>E<sub>2</sub>



Examples: XeF<sub>4</sub>, ICl<sub>4</sub><sup>-</sup>

# Other Types of Hybridization

47

N atomic orbitals always produce N hybrid orbitals

**TABLE 3.2** Hybridization and Molecular Shape\*





Electron arrangement	Number of atomic orbitals	Hybridization of the central atom	Number of hybrid orbitals
linear	2	$sp$	2
trigonal planar	3	$sp^2$	3
tetrahedral	4	$sp^3$	4
trigonal bipyramidal	5	$sp^3d$	5
octahedral	6	$sp^3d^2$	6

Spectroscopic data suggests terminal atoms use hybrid orbitals as well

A terminal Cl uses  $sp^3$  hybridization in the arrangement of its lone pairs?


# Examples

48




# $\pi$ Bond

49






# Ethylene

50



# Ethylene

51



# Characteristics of Multiple Bonds

52


double bond = 1  $\sigma$ -bond + 1  $\pi$ -bond

triple bond = 1  $\sigma$ -bond + 2  $\pi$ -bonds

$\sigma$ -bonds result from head-on overlap of orbitals

$\pi$ -bonds results from side-by-side overlap

Atoms in a single bond can rotate freely, whereas atoms in a double bond are much less likely to:





# Characteristics of Multiple Bonds


53

The shape of ethylene ( $C_2H_4$ )

Experimental evidence:

All six atoms lie in the same plane with  $120^\circ$  bond angles

Suggests trigonal planar structure  $\rightarrow$   $sp^2$  hybridization



Each C:  $sp^2 + p$

s-bonds with the  $sp^2$  orbitals

C — C

4 C — H


p-bond with the leftover p orbital

C — C

# Characteristics of Multiple Bonds





54

The shape of acetylene ( $C_2H_2$ ) is linear  $\rightarrow$   $sp$  hybridization



# Hybridization and Benzene

55





# Molecular Orbital Theory

56


- Lewis Theory
  - ▣ Connectivity, electron tracking
- VSEPR Theory
  - ▣ 3-D Structure around an atom
- Valence Bond Theory
  - ▣ Extended 3-D Structure Information
  - ▣ Delocalization in Molecules (LIMITED)
  - ▣ Illustrates Multiple Bonding
  - ▣ Prediction of Reactivity
- Molecular Orbital Theory
  - ▣ Orbitals as a function of the whole molecule
  - ▣ Delocalization much more thorough
  - ▣ Anti-bonding and Non-bonding electrons
  - ▣ Behavior in a magnetic field
  - ▣ Spectral Data

# Paramagnetic / Diamagnetic

57




no applied magnetic field




applied magnetic field

=




diamagnet



# Constructive and Destructive Interference

58



# Molecular Orbitals


59

MO Theory – electrons occupy **molecular orbitals** spread over *entire* molecule

- All valence electrons are delocalized
- Molecular orbitals are built by adding together atomic orbitals
- Linear combination of atomic orbitals (LCAO)

H – H bond in H<sub>2</sub> with two 1s electrons:

$$\psi = \psi_{A1s} + \psi_{B1s}$$



- LCAO-MO shows constructive interference
- $E_{\text{LCAO-MO}} < E_{\text{AO}}$
- LCAO-MO are bigger than AO

# Molecular Orbitals

60


“Conservation of orbitals”

H<sub>2</sub> bonds with two 1s orbitals – it must have 2 LCAO-MOs

2<sup>nd</sup> LCAO-MO - destructive interference,  
higher PE than atomic orbitals


Nodal plane! – much less e- density  
“internuclearly”

This is an antibonding orbital



$$\psi = \psi_{A1s} + \psi_{B1s}$$

$$\psi = \psi_{A1s} - \psi_{B1s}$$




1<sup>st</sup> LCAO-MO is lower in energy than Atomic orbitals

This is called a bonding orbital




# Sigma Bonds

61



# Molecular Orbital Energy Diagram

62



Shows:

Relative Energies of

Atomic Orbitals

Bonding MOs

Antibonding MOs

Shapes of LCAO-MOs


Type of MO – s or p

Molecular Electron  
Configuration

# Electron Configurations of Diatomic Molecules

63

Shows **ALL valence electrons** using the Building-Up Principle:




Use “Hund’s Rule just like in atomic orbitals!

Electron configuration of  $\text{H}_2 \rightarrow s_{1s}^2$

Based on MO theory – a single electron can hold a bond together

# Bond Order

64



Valence Bond Theory:

Bond order was # bonding pairs

MO Theory, bond order is:

$$\square b = \frac{1}{2} (N - N^*)$$

$N = \# e^-$ s in bonding orbitals

$\square N^* = \# e^-$ s in antibonding orbitals

$\square H_2 \rightarrow b = 1$

$\square He_2 \rightarrow b = 0$

↑ energy

# 2<sup>nd</sup> row Diatomic Molecules

65


Must include both 2s and 2p orbitals in making our MOs

Only orbitals that are close in energy will form MOs

We have 2 (2s) orbitals and 6 (2p) orbitals → 8 atomic orbitals

2s orbitals overlap in the same way as the 1s orbitals of H<sub>2</sub>

2p orbitals overlap in same orientations as in Valence bond theory:





# 2<sup>nd</sup> row Diatomic Molecules

66

We have 2 (2s) orbitals and 6 (2p) orbitals → 8 atomic orbitals





2s orbitals overlap in the same way as the 1s orbitals of H<sub>2</sub>

2p orbitals overlap in same orientations as in Valence bond theory:





# Pi Bonds

67



# 2<sup>nd</sup> row Homonuclear Diatomic Molecules


68






# 2<sup>nd</sup> row Homonuclear Diatomic Molecules

69




- Homonuclear Diatomic MO Energy diagram for all elements right of (and including) oxygen



And their combinations

# 2<sup>nd</sup> row Homonuclear Diatomic Molecules

70




□ Homonuclear Diatomic MO Energy diagram for all elements left of oxygen

$\text{Li}_2$     $\text{Be}_2$     $\text{B}_2$     $\text{C}_2$     $\text{N}_2$


# Relative MO Energy Levels

71




# MO Diagram for Li<sub>2</sub>

72




# MO Diagram for Be<sub>2</sub>

73




# MO Diagram for N<sub>2</sub>

74




# MO Diagram for O<sub>2</sub>

75




# MO Diagram for F<sub>2</sub>

76








bond order = (electrons in bonding MOs – electrons in antibonding MOs) / 2

1	0	1	2	3	2	1	0
$\text{Li-Li}$	$\text{Be-Be}$	$\text{B-B}$	$\text{C=C}$	$\text{N}\equiv\text{N}$	$\text{O=O}$	$\text{F-F}$	$\text{Ne-Ne}$
known in the gas phase	dimer unknown	known in the gas phase paramagnetic diradical	known in the gas phase at high temp.	stable gas	stable gas paramagnetic diradical. Singlet & triplet states	stable gas	dimer unknown


# Relative AO Energy Levels

78



# Heteronuclear Diatomic Molecules

79




Account for  $\Delta EN$   
between AO's

More electronegative  
atoms will have lower  
energy atomic orbitals

Contributions are  
dependent on energy  
level

# MO diagram for HF

80





# Orbitals in Polyatomic Molecules

81


An electron pair in a bonding orbital helps to bind together the whole molecule – can become very complex

Occasionally some MOs are neither bonding nor antibonding –these are called nonbonding orbitals



# MO Diagram for Carbon Monoxide

82



# Organic Molecules

83

Chemists commonly mix VB and MO theories when discussing organic molecules:

VB theory is good for talking about  $\sigma$  bonds

MO theory is better for considering  $\pi$  bonds


Consider Benzene ( $C_6H_6$ ):

VB perspective – each C is  $sp^2$  hybridized

$\sigma$  framework is overlap of  $sp^2$  orbitals


MO perspective – conjugation of  $\pi$  bonds extremely important

$\pi$  MOs account for delocalization of electrons



# $\pi$ Orbitals of Benzene


84





# Coordination Compounds

85




**FIGURE 16.17** When potassium cyanide is added to a solution of iron(II) sulfate, the cyanide ions replace the  $\text{H}_2\text{O}$  ligands of the  $[\text{Fe}(\text{OH}_2)_6]^{2+}$  complex (left) and produce a new complex, the hexacyanoferrate(II) ion,  $[\text{Fe}(\text{CN})_6]^{4-}$  (right). The blue color is due to the polymeric compound called *Prussian blue*, which forms from the cyanoferrate ion.

# Colors of Coordination Compounds

86

Aqueous solutions of  $[\text{Fe}(\text{SCN})(\text{OH}_2)_5]^{2+}$ ,  
 $[\text{Co}(\text{SCN})_4(\text{OH}_2)_2]^{2-}$ ,  $[\text{Cu}(\text{NH}_3)_4(\text{OH}_2)_2]^{2+}$ , and  $[\text{CuBr}_4]^{2-}$




# Biological Importance


87

## □ Metalloproteins`

cobalamin (vitamin B<sub>12</sub>)




Hemoglobin



# Coordination Compounds


88

- ❑ Metals in the d-block form coordination compounds
- ❑ Coordination compound – electrically neutral compound in which at least one of the ions present is a complex
- ❑ Ligands – ions or molecules that are attached to the central metal atom or ion
- ❑ Coordination number – number of ligands attached to the central metal




# Shapes of Complexes


89



1 An octahedral complex




2 A tetrahedral complex




3 A square-planar complex

# Shapes of Complexes

90



4 Dimethylmercury(0)



5 Ferrocene,  $\text{Fe}(\text{C}_5\text{H}_5)_2$

Sandwich compound

# Nomenclature

91

- $[\text{FeCl}(\text{OH}_2)_5]^+$  pentaquachloridoiron(II)
  - ▣ Overall complex charge = +1
  - ▣ Ligand charge =  $\text{Cl}^{-1}$  and  $\text{OH}_2$  (water)
  - ▣ Metal charge + ligand charge = overall complex charge
    - $\text{Fe} + (-1) = +1$
    - $\text{Fe} = +2$
  
- $[\text{Co}(\text{NH}_3)_3(\text{OH}_2)_3]_2(\text{SO}_4)_3$  triamminetriaquacobalt(III) sulfate
  - ▣ Overall complex charge = +3
  - ▣ Ligand charge = 0
  - ▣  $\text{Co} = +3$

# Common Ligands

92





Formula*	Name
<b>Neutral ligands</b>	
$\text{OH}_2$	aqua
$\text{NH}_3$	ammine
$\text{NO}$	nitrosyl
$\text{CO}$	carbonyl
$\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$	ethylenediamine (en) <sup>†</sup>
$\text{NH}_2\text{CH}_2\text{CH}_2\text{NHCH}_2\text{CH}_2\text{NH}_2$	diethylenetriamine (dien) <sup>‡</sup>



# Common Ligands

93

## Anionic ligands



fluorido

chlorido

bromido

iodido

hydroxido

oxido

cyanido- $\kappa C$

isocyanido, cyanido- $\kappa N$

thiocyanato- $\kappa N$

isothiocyanato, thiocyanato- $\kappa S$

nitrito- $\kappa O$

nitro, nitrito- $\kappa N$

carbonato- $\kappa O$

oxalato (ox)<sup>†</sup>

ethylenediaminetetraacetato (edta)<sup>§</sup>

sulfato

# Prefixes

94

- 1 mono
- 2 di (bis)
- 3 tri (tris)
- 4 tetra (tetrakis)
- 5 penta (pentakis)
- 6 hexa (hexakis)
- 7 hepta
- 8 octa

Prefixes in parentheses are used when the ligand already contains a Greek prefix or is polydentate

Example: ethylenediamine  
                  oxalato     (bidentate)

# More Nomenclature


95

- $\text{Na}_2[\text{PtCl}_2(\text{ox})_2]$  sodium dichloridobis(oxalato)platinate(IV)
  - ▣ Overall complex charge = -2
  - ▣ Ligand charge =  $\text{Cl}^{-1}$  and oxalato  $\text{C}_2\text{O}_4^{2-}$
  - ▣ Metal charge + ligand charge = overall complex charge
    - $\text{Pt} + (-6) = -2$
    - $\text{Pt} = +4$
  
- $[\text{CoBr}(\text{NH}_3)_5]\text{SO}_4$  pentaamminebromidocobalt(III) sulfate
  - ▣ Overall complex charge = -2
  - ▣ Ligand charge =  $\text{Br}^{-1}$  and  $\text{NH}_3$
  - ▣ Metal charge + ligand charge = overall complex charge
    - $\text{Co} + (-1) = +2$
    - $\text{Co} = +3$

# Polydentate Ligands


96

- Some ligands attach to the metal more than once



8 Ethylenediamine,  
 $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$

Bidentate ligand




9  $[\text{Co}(\text{en})_3]^{3+}$


# Polydentate Ligands

97

- Some ligands attach to the metal more than once




10 Ethylenediaminetetraacetic acid  
Hexadentate ligand



11 An edta complex

# Coordination Isomers

98



# Coordination Isomers

99

