



(51) International Patent Classification:

A61K 31/405 (2006.01) **C07C 255/41** (2006.01)
A61K 8/49 (2006.01) **C07D 471/04** (2006.01)
A61P 17/14 (2006.01) **C07D 275/03** (2006.01)
A61K 31/404 (2006.01) **C07D 413/12** (2006.01)
C07D 209/18 (2006.01) **C07F 9/38** (2006.01)
C07D 417/06 (2006.01)

(21) International Application Number:

PCT/US2018/040385

(22) International Filing Date:

29 June 2018 (29.06.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/527,775 30 June 2017 (30.06.2017) US
 62/654,095 06 April 2018 (06.04.2018) US

(71) Applicant: **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, Twelfth Floor, Oakland, CA 94607-5200 (US).

(72) Inventors; and

(71) Applicants: **LOWRY, William, E.** [—/US]; University of California Los Angeles, Department of Molecular Cell and Developmental Biology, 621 Charles Young Drive, Los Angeles, CA 90095 (US). **JUNG, Michael, E.** [US/US]; 2335 Manning Avenue, Los Angeles, CA 90064 (US). **CHRISTOFK, Heather** [—/US]; 2532 La Condesa Dr., Los Angeles, CA 90049 (US). **LIU, Xiaoguang** [—/US]; 1333 Harvard St., Apt. 34, Santa Monica, CA 90404 (US).

(74) Agent: **HALSTEAD, David, P.** et al.; Foley Hoag LLP, 155 Seaport Boulevard, Boston, MA 02210-2600 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))
 — before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: COMPOSITIONS AND METHODS FOR MODULATING HAIR GROWTH





FIG. 7A.

(57) Abstract: The present disclosure relates to compounds that are capable of inhibiting the mitochondrial pyruvate carrier and promoting hair growth. The disclosure further relates to methods of promoting hair growth or treating conditions or disorders affecting hair growth, such as baldness or alopecia.



COMPOSITIONS AND METHODS FOR MODULATING HAIR GROWTH

RELATED APPLICATIONS


This application claims the benefit of U.S. Provisional Patent Application No. 62/527,775, filed on June 30, 2017, and U.S. Provisional Patent Application No. 62/654,095, filed on April 6, 2018. The contents of each of these applications is hereby incorporated by reference in its entirety.

BACKGROUND

Hair follicle stem cells (HFSCs) undergo successive rounds of quiescence (telogen) punctuated by brief periods of proliferation correlating with the start of the hair cycle (telogen-anagen transition). Proliferation or activation of HFSCs is well known to be a prerequisite for advancement of the hair cycle. Despite advances in treatment options, baldness and alopecia continue to be conditions that cannot be successfully treated in all individuals. Some of the existing treatments are inconvenient for users, others require surgical intervention or other invasive procedures. Additional therapies are needed.


SUMMARY OF THE INVENTION

In certain aspects, the present disclosure provides compounds of formula I or II:



wherein:

each A is independently CH, CR⁴, or N;



Y is carboxyl, ester, amide, or

Z is CH, CR⁴, or N.

R² is CN or carboxyl;


R³ is H, aryl, aralkyl, or aralkylacyl, and is optionally substituted by one or more R⁵, wherein

each R⁵ is independently selected from alkyl, alkoxy, or halo;


each instance of R⁴ is independently alkyl, carboxyl, halo, hydroxy, ester, or CN;

R^6 is from H, alkyl, or cycloalkyl;
 R^7 is hydrogen, alkyl, halo, hydroxyl, alkoxy, or acyloxy;
 R^{10} is hydrogen or alkyl; and
 n is 0-4;
 or a pharmaceutically acceptable salt thereof.

In certain embodiments, the present disclosure provides compounds of formula III:



wherein:



Y is carboxyl, ester, amide, or ;

R^2 is CN or carboxyl;

R^3 is H, aryl, aralkyl, or aralkylacyl, and is optionally substituted by one or more R^5 , wherein
 each R^5 is independently selected from alkyl, alkoxy, or halo;

each instance of R^4 is independently alkyl, carboxyl, halo, hydroxy, ester, or CN;

R^6 is from H, alkyl, or cycloalkyl; and

R^7 is hydrogen, alkyl, halo, hydroxyl, alkoxy, or acyloxy;


R^{10} is hydrogen or alkyl;

R^{11} is hydrogen or alkyl; and

n is 0-4;

or a pharmaceutically acceptable salt thereof.

In certain embodiments, the present disclosure provides compounds of formula V, VI, or VII:



wherein:

each A is independently CH, CR⁴, or N;

X is NR⁶ or O;

R¹ is H or lower alkyl; or either R¹ and R⁶ or R¹ and R², together with the atoms that separate them, complete a heterocycle;

R² is CN or carboxyl;

R³ is H, aryl, aralkyl, or aralkylacyl, and is optionally substituted by one or more R⁵, wherein each R⁵ is independently selected from alkyl, alkoxy, and halo;


each instance of R⁴ is independently alkyl, carboxyl, halo, hydroxy, or CN;

R⁶ is from H, alkyl, or cycloalkyl; and

R⁷ is hydrogen, alkyl, halo, hydroxyl, alkoxy, or acyloxy;

or a pharmaceutically acceptable salt thereof.

In certain embodiments, the present disclosure provides compounds of formula Va, VIa, or VIIa:



wherein:

X is NR⁶ or O;

R¹ is H or lower alkyl;

R² is CN or carboxyl; or R¹ and R², together with the atoms that separate them, complete a heterocycle;

R³ is H, phenyl, or benzyl, and is optionally substituted by one or more R⁵, wherein each R⁵ is independently selected from alkyl, alkoxy, or halo;

each instance of R⁴ is independently selected from alkyl, carboxyl, halo, hydroxy, or CN; and

R⁶ is selected from H, alkyl, or cycloalkyl;

and pharmaceutically acceptable salts thereof.

In certain aspects, the present disclosure provides a pharmaceutical composition comprising a compound of the present disclosure and a pharmaceutically acceptable excipient.

In certain aspects, the present disclosure provides methods of enhancing lactate production in a cell, comprising contacting the cell with a compound or composition of the disclosure.

In certain aspects, the present disclosure provides methods of inhibiting mitochondrial pyruvate oxidation in a cell, comprising contacting the cell with a mitochondrial pyruvate oxidation (MPO) inhibitor, such as a compound of the present disclosure. In certain embodiments, the MPO inhibitor is a mitochondrial pyruvate carrier (MPC) inhibitor. In certain embodiments, inhibiting mitochondrial pyruvate oxidation in a cell has the effect of enhancing lactate production in a cell and/or enhancing the activity of LDH in a cell, and promoting hair growth, as described herein.

In certain aspects, the present disclosure provides methods of enhancing lactate production in a cell, comprising contacting the cell with an MPO inhibitor, such as a compound of the present disclosure. In certain embodiments, the MPO inhibitor is a mitochondrial pyruvate carrier (MPC) inhibitor.

In certain aspects, the present disclosure provides methods of enhancing the activity of LDH in a cell, comprising contacting the cell with an MPO inhibitor, such as a compound of the present disclosure. In certain embodiments, the MPO inhibitor is a mitochondrial pyruvate carrier (MPC) inhibitor.

In certain aspects, the present disclosure provides methods of enhancing the activity of lactic acid dehydrogenase (LDH) in a cell, comprising contacting the cell with an MPO inhibitor, such as a compound of the present disclosure. In certain embodiments, the MPO inhibitor is a mitochondrial pyruvate carrier (MPC) inhibitor. In certain aspects, the present disclosure provides methods of promoting hair growth or treating a hair growth condition or disorder such as baldness or alopecia, comprising administering to a patient a compound or composition as disclosed herein.

In certain aspects, the present disclosure provides methods of promoting hair growth or treating a hair growth condition or disorder such as baldness or alopecia, comprising administering to a patient an MPO inhibitor (e.g., topically, such as with a pharmaceutical composition formulated for topical application), such as a compound of the present disclosure. In certain embodiments, the present disclosure provides methods of promoting hair growth or treating a hair growth condition or disorder such as baldness or alopecia, comprising administering to a patient an MPC inhibitor (e.g., topically, such as with a pharmaceutical composition formulated for topical application), such a compound of the

present disclosure. In certain embodiments, inhibiting mitochondrial pyruvate oxidation or the mitochondrial pyruvate carrier in a cell has the effect of enhancing lactate production and/or enhancing the activity of LDH in a cell, and promoting hair growth, as described herein.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGs. 1A-1E show Lactate dehydrogenase activity is enriched in HFSCs. **FIG. 1A:** IHC staining for Ldha expression across the hair cycle shows Ldha protein confined to the HFSC niche, the bulge, indicated by the bracket. IHC staining for Sox9 on serial sections demarcates the HFSC population. Scale bar indicates 20 micrometers. **FIG. 1B:** Immunoblotting on FACS-isolated HFSC populations ($\alpha 6\text{low}/\text{Cd}34+$ and $\alpha 6\text{hi}/\text{Cd}34+$) versus total epidermis (Epi) shows differential expression of Ldha in the stem cell niche. Sox9 is a marker of HFSCs, and β -actin is a loading control. **FIG. 1C:** Colorimetric assay for Ldh enzyme activity in the epidermis shows highest activity in the bulge (brackets) and subcuticular muscle layer (bracket). This activity is enriched in the bulge across different stages of the hair cycle. Activity is indicated by purple color; pink is a nuclear counterstain. Note also that developing hair shafts in pigmented mice show strong deposits of melanin as observed here; hair shafts never displayed any purple stain indicative of Ldh activity. Scale bars indicate 50 micrometers. **FIG. 1D:** Ldh activity in sorted cell populations, measured using a plate reader-based assay, also shows the highest Ldh activity in two separate HFSC populations ($\alpha 6\text{hi}/\text{Cd}34$ and $\alpha 6\text{low}/\text{Cd}34$) compared to epidermal cells (Epi) and fibroblasts (FBs). Each bar represents the average signal for each cell type where $n=9$ mice pooled from 3 independent experiments. Shown as mean \pm SEM. Paired t-test was performed, $p < 0.05$ shown for each cell type versus epidermal cells. **FIG. 1E:** HFSCs and epidermal cells were isolated during telogen (day 50) by FACS, and metabolites were extracted and analyzed by LC-MS. Heatmaps show relative levels of glycolytic and TCA cycle metabolites from cells isolated from different mice in independent experiments with cells from three animals in each. Asterisks indicate significant difference in metabolite levels between epidermal cells and HFSCs. For e, paired t-test was performed; * denotes $p < 0.05$, ** denotes $p < 0.01$, *** denotes $p < 0.001$, ns denotes $p > 0.05$, and $n=9$ mice pooled from 3 independent experiments.

FIGs. 2A-2C show the validation of key reagents and assays. **FIG. 2A:** IHC with antibody recognizing specifically Ldha (same as used in **FIG. 1A**). bottom, IHC with

antibody recognizing multiple isoforms of Ldh protein. Scale bars indicate 20 micrometers.

FIG. 2B: the sorting strategy employed to isolate two populations of cells from the bulge.

This particular sort was used to isolate the protein samples shown by western blot in **FIG. 1B**.

FIG. 2C: Validation of colorimetric Ldh enzyme activity assay. The highest Ldh enzyme activity was observed in HFSC bulge and in the muscle. Activity indicated by purple stain; pink color is nuclear fast red counterstain. In absence of substrate lactate there was no detectable activity (purple stain). right, Additional validation of colorimetric Ldh enzyme activity assay. Enzyme activity inhibited by treating skin with HCl before addition of staining solution with substrate lactate. No Ldh activity (purple stain) detected. Skin in which enzyme activity is not inhibited by Hydrochloric Acid (HCl) shows highest Ldh enzyme activity in HFSC bulge and in the muscle. Scale bars indicate 50 micrometers.

FIGs. 3A-3E show that Ldh activity increases during HFSC activation. **FIG. 3A:** GSEA on RNA-seq transcriptome data from HFSCs versus total epidermis shows enrichment for Glycolysis related genes in HFSCs (NES = 1.72). **FIG. 3B:** GSEA on microarray transcriptome data from HFSCs versus total epidermis shows enrichment for Glycolysis related genes in HFSCs (NES = 1.45). Results were generated from three mice of each condition. **FIG. 3C:** RNA-seq data from HFSCs sorted during telogen or telogen-anagen transition show induction of *Ldha*²¹. Data represent the average of three separate animals at each timepoint. **FIG. 3D:** Ldh activity in sorted stem cell populations, measured using a plate reader-based assay, shows elevated Ldh activity as stem cells become activated in telogen to anagen transition (Tel-Ana). Each bar represents the average signal for each condition where n=9 mice pooled from 3 independent experiments. Shown as mean \pm SEM. Paired t-test was performed, $p < 0.05$. **FIG. 3E:** Heatmap showing relative levels of glycolytic and TCA cycle metabolites extracted from quiescent (Telogen, day 50), activated (Telogen-Anagen, day 70) and HFSCs that have returned to the quiescent state (Anagen, day 90). Data shown were generated from n=3 animals per timepoint in 3 independent experiments.

FIGs. 4A-4B show validation of hair cycle stage measurements. **FIG. 4A:** Analysis of RNA-seq data to validate that HFSCs in telogen-anagen transition were in fact in such a transition. The telogen-anagen transition is known to be driven by Shh (Gli factors are targets) and Wnt (Lef1, Axin, Ccnd1 are targets) signaling, and correlate with increased proliferation (Ki67 and Pcna). In addition, Sox4 was previously identified as a regulator of the telogen-anagen transition. n=3 mice per timepoint. Shown as mean \pm SEM. Paired t-test

was performed, $p < 0.05$. **FIG. 4B**: staining for Ki-67 marks dividing cells during various stages of the hair cycle. Brackets indicate the HFSC niche. Scale bars indicate 100 micrometers.

FIGs. 5A-5G show that deletion of *Mpc1* increases lactate production and activation of HFSCs. **FIG. 5A**: *Mpc1*^{fl/fl} animals show pigmentation and hair growth, consistent with entry into the anagen cycle at 8.5 weeks, whereas *Mpc1*^{+/+} animals do not show dorsal pigmentation and hair growth this early. Animals shown are representative of at least 12 animals of each genotype. **FIG. 5B**: FACS isolation of HFSC bulge populations in *Mpc1*^{+/+} versus *Mpc1*^{fl/fl} mice followed by western blotting shows successful deletion of *Mpc1* protein in the stem cell niche. β -actin is a loading control. **FIG. 5C**: Plate reader assay for Ldh activity on sorted HFSC populations shows elevated activity in *Mpc1*^{fl/fl} HFSCs compared to *Mpc1*^{+/+} HFSCs. Each bar represents the average signal for each genotype where $n=9$ mice pooled from 3 independent experiments. Shown as mean \pm SEM. Paired t-test was performed, $p < 0.05$. **FIG. 5D**: Histology on WT versus *Mpc1* deletion skin shows induction of anagen in absence of *Mpc1*. Scale bars indicate 100 micrometers. Quantification of phenotype at right shows percentage of dorsal follicles in telogen, telogen to anagen transition and anagen in *Mpc1*^{+/+} mice versus *Mpc1*^{fl/fl} mice ($n = 250$ follicles from 3 mice per genotype). Shown as mean \pm SEM. Paired t-test was performed, $p < 0.05$. **FIG. 5E**: Immunohistochemistry staining for Ki-67, a marker of proliferation that is only active in HFSCs at the beginning of a new hair cycle, is only present in *Mpc1*^{fl/fl} HFSCs at 8.5 weeks, consistent with their accelerated entry into a new hair cycle. Phospho-S6, another marker that is only active in HFSCs at the beginning of a new hair cycle, is only present in *Mpc1*^{fl/fl} HFSCs. Staining for Sox9 shows that HFSCs are present in *Mpc1* deleted niche. Images taken at 60X magnification. **FIG. 5F**: Deletion of *Mpc1* in mice bearing the *Lgr5*CreER allele shows strong induction of the hair cycle. Note that red boxes indicate areas of new hair growth. Results are representative of at least 9 animals per genotype. **FIG. 5G**: Quantification of pigmentation in the indicated genotypes across three independent litters ($n = 5$ mice per genotype).

FIGs. 6A-6D show the effects of long term deletion of *Mpc1* in HFSCs. **FIG. 6A**: Six months after initiation of deletion of *Mpc1* in HFSCs (*K15CrePR*;*Mpc1*^{fl/fl}), mice lacking *Mpc1* show no deleterious effects as measured by the hair cycle (left), pathology (middle, H and E), or staining for HFSCs (right, Sox9). Scale bars indicate 100 micrometers in middle panel, and 50 micrometers in right panel. Images are representative of at least 12 animals per

genotype. **FIG. 6B:** To demonstrate that the deletion of *Mpc1* promotes proliferation specifically in HFSCs, we used *K15CrePR;Ldha^{fl/fl}* mice bearing a lox-stop-lox-Tomato allele to look at K15+ HFSCs and proliferation with and without *Mpc1* deletion (left). In addition, we took advantage of the ires-GFP within the *Lgr5CreER* allele to stain for Ki-67 and GFP and look for co-localization with and without *Mpc1* deletion (right). White brackets denote bulge area. Scale bars represent 20 micrometers. **FIG. 6C:** Deletion of *Mpc1* in mice bearing the *Lgr6CreER* allele shows no premature induction of the hair cycle. **FIG. 6D:** Ldh activity assay on sorted HFSCs from either control or *Lgr6CreER* mediated *Mpc1* deletion mice showed increased activity in cells lacking *Mpc1*. n=6 mice per genotype pooled from 2 independent experiments. Shown as mean \pm SEM. Paired t-test was performed, $p < 0.05$.

FIGs. 7A-7D show that pharmacological inhibition of *Mpc1* promotes HFSC activation. **FIG. 7A:** Animals treated topically with UK-5099 (20 μ M) show pigmentation and hair growth, indicative of entry into anagen, after 8 days of treatment. Full anagen, indicated by full coat of hair, is achieved after 14 days of treatment. Mice treated topically with vehicle control do not show pigmentation nor hair growth even after 12 days of treatment. right, Skin pathology showing that UK-5099 animals enter an accelerated anagen at 8 weeks typified by down growth of the follicle and hypodermal thickening, while vehicle control treated animals showed neither and remained in telogen. Images shown are representative of at least 14 mice from 7 independent experiments. Scale bars indicate 100 micrometers. **FIG. 7B:** Graph showing time to observed phenotype in vehicle versus UK-5099 treated mice. n = 6 mice per condition. Shown as mean \pm SEM. **FIG. 7C:** Ldh enzyme activity assay in the epidermis shows strong activity in HFSCs in vehicle control and UK-5099 treated animals. Ldh enzyme activity also seen in interfollicular epidermis of UK-5099 treated animals. Ldh activity is indicated by purple stain; pink is nuclear fast red counterstain. Scale bars indicate 50 micrometers. **FIG. 7D:** Metabolomic analysis of Lactate on HFSCs isolated from UK-5099 treated skin for 48 hours; Each bar represents the average signal for each condition where n=9 mice pooled from 3 independent experiments. Shown as mean \pm SEM. Paired t-test was performed, $p < 0.05$.

FIG. 8 shows the effect on lactate production of certain *Mpc1* inhibitors described herein.

FIG. 9 shows the effect on lactate production of the certain *Mpc1* inhibitor described herein.

FIG. 10 shows the EC50 calculation for UK5099 and JXL020.

FIG. 11 shows that the Mpc1 inhibitors of the present invention induce hair growth.

FIG. 12 shows the effect on lactate production of certain Mpc1 inhibitors described herein.

FIG. 13 shows the effect on total cell count of certain Mpc1 inhibitors described herein, normalized to DMSO treatment.

FIG. 14 shows the effect on cell lactate production of certain Mpc1 inhibitors described herein, normalized to DMSO treatment.

FIG. 15 shows the effect on total cell count of certain Mpc1 inhibitors described herein, normalized to DMSO treatment.

FIG. 16 shows the effect on cell lactate production of certain Mpc1 inhibitors described herein, normalized to DMSO treatment.

FIG. 17 shows the effect on total cell count of certain Mpc1 inhibitors described herein, normalized to DMSO treatment.


FIG. 18 shows the effect on cell lactate production of certain Mpc1 inhibitors described herein, normalized to DMSO treatment.

FIG. 19 shows the effect on total cell count of certain Mpc1 inhibitors described herein, normalized to DMSO treatment.

FIG. 20 shows the role of MPC in the oxidation of pyruvate to acetyl coenzyme A.


DETAILED DESCRIPTION OF THE INVENTION

In certain aspects, the present disclosure provides compounds of formula I or II:



wherein:

each A is independently CH, CR⁴, or N;



Y is carboxyl, ester, amide, or

Z is CH, CR⁴, or N.


R² is CN or carboxyl;

R^3 is H, aryl, aralkyl, or aralkylacyl, and is optionally substituted by one or more R^5 , wherein each R^5 is independently selected from alkyl, alkoxy, or halo;
 each instance of R^4 is independently alkyl, carboxyl, halo, hydroxy, ester, or CN;
 R^6 is from H, alkyl, or cycloalkyl;
 R^7 is hydrogen, alkyl, halo, hydroxyl, alkoxy, or acyloxy;
 R^{10} is hydrogen or alkyl; and
 n is 0-4;
 or a pharmaceutically acceptable salt thereof.


In certain embodiments, the compound is a compound of formula I. In certain embodiments, the compound is a compound of formula II.

In certain embodiments of Formula I or II, Z is CH or N.

In certain embodiments, the present disclosure provides compounds of formula III or IV:



wherein,





Y is carboxyl, ester, amide, or ;

R^2 is CN or carboxyl;

R^3 is H, aryl, aralkyl, or aralkylacyl, and is optionally substituted by one or more R^5 , wherein each R^5 is independently selected from alkyl, alkoxy, or halo;
 each instance of R^4 is independently alkyl, carboxyl, halo, hydroxy, ester, or CN;
 R^6 is from H, alkyl, or cycloalkyl; and
 R^7 is hydrogen, alkyl, halo, hydroxyl, alkoxy, or acyloxy;
 R^{10} is hydrogen or alkyl;
 R^{11} is hydrogen or alkyl; and
 n is 0-4;
 or a pharmaceutically acceptable salt thereof.


In certain embodiments, the compound is a compound of formula III.



In certain embodiments of Formula I, II, and III, Y is wherein Y is . In certain embodiments, R¹⁰ is H. In certain embodiments, R¹⁰ is alkyl (e.g., ethyl). In certain embodiments Y is ester or amido.

In certain embodiments of Formula I, II, or III, R¹¹ is alkyl (e.g., methyl).

In certain embodiments, the present disclosure provides compounds of formula V, VI, or VII:



wherein:

each A is independently CH, CR⁴, or N;

X is NR⁶ or O;

R¹ is H or lower alkyl; or either R¹ and R⁶ or R¹ and R², together with the atoms that separate them, complete a heterocycle;

R² is CN or carboxyl;

R³ is H, aryl, aralkyl, or aralkylacyl, and is optionally substituted by one or more R⁵, wherein each R⁵ is independently selected from alkyl, alkoxy, and halo;

each instance of R⁴ is independently alkyl, carboxyl, halo, hydroxy, or CN;


R⁶ is from H, alkyl, or cycloalkyl; and

R⁷ is hydrogen, alkyl, halo, hydroxyl, alkoxy, or acyloxy; or a pharmaceutically acceptable salt thereof.

In certain embodiments, the compound is a compound of formula V. In certain embodiments, the compound is a compound of formula VI. In certain embodiments, the compound is a compound of formula VII.

In certain embodiments of formula I, II, III, V, VI, or VII, at least one A is N, preferably no more than two occurrences of A are N. In certain preferred embodiments, exactly one A is N, preferably the A bound to the same carbon as NR³.

In certain embodiments, the present disclosure provides compounds of formula Va, VIa, or VIIa:



wherein:

X is NR⁶ or O;

R¹ is H or lower alkyl;

R² is CN or carboxyl; or R¹ and R², together with the atoms that separate them, complete a heterocycle;

R³ is H, phenyl, or benzyl, and is optionally substituted by one or more R⁵, wherein each R⁵ is independently selected from alkyl, alkoxy, or halo;

each instance of R⁴ is independently selected from alkyl, carboxyl, halo, hydroxy, or CN; and

R⁶ is selected from H, alkyl, or cycloalkyl;

and pharmaceutically acceptable salts thereof.

In certain embodiments, the compound is a compound of formula Va. In certain embodiments, the compound is a compound of formula VIa. In certain embodiments, the compound is a compound of formula VIIa.

In certain embodiments of formula V, VI, VII, Va, VIa, or VIIa, X is NH. In certain embodiments, X is O.

In certain embodiments of formula V, VI, VII, Va, VIa, or VIIa, R¹ is H. In certain embodiments, R¹ is lower alkyl. In certain embodiments, R¹ and R⁶, together with the atoms that separate them, complete a heterocycle (e.g., morpholinyl).

In certain embodiments of formula V, VI, VII, Va, VIa, or VIIa, R⁶ is hydrogen.


In certain embodiments of formula V, VI, VII, Va, VIa, or VIIa, R² is CN. In certain embodiments, R² is carboxyl. In certain embodiments, R¹ and R², together with the atoms that separate them, complete a heterocycle selected from thiazolidine-2,4-dione-5-ylidene or 2-iminothiazolidin-4-one-5-ylidene.

In certain embodiments of formula I, II, III, V, VI, VII, Va, VIa, or VIIa, R³ is H. In certain embodiments, R³ is phenyl. In certain embodiments, R³ is phenyl and is substituted by one or more R⁵. In certain embodiments, R³ is substituted by one R⁵, and wherein R⁵ is an alkoxy. In certain embodiments, R³ is aralkyl (e.g., benzyl or phenethyl). In certain embodiments, R³ is aralkylacyl (e.g., phenylacetyl). In certain embodiments, R³ is benzyl. In


certain embodiments, R^3 is benzyl and is substituted by one or more R^5 . In certain embodiments, R^3 is aralkyl (e.g., benzyl or phenethyl) and is substituted by one or more R^5 (preferably on the phenyl ring). In certain embodiments, R^3 is aralkylacyl (e.g., phenylacetyl), and is substituted by one or more R^5 (preferably on the phenyl ring). In certain embodiments, R^3 is substituted by one or two R^5 , and wherein each R^5 is independently selected from fluoroalkyl or fluoro. In certain embodiments, R^3 is substituted by two R^5 , and wherein each R^5 is trifluoromethyl.

In certain embodiments of formula Va, VIa, or VIIa, n is 0.

In certain preferred embodiments, the present disclosure provides compounds of formula Vb




In certain embodiments, the present disclosure provides compounds of formula Vc




In certain embodiments of formula Va, VIa, VIIa, or Vb, n is 1.

In certain preferred embodiments, the present disclosure provides compounds of formula Vd



In certain embodiments, the present disclosure provides compounds of formula Ve



In certain embodiments of formula Vb, Vd or Ve, R^4 is selected from halo or haloalkyl. In certain preferred embodiments, R^4 is halo (e.g., chloro or bromo). In other preferred embodiments, R^4 is carboxyl or ester.


In certain embodiments of formula VI or VIa, n is 0. In certain embodiments, n is 2, and R^4 is selected from halo or haloalkyl.


In certain embodiments of formula I, II, III, V, VI, VII, Va, Vb, Vc, Vd, Ve, VIa, or VIIa, R^7 is hydrogen, hydroxyl, halo (e.g., chloro), or acyloxy (e.g., acetyloxy). In certain embodiments, R^7 is hydroxyl, halo (e.g., chloro), or acyloxy (e.g., acetyloxy).



















In certain embodiments of formula I, II, III, V, VI, VII, Va, Vb, Vc, Vd, Ve, VIa, or VIIa, the compound is not JXL001.






In certain aspects, the compound of the disclosure is a compound of Table 1.














Table 1: Exemplary Compounds of the Present Invention







**JXL024****JXL025****JXL026****JXL027****JXL028****JXL029****JXL030****JXL031****JXL032****JXL033****JXL034****JXL035****JXL036****JXL037****JXL038****JXL039****JXL040****JXL041**









In certain aspects, the present disclosure provides a pharmaceutical composition comprising a compound of the present disclosure and a pharmaceutically acceptable excipient. In certain embodiments, the pharmaceutical composition is formulated for topical administration.

In certain aspects, the present disclosure provides methods of enhancing lactate production in a cell, comprising contacting the cell with a compound or composition of the disclosure. In certain embodiments, the cell is a hair follicle stem cell.

In certain aspects, the present disclosure provides methods of promoting hair growth or treating hair growth conditions or disorders such as baldness or alopecia, comprising administering to a patient a compound or composition as disclosed herein.

In certain aspects, the present disclosure provides methods of enhancing lactate production in a cell, comprising contacting the cell with a mitochondrial pyruvate carrier (MPC) inhibitor. In certain embodiments, the MPC is MPC1.

In certain aspects, the present disclosure provides methods of promoting hair growth or treating a hair growth condition or disorder such as baldness or alopecia, comprising administering to a patient a mitochondrial pyruvate carrier (MPC) inhibitor (e.g., topically,

such as with a pharmaceutical composition formulated for topical application). In certain embodiments, the MPC is MPC1.

In certain aspects, the present disclosure provides methods of inhibiting mitochondrial pyruvate oxidation in a cell, comprising contacting the cell with a mitochondrial oxidation (MPO) inhibitor.

In certain aspects, the present disclosure provides methods of promoting hair growth or treating a hair growth condition or disorder such as baldness or alopecia, comprising administering to a patient a mitochondrial oxidation (MPO) inhibitor (e.g., topically, such as with a pharmaceutical composition formulated for topical application).

In certain embodiments, the MPC or MPO inhibitor is a compound of the disclosure.

Discussion

Numerous studies have uncovered unique gene expression signatures in hair follicle stem cells (HFSCs) versus other follicle cells or cells of the interfollicular epidermis. Many of these signatures are regulated by transcription factors that were later shown to play important roles in HFSC homeostasis.

Lactate dehydrogenase is most commonly encoded by the *Ldha* and *Ldhb* genes in mammals, the protein products of which form homo- or hetero-tetramers to catalyze the NADH-dependent reduction of pyruvate to lactate and NAD⁺-dependent oxidation of lactate to pyruvate. By immunostaining, it has been discovered that *Ldha* is enriched in quiescent HFSCs *in situ* (telogen) (FIG. 1A), and performing immunohistochemistry staining (IHC) with an antibody that recognizes both *Ldha* and *Ldhb* showed that only *Ldha* appears to be localized to the HFSC niche (FIG. 2A).

IHC analysis also showed *Ldha* expression was enriched in HFSCs (Sox9⁺) at three stages of the hair cycle (FIG. 1A). Consistently, immunoblotting of lysates from sorted cells showed strong expression of *Ldha* in the basal HFSCs (α 6HiCD34⁺), and suprabasal (α 6LoCD34⁺) HFSC populations relative to total epidermis (Fig 1B) (Sorting strategy is outlined in Fig 2B).

To determine whether *Ldha* expression patterns correlate with activity of the Ldh enzyme, a colorimetric-based enzymatic assay was used to assess Ldh activity capacity *in situ*. Typically performed on protein lysates or aliquots with a plate reader, the Ldh activity assay was adapted to work *in situ* on frozen tissue sections. Note that since both the *in situ* and *in vitro* Ldh activity assays employ use of excess substrate (lactate), the results from these assays reflect the capacity for Ldh activity, and not the steady-state activity.

Applying this assay to skin samples demonstrated that Ldh activity capacity was significantly higher in HFSCs, consistent with the expression pattern of *Ldha* (FIG. 1C). Furthermore, Ldh activity was enriched in HFSCs across the hair cycle (FIG. 1C). As a control, assays conducted without the enzymatic substrate (lactate) or on acid-treated tissue yielded zero activity (FIG. 2C). To further validate these results, we sorted epidermal populations, generated cell lysates on the sorted cells, and performed a similar colorimetric-based enzymatic assay on the sorted cell lysates, which also showed increased Ldh activity in HFSCs (FIG. 1D). To better characterize the metabolism of HFSCs, we performed metabolomics analysis on sorted populations from mouse skin by liquid chromatography-mass spectrometry (LC-MS) (Fig 1e). Several glycolytic metabolites, including glucose/fructose-6-phosphate, fructose-bisphosphate, dihydroxyacetone phosphate, 3-phosphoglycerate, and lactate, were routinely higher in HFSCs relative to total epidermis across three independent experiments (isolated from different mice on different days). Conversely, most TCA cycle metabolites were not consistently different between the epidermis and HFSCs (FIG. 1E). Collectively these results suggest that while all cells in the epidermis use the TCA cycle extensively to generate energy, HFSCs also have increased *Ldha* expression, Ldh activity, and glycolytic metabolism.


Measuring metabolism across the hair cycle therefore would capture any dynamic changes that occur in HFSCs that correlate with activation or quiescence. Analysis of RNA-seq data from HFSCs isolated during either telogen or the telogen-anagen transition demonstrated not only that *Ldha* is the predominant *Ldh* isoform expressed in HFSCs (FIG. 3), but is also induced during the telogen-anagen transition (FIGs. 3A and 3B (NIHGEORGSE67404 and GSE51635)). To confirm that the cells analyzed by RNA-seq were indeed either in telogen or the telogen to anagen transition, important markers of this transition were assessed including the Shh and Wnt pathways (*Gli1*, 2, 3; *Lef1*, *Axin1*, *Axin2*, *Ccnd1*) as well as proliferation markers (*Ki-67*, *Pcna* and *Sox4*) (Fig 4A).

The *in vitro* Ldh activity assay on lysates from sorted HFSCs uncovered a modest induction of Ldh activity correlating with the telogen to anagen transition (Fig 3D). Hair cycle staging was validated by Ki-67 immunostaining to determine HFSC activation (Fig 4B). Additionally, measurements of steady-state metabolites extracted from sorted HFSCs showed an increase in lactate in HFSCs as they transition from telogen to telogen-anagen transition, and then decrease again in anagen as HFSCs return to quiescence (Fig 3E).

To determine whether induction of lactate production could affect HFSC activation or the hair cycle, we crossed *K15CrePR* animals to those floxed for mitochondrial pyruvate carrier 1 (*Mpc1*) (*K15CrePR;Mpc1^{fl/fl}*). *Mpc1*, as a heterodimer with *Mpc2*, forms the mitochondrial pyruvate carrier MPC, a transporter on the inner mitochondrial membrane required for pyruvate entry into the mitochondria. Loss of function of *Mpc1* has been shown to drive lactate production through enhanced conversion of pyruvate to lactate by *Ldh*. Furthermore, inhibition of MPC results in a decrease in mitochondrial pyruvate oxidation (MPO) to acetyl coenzyme A (FIG. 20).

In animals with *Mpc1* deletion in HFSCs, we observed a strong acceleration of the ventral and dorsal hair cycles with all the typical features of a telogen-anagen transition (FIG. 5A) (n = 12 littermate pairs). Mifepristone treated *K15CrePR;Mpc1^{fl/fl}* animals were the only to show any signs of dorsal anagen by day 70. Western blotting on sorted HFSCs validated the loss of *Mpc1* protein (FIG. 5B). Importantly, purified HFSCs lacking *Mpc1* showed a strong induction of *Ldh* activity (FIG. 5C). Quantification of the dorsal hair cycle across three pairs of littermates showed a strong induction of anagen in backskin lacking *Mpc1* (FIG. 5D, right), and histology showed that the anagen induction was normal in appearance with a typical hypodermal expansion (FIG. 5D). Immunostaining demonstrated the induction in *Mpc1-null* HFSCs of various markers of hair cycle activation such as Ki-67 and pS6, while *Sox9* expression was unaffected (FIG. 5E). Long term deletion of *Mpc1* did not lead to aberrant follicles or exhaustion of HFSCs as judged by pathology and staining for *Sox9* (FIG. 6A). Furthermore, deletion of *Mpc1* with *Lgr5CreER* showed a very similar phenotype as deletion with *K15CrePR* (FIG. 5F and 5G), validating the fact that deletion of this protein in HFSCs leads to their activation (n = 12 pairs of littermates). Finally, immunofluorescence for the *Ires-GFP* of the *Lgr5CreER* transgene along with Ki-67 and lineage tracing with *K15CrePR;Mpc1^{fl/fl};Isl-Tomato* mice also demonstrated that the HFSCs were indeed proliferative following induction of *Mpc1* deletion by tamoxifen or mifepristone (FIG. 6B).

UK-5099 (also designated herein as JXL001) is a well-established pharmacological inhibitor of the mitochondrial pyruvate carrier and is known to promote lactate production as a result in various settings. UK-5099 has the following structure:



Topical treatment of animals in telogen (day 50) with UK-5099 led to a robust acceleration of the hair cycle, as well as minor hyperproliferation of the interfollicular epidermis (FIG. 7A). Quantification of the hair cycle across at least 6 pairs of animals (vehicle vs UK-5099) indicated a strong acceleration of the hair cycle, in as few as 6-9 days (FIG. 7B). Similar to genetic deletion of *Mpc1*, pharmacological blockade of the mitochondrial pyruvate carrier by UK-5099 for 48 hours during telogen promoted increased Ldh activity in HFSCs and the interfollicular epidermis, consistent with increased capacity for lactate production (FIG. 7C). Finally, metabolomic analysis demonstrated that topical application of UK-5099 increases total levels of lactate in sorted HFSCs (FIG. 7D).

Compounds were synthesized that could topically promote increased lactate levels and therefore drive the hair cycle.

The compounds were generally prepared by reaction of the corresponding aldehydes, e.g., for JXL001, 1-phenylindole-3-carboxaldehyde, with ethyl cyanoacetate in the presence of 40% aq. L-proline to give exclusively the *E*-isomer of the ethyl 2-cyano-3-(1-phenylindol-3-yl)propenoate, e.g., JXL004. Hydrolysis of the ester with mild lithium hydroxide afforded the *E*-isomer of the acid, e.g., JXL001. All of the other compounds were prepared by analogous methods using the specific aldehyde. The two heterocyclic compounds, JXL023 and JXL024, were prepared from the condensation of 1-phenyl-indole-3-carboxaldehyde with thiazolidine-2,4-dione and 2-iminothiazolidin-4-one. The structures of all of the compounds were determined using normal organic chemistry methods, especially high field proton, carbon, and fluorine NMR spectra. In particular, $^3\text{J}_{\text{C-H}}$ coupling measurements demonstrated that the compounds all had the *E*-stereochemistry about the key carbon-carbon double bond.

To determine whether these compounds could promote cellular lactate production, we treated cultured epithelial cells with the compounds and measured lactate levels in the culture media using a Nova Biomedical BioProfile Basic Analyzer. Briefly, cultured epithelial cells were treated with DMSO, UK-5099 (also called JXL001), or certain of the exemplary compounds disclosed herein for 24-30 hours, and media lactate levels were measured and

normalized to cell number and duration of the experiment to acquire a cellular lactate production rate (nmol lactate, million cells, hour). The results are shown in FIGs. 8 and 9.

Lactate production rates of treated cells are shown in FIG. 8. As expected since they are UK-5099 analogues, most of the novel compounds assayed increased lactate production. A separate assay was performed to calculate the EC₅₀ of some of the compounds as shown in FIG. 10.

To determine the efficacy of the compounds on the hair cycle, mice were shaved at postnatal day 50, and topically treated with a compound disclosed herein suspended in lotion in every other day for 3 weeks. As seen in FIG. 11, all the analogues that showed the ability to promote lactate production in the in vitro assay were also able to stimulate hair growth over the course of 2 weeks.

Pharmaceutical Compositions

The compositions and methods of the present invention may be utilized to treat an individual in need thereof. In certain embodiments, the individual is a mammal such as a human, or a non-human mammal. When administered to an animal, such as a human, the composition or the compound is preferably administered as a pharmaceutical composition comprising, for example, a compound of the invention and a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers are well known in the art and include, for example, aqueous solutions such as water or physiologically buffered saline or other solvents or vehicles such as glycols, glycerol, oils such as olive oil, or injectable organic esters. In preferred embodiments, when such pharmaceutical compositions are for human administration, particularly for invasive routes of administration (i.e., routes, such as injection or implantation, that circumvent transport or diffusion through an epithelial barrier), the aqueous solution is pyrogen-free, or substantially pyrogen-free. The excipients can be chosen, for example, to effect delayed release of an agent or to selectively target one or more cells, tissues or organs. The pharmaceutical composition can be in dosage unit form such as tablet, capsule (including sprinkle capsule and gelatin capsule), granule, lyophile for reconstitution, powder, solution, syrup, suppository, injection or the like. The composition can also be present in a transdermal delivery system, e.g., a skin patch. The composition can also be present in a solution suitable for topical administration, such as a lotion, cream, or ointment.

A pharmaceutically acceptable carrier can contain physiologically acceptable agents that act, for example, to stabilize, increase solubility or to increase the absorption of a compound such as a compound of the invention. Such physiologically acceptable agents include, for example, carbohydrates, such as glucose, sucrose or dextrans, antioxidants, such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins or other stabilizers or excipients. The choice of a pharmaceutically acceptable carrier, including a physiologically acceptable agent, depends, for example, on the route of administration of the composition. The preparation or pharmaceutical composition can be a selfemulsifying drug delivery system or a selfmicroemulsifying drug delivery system. The pharmaceutical composition (preparation) also can be a liposome or other polymer matrix, which can have incorporated therein, for example, a compound of the invention. Liposomes, for example, which comprise phospholipids or other lipids, are nontoxic, physiologically acceptable and metabolizable carriers that are relatively simple to make and administer.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

The phrase "pharmaceutically acceptable carrier" as used herein means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer

solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

A pharmaceutical composition (preparation) can be administered to a subject by any of a number of routes of administration including, for example, orally (for example, drenches as in aqueous or non-aqueous solutions or suspensions, tablets, capsules (including sprinkle capsules and gelatin capsules), boluses, powders, granules, pastes for application to the tongue); absorption through the oral mucosa (e.g., sublingually); subcutaneously; transdermally (for example as a patch applied to the skin); and topically (for example, as a cream, ointment or spray applied to the skin). The compound may also be formulated for inhalation. In certain embodiments, a compound may be simply dissolved or suspended in sterile water. Details of appropriate routes of administration and compositions suitable for same can be found in, for example, U.S. Pat. Nos. 6,110,973, 5,763,493, 5,731,000, 5,541,231, 5,427,798, 5,358,970 and 4,172,896, as well as in patents cited therein.

The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will vary depending upon the host being treated, the particular mode of administration. The amount of active ingredient that can be combined with a carrier material to produce a single dosage form will generally be that amount of the compound which produces a therapeutic effect. Generally, out of one hundred percent, this amount will range from about 1 percent to about ninety-nine percent of active ingredient, preferably from about 5 percent to about 70 percent, most preferably from about 10 percent to about 30 percent.

Methods of preparing these formulations or compositions include the step of bringing into association an active compound, such as a compound of the invention, with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a compound of the present invention with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

Formulations of the invention suitable for oral administration may be in the form of capsules (including sprinkle capsules and gelatin capsules), cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), lyophile, powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert

base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of a compound of the present invention as an active ingredient. Compositions or compounds may also be administered as a bolus, electuary or paste.

To prepare solid dosage forms for oral administration (capsules (including sprinkle capsules and gelatin capsules), tablets, pills, dragees, powders, granules and the like), the active ingredient is mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, cetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; (10) complexing agents, such as, modified and unmodified cyclodextrins; and (11) coloring agents. In the case of capsules (including sprinkle capsules and gelatin capsules), tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

The tablets, and other solid dosage forms of the pharmaceutical compositions, such as dragees, capsules (including sprinkle capsules and gelatin capsules), pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient therein using, for example,

hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions that can be dissolved in sterile water, or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. The active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

Liquid dosage forms useful for oral administration include pharmaceutically acceptable emulsions, lyophiles for reconstitution, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, cyclodextrins and derivatives thereof, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

Dosage forms for the topical or transdermal administration include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants that may be required.

The ointments, pastes, creams and gels may contain, in addition to an active compound, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch,

tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

Powders and sprays can contain, in addition to an active compound, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

Transdermal patches have the added advantage of providing controlled delivery of a compound of the present invention to the body. Such dosage forms can be made by dissolving or dispersing the active compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate of such flux can be controlled by either providing a rate controlling membrane or dispersing the compound in a polymer matrix or gel.

The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion. Pharmaceutical compositions suitable for parenteral administration comprise one or more active compounds in combination with one or more pharmaceutically acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

Examples of suitable aqueous and nonaqueous carriers that may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents that delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution, which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsulated matrices of the subject compounds in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions that are compatible with body tissue.

For use in the methods of this invention, active compounds can be given per se or as a pharmaceutical composition containing, for example, 0.1 to 99.5% (more preferably, 0.5 to 90%) of active ingredient in combination with a pharmaceutically acceptable carrier.

Methods of introduction may also be provided by rechargeable or biodegradable devices. Various slow release polymeric devices have been developed and tested *in vivo* in recent years for the controlled delivery of drugs, including proteinaceous biopharmaceuticals. A variety of biocompatible polymers (including hydrogels), including both biodegradable and non-degradable polymers, can be used to form an implant for the sustained release of a compound at a particular target site.

Actual dosage levels of the active ingredients in the pharmaceutical compositions may be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

The selected dosage level will depend upon a variety of factors including the activity of the particular compound or combination of compounds employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound(s) being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compound(s) employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the therapeutically effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the pharmaceutical composition or compound at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. By “therapeutically effective amount” is meant the amount of a compound that is sufficient to elicit the desired therapeutic effect. It is generally understood that the effective amount of the compound will vary according to the weight, sex, age, and medical history of the subject. Other factors which influence the effective amount may include, but are not limited to, the severity of the patient's condition, the disorder being treated, the stability of the compound, and, if desired, another type of therapeutic agent being administered with the compound of the invention. A larger total dose can be delivered by multiple administrations of the agent. Methods to determine efficacy and dosage are known to those skilled in the art (Isselbacher et al. (1996) Harrison's Principles of Internal Medicine 13 ed., 1814-1882, herein incorporated by reference).

In general, a suitable daily dose of an active compound used in the compositions and methods of the invention will be that amount of the compound that is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above.

If desired, the effective daily dose of the active compound may be administered as one, two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms. In certain embodiments of the present invention, the active compound may be administered two or three times daily. In preferred embodiments, the active compound will be administered once daily.

The patient receiving this treatment is any animal in need, including primates, in particular humans; and other mammals such as equines, cattle, swine, sheep, cats, and dogs; poultry; and pets in general.

In certain embodiments, compounds of the invention may be used alone or conjointly administered with another type of therapeutic agent.

The present disclosure includes the use of pharmaceutically acceptable salts of compounds of the invention in the compositions and methods of the present invention. In certain embodiments, contemplated salts of the invention include, but are not limited to, alkyl, dialkyl, trialkyl or tetra-alkyl ammonium salts. In certain embodiments, contemplated salts of the invention include, but are not limited to, L-arginine, benenthamine, benzathine, betaine, calcium hydroxide, choline, deanol, diethanolamine, diethylamine, 2-(diethylamino)ethanol, ethanolamine, ethylenediamine, N-methylglucamine, hydrabamine, 1H-imidazole, lithium, L-lysine, magnesium, 4-(2-hydroxyethyl)morpholine, piperazine, potassium, 1-(2-hydroxyethyl)pyrrolidine, sodium, triethanolamine, tromethamine, and zinc salts. In certain embodiments, contemplated salts of the invention include, but are not limited to, Na, Ca, K, Mg, Zn or other metal salts. In certain embodiments, contemplated salts of the invention include, but are not limited to, 1-hydroxy-2-naphthoic acid, 2,2-dichloroacetic acid, 2-hydroxyethanesulfonic acid, 2-oxoglutaric acid, 4-acetamidobenzoic acid, 4-aminosalicylic acid, acetic acid, adipic acid, l-ascorbic acid, l-aspartic acid, benzenesulfonic acid, benzoic acid, (+)-camphoric acid, (+)-camphor-10-sulfonic acid, capric acid (decanoic acid), caproic acid (hexanoic acid), caprylic acid (octanoic acid), carbonic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, d-glucoheptonic acid, d-gluconic acid, d-glucuronic acid, glutamic acid, glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, hydrobromic acid, hydrochloric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid, l-malic acid, malonic acid, mandelic acid, methanesulfonic acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, nicotinic acid, nitric acid, oleic acid, oxalic acid, palmitic acid, pamoic acid, phosphoric acid, proprionic acid, l-pyroglutamic acid, salicylic acid, sebacic acid, stearic acid, succinic acid, sulfuric acid, l-tartaric acid, thiocyanic acid, p-toluenesulfonic acid, trifluoroacetic acid, and undecylenic acid acid salts.

The pharmaceutically acceptable acid addition salts can also exist as various solvates, such as with water, methanol, ethanol, dimethylformamide, and the like. Mixtures of such

solvates can also be prepared. The source of such solvate can be from the solvent of crystallization, inherent in the solvent of preparation or crystallization, or adventitious to such solvent.

Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions.

Examples of pharmaceutically acceptable antioxidants include: (1) water-soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal-chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

Definitions

Unless otherwise defined herein, scientific and technical terms used in this application shall have the meanings that are commonly understood by those of ordinary skill in the art. Generally, nomenclature used in connection with, and techniques of, chemistry, cell and tissue culture, molecular biology, cell and cancer biology, neurobiology, neurochemistry, virology, immunology, microbiology, pharmacology, genetics and protein and nucleic acid chemistry, described herein, are those well known and commonly used in the art.

The methods and techniques of the present disclosure are generally performed, unless otherwise indicated, according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout this specification. See, e.g. "Principles of Neural Science", McGraw-Hill Medical, New York, N.Y. (2000); Motulsky, "Intuitive Biostatistics", Oxford University Press, Inc. (1995); Lodish et al., "Molecular Cell Biology, 4th ed.", W. H. Freeman & Co., New York (2000); Griffiths et al., "Introduction to Genetic Analysis, 7th ed.", W. H. Freeman & Co., N.Y. (1999); and Gilbert et al., "Developmental Biology, 6th ed.", Sinauer Associates, Inc., Sunderland, MA (2000).

Chemistry terms used herein, unless otherwise defined herein, are used according to conventional usage in the art, as exemplified by “The McGraw-Hill Dictionary of Chemical Terms”, Parker S., Ed., McGraw-Hill, San Francisco, C.A. (1985).

All of the above, and any other publications, patents and published patent applications referred to in this application are specifically incorporated by reference herein. In case of conflict, the present specification, including its specific definitions, will control.

The term “agent” is used herein to denote a chemical compound (such as an organic or inorganic compound, a mixture of chemical compounds), a biological macromolecule (such as a nucleic acid, an antibody, including parts thereof as well as humanized, chimeric and human antibodies and monoclonal antibodies, a protein or portion thereof, e.g., a peptide, a lipid, a carbohydrate), or an extract made from biological materials such as bacteria, plants, fungi, or animal (particularly mammalian) cells or tissues. Agents include, for example, agents whose structure is known, and those whose structure is not known. The ability of such agents to inhibit AR or promote AR degradation may render them suitable as “therapeutic agents” in the methods and compositions of this disclosure.

A “patient,” “subject,” or “individual” are used interchangeably and refer to either a human or a non-human animal. These terms include mammals, such as humans, primates, livestock animals (including bovines, porcines, etc.), companion animals (e.g., canines, felines, etc.) and rodents (e.g., mice and rats).

“Treating” a condition or patient refers to taking steps to obtain beneficial or desired results, including clinical results. As used herein, and as well understood in the art, “treatment” is an approach for obtaining beneficial or desired results, including clinical results. Beneficial or desired clinical results can include, but are not limited to, alleviation or amelioration of one or more symptoms or conditions, diminishment of extent of disease, stabilized (i.e. not worsening) state of disease, preventing spread of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. “Treatment” can also mean prolonging survival as compared to expected survival if not receiving treatment.

The term “preventing” is art-recognized, and when used in relation to a condition, such as a local recurrence (e.g., pain), a disease such as cancer, a syndrome complex such as heart failure or any other medical condition, is well understood in the art, and includes administration of a composition which reduces the frequency of, or delays the onset of, symptoms of a medical condition in a subject relative to a subject which does not receive the

composition. Thus, prevention of cancer includes, for example, reducing the number of detectable cancerous growths in a population of patients receiving a prophylactic treatment relative to an untreated control population, and/or delaying the appearance of detectable cancerous growths in a treated population versus an untreated control population, e.g., by a statistically and/or clinically significant amount.

“Administering” or “administration of” a substance, a compound or an agent to a subject can be carried out using one of a variety of methods known to those skilled in the art. For example, a compound or an agent can be administered, intravenously, arterially, intradermally, intramuscularly, intraperitoneally, subcutaneously, ocularly, sublingually, orally (by ingestion), intranasally (by inhalation), intraspinally, intracerebrally, and transdermally (by absorption, e.g., through a skin duct). A compound or agent can also appropriately be introduced by rechargeable or biodegradable polymeric devices or other devices, e.g., patches and pumps, or formulations, which provide for the extended, slow or controlled release of the compound or agent. Administering can also be performed, for example, once, a plurality of times, and/or over one or more extended periods.

Appropriate methods of administering a substance, a compound or an agent to a subject will also depend, for example, on the age and/or the physical condition of the subject and the chemical and biological properties of the compound or agent (e.g., solubility, digestibility, bioavailability, stability and toxicity). In some embodiments, a compound or an agent is administered orally, e.g., to a subject by ingestion. In some embodiments, the orally administered compound or agent is in an extended release or slow release formulation, or administered using a device for such slow or extended release.

As used herein, the phrase “conjoint administration” refers to any form of administration of two or more different therapeutic agents such that the second agent is administered while the previously administered therapeutic agent is still effective in the body (e.g., the two agents are simultaneously effective in the patient, which may include synergistic effects of the two agents). For example, the different therapeutic compounds can be administered either in the same formulation or in separate formulations, either concomitantly or sequentially. Thus, an individual who receives such treatment can benefit from a combined effect of different therapeutic agents.

A “therapeutically effective amount” or a “therapeutically effective dose” of a drug or agent is an amount of a drug or an agent that, when administered to a subject, will have the intended therapeutic effect. The full therapeutic effect does not necessarily occur by

administration of one dose, and may occur only after administration of a series of doses. Thus, a therapeutically effective amount may be administered in one or more administrations. The precise effective amount needed for a subject will depend upon, for example, the subject's size, health and age, and the nature and extent of the condition being treated, such as cancer or MDS. The skilled worker can readily determine the effective amount for a given situation by routine experimentation.

The term "acyl" is art-recognized and refers to a group represented by the general formula hydrocarbylC(O)-, preferably alkylC(O)-.

The term "acylamino" is art-recognized and refers to an amino group substituted with an acyl group and may be represented, for example, by the formula hydrocarbylC(O)NH-.

The term "acyloxy" is art-recognized and refers to a group represented by the general formula hydrocarbylC(O)O-, preferably alkylC(O)O-.

The term "alkoxy" refers to an alkyl group having an oxygen attached thereto. Representative alkoxy groups include methoxy, ethoxy, propoxy, tert-butoxy and the like.

The term "alkoxyalkyl" refers to an alkyl group substituted with an alkoxy group and may be represented by the general formula alkyl-O-alkyl.

The term "alkyl" refers to saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl-substituted cycloalkyl groups, and cycloalkyl-substituted alkyl groups. In preferred embodiments, a straight chain or branched chain alkyl has 30 or fewer carbon atoms in its backbone (e.g., C₁₋₃₀ for straight chains, C₃₋₃₀ for branched chains), and more preferably 20 or fewer.


Moreover, the term "alkyl" as used throughout the specification, examples, and claims is intended to include both unsubstituted and substituted alkyl groups, the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone, including haloalkyl groups such as trifluoromethyl and 2,2,2-trifluoroethyl, etc.

The term "C_{x-y}" or "C_x-C_y", when used in conjunction with a chemical moiety, such as, acyl, acyloxy, alkyl, alkenyl, alkynyl, or alkoxy is meant to include groups that contain from x to y carbons in the chain. C₀alkyl indicates a hydrogen where the group is in a terminal position, a bond if internal. A C₁₋₆alkyl group, for example, contains from one to six carbon atoms in the chain.

The term "alkylamino", as used herein, refers to an amino group substituted with at least one alkyl group.


The term “alkylthio”, as used herein, refers to a thiol group substituted with an alkyl group and may be represented by the general formula alkylS-

The term “amide”, as used herein, refers to a group



wherein R^9 and R^{10} each independently represent a hydrogen or hydrocarbonyl group, or R^9 and R^{10} taken together with the N atom to which they are attached complete a heterocycle having from 4 to 8 atoms in the ring structure.

The terms “amine” and “amino” are art-recognized and refer to both unsubstituted and substituted amines and salts thereof, e.g., a moiety that can be represented by




wherein R^9 , R^{10} , and $\text{R}^{10'}$ each independently represent a hydrogen or a hydrocarbonyl group, or R^9 and R^{10} taken together with the N atom to which they are attached complete a heterocycle having from 4 to 8 atoms in the ring structure.

The term “aminoalkyl”, as used herein, refers to an alkyl group substituted with an amino group.

The term “aralkyl”, as used herein, refers to an alkyl group substituted with an aryl group.

The term “aryl” as used herein include substituted or unsubstituted single-ring aromatic groups in which each atom of the ring is carbon. Preferably the ring is a 5- to 7-membered ring, more preferably a 6-membered ring. The term “aryl” also includes polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings wherein at least one of the rings is aromatic, e.g., the other cyclic rings can be cycloalkyls, cycloalkenyls, cycloalkynyls, aryls, heteroaryls, and/or heterocyclyls. Aryl groups include benzene, naphthalene, phenanthrene, phenol, aniline, and the like.

The term “carbamate” is art-recognized and refers to a group



wherein R^9 and R^{10} independently represent hydrogen or a hydrocarbonyl group.

The term “carbocyclalkyl”, as used herein, refers to an alkyl group substituted with a carbocycle group.

The terms “carbocycle”, “carbocyclyl”, and “carbocyclic”, as used herein, refers to a non-aromatic saturated or unsaturated ring in which each atom of the ring is carbon. Preferably a carbocycle ring contains from 3 to 10 atoms, more preferably from 5 to 7 atoms.

The term “carbocyclalkyl”, as used herein, refers to an alkyl group substituted with a carbocycle group.

The term “carbonate” is art-recognized and refers to a group $-\text{OCO}_2-$.

The term “carboxy”, as used herein, refers to a group represented by the formula $-\text{CO}_2\text{H}$.

The term “ester”, as used herein, refers to a group $-\text{C}(\text{O})\text{OR}^9$ wherein R^9 represents a hydrocarbyl group.

The term “ether”, as used herein, refers to a hydrocarbyl group linked through an oxygen to another hydrocarbyl group. Accordingly, an ether substituent of a hydrocarbyl group may be hydrocarbyl-O-. Ethers may be either symmetrical or unsymmetrical. Examples of ethers include, but are not limited to, heterocycle-O-heterocycle and aryl-O-heterocycle. Ethers include “alkoxyalkyl” groups, which may be represented by the general formula alkyl-O-alkyl.

The terms “halo” and “halogen” as used herein means halogen and includes chloro, fluoro, bromo, and iodo.

The terms “hetaralkyl” and “heteroaralkyl”, as used herein, refers to an alkyl group substituted with a hetaryl group.

The terms “heteroaryl” and “hetaryl” include substituted or unsubstituted aromatic single ring structures, preferably 5- to 7-membered rings, more preferably 5- to 6-membered rings, whose ring structures include at least one heteroatom, preferably one to four heteroatoms, more preferably one or two heteroatoms. The terms “heteroaryl” and “hetaryl” also include polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings wherein at least one of the rings is heteroaromatic, e.g., the other cyclic rings can be cycloalkyls, cycloalkenyls, cycloalkynyls, aryls, heteroaryls, and/or heterocyclyls. Heteroaryl groups include, for example, pyrrole, furan, thiophene, imidazole, oxazole, thiazole, pyrazole, pyridine, pyrazine, pyridazine, and pyrimidine, and the like.

The term “heteroatom” as used herein means an atom of any element other than carbon or hydrogen. Preferred heteroatoms are nitrogen, oxygen, and sulfur.

The term “heterocyclalkyl”, as used herein, refers to an alkyl group substituted with a heterocycle group.

The terms “heterocycl”, “heterocycle”, and “heterocyclic” refer to substituted or unsubstituted non-aromatic ring structures, preferably 3- to 10-membered rings, more preferably 3- to 7-membered rings, whose ring structures include at least one heteroatom, preferably one to four heteroatoms, more preferably one or two heteroatoms. The terms “heterocycl” and “heterocyclic” also include polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings wherein at least one of the rings is heterocyclic, e.g., the other cyclic rings can be cycloalkyls, cycloalkenyls, cycloalkynyls, aryls, heteroaryl, and/or heterocycls. Heterocycl groups include, for example, piperidine, piperazine, pyrrolidine, morpholine, lactones, lactams, and the like.

The term “hydrocarb”, as used herein, refers to a group that is bonded through a carbon atom that does not have a =O or =S substituent, and typically has at least one carbon-hydrogen bond and a primarily carbon backbone, but may optionally include heteroatoms. Thus, groups like methyl, ethoxyethyl, 2-pyridyl, and even trifluoromethyl are considered to be hydrocarb for the purposes of this application, but substituents such as acetyl (which has a =O substituent on the linking carbon) and ethoxy (which is linked through oxygen, not carbon) are not. Hydrocarb groups include, but are not limited to aryl, heteroaryl, carbocycle, heterocycle, alkyl, alkenyl, alkynyl, and combinations thereof.


The term “hydroxyalkyl”, as used herein, refers to an alkyl group substituted with a hydroxy group.

The term “lower” when used in conjunction with a chemical moiety, such as, acyl, acyloxy, alkyl, alkenyl, alkynyl, or alkoxy is meant to include groups where there are ten or fewer atoms in the substituent, preferably six or fewer. A “lower alkyl”, for example, refers to an alkyl group that contains ten or fewer carbon atoms, preferably six or fewer. In certain embodiments, acyl, acyloxy, alkyl, alkenyl, alkynyl, or alkoxy substituents defined herein are respectively lower acyl, lower acyloxy, lower alkyl, lower alkenyl, lower alkynyl, or lower alkoxy, whether they appear alone or in combination with other substituents, such as in the recitations hydroxyalkyl and aralkyl (in which case, for example, the atoms within the aryl group are not counted when counting the carbon atoms in the alkyl substituent).

The terms “polycyclyl”, “polycycle”, and “polycyclic” refer to two or more rings (e.g., cycloalkyls, cycloalkenyls, cycloalkynyls, aryls, heteroaryls, and/or heterocyclyls) in which two or more atoms are common to two adjoining rings, e.g., the rings are “fused rings”. Each of the rings of the polycycle can be substituted or unsubstituted. In certain embodiments, each ring of the polycycle contains from 3 to 10 atoms in the ring, preferably from 5 to 7.

The term “sulfate” is art-recognized and refers to the group $-\text{OSO}_3\text{H}$, or a pharmaceutically acceptable salt thereof.

The term “sulfonamide” is art-recognized and refers to the group represented by the general formulae



wherein R^9 and R^{10} independently represents hydrogen or hydrocarbyl.

The term “sulfoxide” is art-recognized and refers to the group $-\text{S}(\text{O})-$.

The term “sulfonate” is art-recognized and refers to the group SO_3H , or a pharmaceutically acceptable salt thereof.

The term “sulfone” is art-recognized and refers to the group $-\text{S}(\text{O})_2-$.

The term “substituted” refers to moieties having substituents replacing a hydrogen on one or more carbons of the backbone. It will be understood that “substitution” or “substituted with” includes the implicit proviso that such substitution is in accordance with permitted valence of the substituted atom and the substituent, and that the substitution results in a stable compound, e.g., which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, etc. As used herein, the term “substituted” is contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and non-aromatic substituents of organic compounds. The permissible substituents can be one or more and the same or different for appropriate organic compounds. For purposes of this invention, the heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valences of the heteroatoms. Substituents can include any substituents described herein, for example, a halogen, a hydroxyl, a carbonyl (such as a carboxyl, an alkoxycarbonyl, a formyl, or an acyl), a thiocarbonyl (such as a thioester, a


thioacetate, or a thioformate), an alkoxyl, a phosphoryl, a phosphate, a phosphonate, a phosphinate, an amino, an amido, an amidine, an imine, a cyano, a nitro, an azido, a sulfhydryl, an alkylthio, a sulfate, a sulfonate, a sulfamoyl, a sulfonamido, a sulfonyl, a heterocyclyl, an aralkyl, or an aromatic or heteroaromatic moiety. It will be understood by those skilled in the art that the moieties substituted on the hydrocarbon chain can themselves be substituted, if appropriate.

The term “thioalkyl”, as used herein, refers to an alkyl group substituted with a thiol group.

The term “thioester”, as used herein, refers to a group $-C(O)SR^9$ or $-SC(O)R^9$ wherein R^9 represents a hydrocarbyl.

The term “thioether”, as used herein, is equivalent to an ether, wherein the oxygen is replaced with a sulfur.

The term “urea” is art-recognized and may be represented by the general formula



wherein R^9 and R^{10} independently represent hydrogen or a hydrocarbyl.

The term “modulate” as used herein includes the inhibition or suppression of a function or activity (such as cell proliferation) as well as the enhancement of a function or activity.

The phrase “pharmaceutically acceptable” is art-recognized. In certain embodiments, the term includes compositions, excipients, adjuvants, polymers and other materials and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

“Pharmaceutically acceptable salt” or “salt” is used herein to refer to an acid addition salt or a basic addition salt which is suitable for or compatible with the treatment of patients.

The term “pharmaceutically acceptable acid addition salt” as used herein means any non-toxic organic or inorganic salt of any base compounds represented by formula I or II. Illustrative inorganic acids which form suitable salts include hydrochloric, hydrobromic, sulfuric and phosphoric acids, as well as metal salts such as sodium monohydrogen orthophosphate and potassium hydrogen sulfate. Illustrative organic acids that form suitable

salts include mono-, di-, and tricarboxylic acids such as glycolic, lactic, pyruvic, malonic, succinic, glutaric, fumaric, malic, tartaric, citric, ascorbic, maleic, benzoic, phenylacetic, cinnamic and salicylic acids, as well as sulfonic acids such as p-toluene sulfonic and methanesulfonic acids. Either the mono or di-acid salts can be formed, and such salts may exist in either a hydrated, solvated or substantially anhydrous form. In general, the acid addition salts of compounds of formula I or II are more soluble in water and various hydrophilic organic solvents, and generally demonstrate higher melting points in comparison to their free base forms. The selection of the appropriate salt will be known to one skilled in the art. Other non-pharmaceutically acceptable salts, e.g., oxalates, may be used, for example, in the isolation of compounds of formula I or II for laboratory use, or for subsequent conversion to a pharmaceutically acceptable acid addition salt.

The term “pharmaceutically acceptable basic addition salt” as used herein means any non-toxic organic or inorganic base addition salt of any acid compounds represented by formula I or II or any of their intermediates. Illustrative inorganic bases which form suitable salts include lithium, sodium, potassium, calcium, magnesium, or barium hydroxide. Illustrative organic bases which form suitable salts include aliphatic, alicyclic, or aromatic organic amines such as methylamine, trimethylamine and picoline or ammonia. The selection of the appropriate salt will be known to a person skilled in the art.

Many of the compounds useful in the methods and compositions of this disclosure have at least one stereogenic center in their structure. This stereogenic center may be present in a R or a S configuration, said R and S notation is used in correspondence with the rules described in Pure Appl. Chem. (1976), 45, 11-30. The disclosure contemplates all stereoisomeric forms such as enantiomeric and diastereoisomeric forms of the compounds, salts, prodrugs or mixtures thereof (including all possible mixtures of stereoisomers). See, e.g., WO 01/062726.

Furthermore, certain compounds which contain alkenyl groups may exist as Z (zusammen) or E (entgegen) isomers. In each instance, the disclosure includes both mixture and separate individual isomers.

Some of the compounds may also exist in tautomeric forms. Such forms, although not explicitly indicated in the formulae described herein, are intended to be included within the scope of the present disclosure.

“Prodrug” or “pharmaceutically acceptable prodrug” refers to a compound that is metabolized, for example hydrolyzed or oxidized, in the host after administration to form the

compound of the present disclosure (e.g., compounds of formula I or II). Typical examples of prodrugs include compounds that have biologically labile or cleavable (protecting) groups on a functional moiety of the active compound. Prodrugs include compounds that can be oxidized, reduced, aminated, deaminated, hydroxylated, dehydroxylated, hydrolyzed, dehydrolyzed, alkylated, dealkylated, acylated, deacylated, phosphorylated, or dephosphorylated to produce the active compound. Examples of prodrugs using ester or phosphoramidate as biologically labile or cleavable (protecting) groups are disclosed in U.S. Patents 6,875,751, 7,585,851, and 7,964,580, the disclosures of which are incorporated herein by reference. The prodrugs of this disclosure are metabolized to produce a compound of formula I or II. The present disclosure includes within its scope, prodrugs of the compounds described herein. Conventional procedures for the selection and preparation of suitable prodrugs are described, for example, in "Design of Prodrugs" Ed. H. Bundgaard, Elsevier, 1985.


The phrase "pharmaceutically acceptable carrier" as used herein means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filter, diluent, excipient, solvent or encapsulating material useful for formulating a drug for medicinal or therapeutic use.

The term "Log of solubility", "LogS" or "logS" as used herein is used in the art to quantify the aqueous solubility of a compound. The aqueous solubility of a compound significantly affects its absorption and distribution characteristics. A low solubility often goes along with a poor absorption. LogS value is a unit stripped logarithm (base 10) of the solubility measured in mol/liter.

EXAMPLES

The invention now being generally described, it will be more readily understood by reference to the following examples which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

Example 1: Preparation of Exemplary Compounds



To the solution of indole-3-carboxaldehyde (2.8 mmol, 411 mg) in dry DMF (6 mL) were added Cu_2O (0.3 equiv, 0.84 mmol, 120 mg), K_2CO_3 (2.0 equiv, 5.6 mmol, 774 mg), and iodobenzene (2.0 equiv, 5.6 mmol, 624 μL) sequentially. The reaction was stirred and refluxed for 24 h, at which point TLC indicated that the reaction was completed. After it was cooled to 21 °C, the reaction mixture was filtrated through a Celite pad eluting with ethyl acetate. The filtrate was washed by saturated NaCl solution and organic phase was dried by sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes/EtOAc = 8:1) to provide the desired product. yield: 89%, 550.7 mg.


To the solution of 1-phenyl-indole-3-carbaldehyde (1 mmol, 221 mg) in ethanol (1 mL) were added ethyl 2-cyanoacetate (1.3 equiv, 1.3 mmol, 140 μL) and L-proline (40 mol%, 0.4 mmol, 58 mg). The reaction was stirred at 21 °C for 12 h and yellow solid precipitated gradually. After completion of the reaction, ice-cold water (2 mL) was added into the reaction vial. The solid was separated by Buchner funnel filtration and washed with water (2 mL \times 3) and dried to afford the desired product. yield: 95%, 300 mg.

To the solution of (*E*)-ethyl 2-cyano-3-(1-phenyl-1*H*-indol-3-yl)acrylate (0.32 mmol, 100 mg) in THF (2 mL) was added 0.5N LiOH solution (3 equiv, 0.6 mmol, 1.2 mL). The reaction mixture was stirred at 21 °C for 1 h. After reaction completion shown by TLC, THF was evaporated. Concentrated HCl was added dropwise to acidify the reaction mixture until the pH was lower than 1, meanwhile yellow solid precipitated. Ice-cold water (5 mL) was added to the reaction mixture and the solid was separated by Buchner funnel filtration and washed with water (5 mL \times 3). After dried by vacuum, the solid was washed by 2 mL of solvent mixture (hexanes/EtOAc = 5:1) 5 to 10 times and monitored by TLC until non-polar impurities disappear (The non-polar compound was the retro-Aldol condensation product, which can be recovered from the filtrate). Finally, the purity of the product was checked by NMR. yield: 65%, 60 mg.


(E)-2-Cyano-3-(1-phenyl-1H-indol-3-yl)acrylic acid (JXL001/UK5099)

¹H NMR (500 MHz, DMSO-d₆) δ 8.59 (s, 1H), 8.56 (s, 1H), 8.06 (m, 1H), 7.65 (m, 4H), 7.53 (m, 2H), 7.34 (m, 2H).

¹³C NMR (126 MHz, DMSO-d₆) δ 164.5, 145.6, 137.7, 136.3, 133.6, 130.5, 128.9, 128.0, 125.0, 124.9, 123.3, 119.9, 118.4, 111.9, 96.7.




The following compounds were synthesized by a route similar to that described for JXL001: JXL002, JXL003, JXL004, JXL005, JXL006, JXL007, JXL012, JXL013, JXL014, JXL021, JXL025, JXL026, JXL027, JXL028, JXL029, JXL035, JXL093.

**(E)-2-Cyano-3-(1H-indol-3-yl)acrylic acid (JXL002)**


¹H NMR (500 MHz, DMSO-d₆) δ 12.48 (s, 1H), 8.51 (s, 1H), 8.49 (s, 1H), 7.91 (d, *J* = 6.5 Hz, 1H), 7.53 (d, *J* = 7.0 Hz, 1H), 7.23 (m, 2H).

¹³C NMR (126 MHz, DMSO-d₆) δ 165.0, 146.5, 136.5, 132.4, 127.3, 123.9, 122.4, 118.9, 118.8, 113.2, 110.2, 94.0.

**Ethyl (E)-2-cyano-3-(1H-indol-3-yl)acrylate**

¹H NMR (500 MHz, CDCl₃) δ 12.55 (s, 1H), 8.53 (s, 1H), 8.52 (s, 1H), 7.92 (d, *J* = 7.6 Hz, 1H), 7.53 (d, *J* = 7.8 Hz, 1H), 7.26 (app. t, *J* = 7.4 Hz, 1H), 7.22 (app. t, *J* = 7.4 Hz, 1H), 4.24 (q, *J* = 7.0 Hz, 2H), 1.26 (t, *J* = 7.0 Hz, 3H).


^{13}C NMR (126 MHz, CDCl_3) δ 163.6, 147.0, 136.6, 133.0, 127.3, 124.0, 122.5, 118.9, 118.4, 113.3, 110.3, 92.6, 61.8, 14.5.



Ethyl (*E*)-2-cyano-3-(1-phenyl-1*H*-indol-3-yl)acrylate (JXL004)

^1H NMR (500 MHz, CDCl_3) δ 8.71 (s, 1H), 8.66 (s, 1H), 7.90 (d, $J = 7.2$ Hz, 1H), 7.54 (m, 6H), 7.36 (m, 2H), 4.39 (q, $J = 7.1$ Hz, 2H), 1.42 (t, $J = 7.1$ Hz, 3H).


^{13}C NMR (126 MHz, CDCl_3) δ 163.6, 145.6, 137.8, 136.4, 133.2, 129.9, 128.5, 124.8, 124.4, 123.0, 118.5, 117.9, 111.6, 111.5, 95.4, 62.0, 14.3.



Ethyl (*E*)-3-(6-chloro-1-phenyl-1*H*-indol-3-yl)-2-cyanoacrylate (JXL005)

^1H NMR (500 MHz, CDCl_3) δ 8.67 (s, 1H), 8.58 (s, 1H), 7.81 (d, $J = 8.5$ Hz, 1H), 7.60 (m, 2H), 7.52 (m, 4H), 7.34 (d, $J = 8.4$ Hz, 1H), 4.39 (q, $J = 7.1$ Hz, 2H), 1.41 (t, $J = 7.1$ Hz, 3H).

^{13}C NMR (126 MHz, CDCl_3) δ 163.4, 145.1, 137.3, 136.8, 133.5, 130.5, 130.0, 128.8, 126.8, 124.8, 123.6, 119.5, 117.5, 111.6, 111.4, 96.4, 62.1, 14.2.




Ethyl (*E*)-2-cyano-3-(1-(2-methoxyphenyl)-1*H*-indol-3-yl)acrylate (JXL006)

^1H NMR (500 MHz, CDCl_3) δ 8.67 (s, 1H), 8.66 (s, 1H), 7.89 (d, $J = 7.8$ Hz, 1H), 7.49 (app. t, $J = 8.6$ Hz, 1H), 7.41 (d, $J = 7.6$ Hz, 1H), 7.35 (app. t, $J = 7.3$ Hz, 1H), 7.30 (t, J

= 7.5 Hz, 1H), 7.23 (d, J = 8.1 Hz, 1H), 7.13 (m, 2H), 4.39 (q, J = 7.1 Hz, 2H), 3.81 (s, 3H), 1.41 (t, J = 7.1 Hz, 3H).


^{13}C NMR (126 MHz, CDCl_3) δ 163.8, 154.2, 146.0, 137.2, 135.2, 130.3, 127.8, 126.0, 124.0, 122.6, 120.8, 118.2, 118.0, 112.3, 111.8, 111.0, 94.7, 61.8, 55.7, 14.3.



Ethyl (*E*)-2-cyano-3-(1-(4-methoxyphenyl)-1*H*-indol-3-yl)acrylate (JXL007)

^1H NMR (500 MHz, CDCl_3) δ 8.65 (s, 1H), 8.64 (s, 1H), 7.89 (d, J = 7.2 Hz, 1H), 7.44 (m, 3H), 7.35 (m, 2H), 7.07 (d, J = 8.8 Hz, 2H), 4.38 (q, J = 7.1 Hz, 2H), 3.90 (s, 3H), 1.41 (t, J = 7.1 Hz, 3H).


^{13}C NMR (126 MHz, CDCl_3) δ 163.8, 159.5, 145.6, 136.8, 133.5, 130.6, 128.3, 126.2, 124.2, 122.9, 118.4, 118.0, 115.0, 111.5, 111.1, 94.9, 61.9, 55.6, 14.2.



(*E*)-2-Cyano-3-(1-(4-methoxyphenyl)-1*H*-indol-3-yl)acrylic acid (JXL012)

^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.54 (s, 1H), 8.52 (s, 1H), 8.05 (d, J = 7.7 Hz, 1H), 7.58 (app. d, J = 8.7 Hz, 2H), 7.44 (m, 1H), 7.33 (m, 2H), 7.16 (app. d, J = 8.7 Hz, 2H), 3.82 (s, 3H).


^{13}C NMR (126 MHz, $\text{DMSO}-d_6$) δ 164.7, 160.5, 136.7, 133.9, 130.5, 127.8, 126.6, 124.8, 123.2, 120.0, 119.5, 118.6, 115.6, 115.4, 111.9, 110.9, 55.9.



(E)-3-(6-Chloro-1-phenyl-1H-indol-3-yl)-2-cyanoacrylic acid (JXL013)

¹H NMR (500 MHz, DMSO-*d*₆) δ 13.58 (br. s, 1H), 8.59 (s, 1H), 8.54 (s, 1H), 8.11 (d, *J* = 7.5 Hz, 1H), 7.67 (m, 4H), 7.53 (m, 2H), 7.35 (d, *J* = 7.5 Hz, 1H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.3, 145.4, 137.3, 136.7, 134.4, 130.6, 129.6, 129.1, 126.7, 125.1, 123.5, 121.5, 118.2, 111.6, 111.3, 97.9.



(E)-2-Cyano-3-(1-(2-methoxyphenyl)-1H-indol-3-yl)acrylic acid (JXL014)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.55 (s, 1H), 8.47 (s, 1H), 8.02 (d, *J* = 7.4 Hz, 1H), 7.54 (m, 2H), 7.31 (m, 3H), 7.15 (m, 2H), 3.74 (s, 3H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.6, 154.2, 145.8, 137.3, 135.1, 131.1, 128.2, 127.3, 125.7, 124.6, 123.0, 121.5, 119.3, 118.5, 113.6, 112.2, 110.8, 96.1, 56.2.



2-((1-Phenyl-1H-indol-3-yl)methylene)malononitrile (JXL021)

¹H NMR (500 MHz, CDCl₃) δ 8.62 (s, 1H), 8.13 (s, 1H), 7.80 (d, *J* = 7.4 Hz, 1H), 7.61 (m, 2H), 7.53 (m, 4H), 7.40 (m, 2H).


¹³C NMR (126 MHz, CDCl₃) δ 149.8, 137.3, 136.5, 133.6, 130.1, 128.9, 127.7, 124.9, 124.8, 123.7, 118.2, 115.1, 115.0, 111.9, 111.8, 73.7.



2-(Ethoxycarbonyl)-3-(1-phenyl-1H-indol-3-yl)acrylic acid (a mixture of *E/Z* isomers, 1:1 ratio) (JXL025)

¹H NMR (500 MHz, DMSO-d₆) δ 13.08 (br. s, 1H), 7.87 (m, 3H), 7.61 (m, 4H), 7.52 (m, 2H), 7.30 (m, 2H), 4.26 (m, 2H), 1.23 (m, 3H).


¹³C NMR (126 MHz, DMSO-d₆) δ 168.9, 167.8, 166.1, 164.9, 138.3, 136.1, 132.9, 132.1, 131.1, 130.9, 130.8, 128.5, 124.9, 124.4, 122.6, 122.2, 119.5, 111.6, 111.0, 61.7, 61.3, 14.7, 14.4.



(*E*)-2-Cyano-3-(4-fluoro-1-phenyl-1H-indol-3-yl)acrylic acid (JXL026)

¹H NMR (500 MHz, DMSO-d₆) δ 8.65 (s, 1H), 8.61 (s, 1H), 7.67 (m, 4H), 7.57 (m, 1H), 7.36 (m, 2H), 7.19 (m, 1H).


¹³C NMR (126 MHz, DMSO-d₆) δ 164.4, 156.8 (d, *J*_{C-F} = 245.6 Hz), 146.4, 138.7, 137.6, 133.4, 130.7, 129.5, 125.9, 125.4, 118.2, 116.3, 116.2, 109.5 (d, *J*_{C-F} = 34.5 Hz), 109.2 (d, *J*_{C-F} = 23.2 Hz), 98.0.



(*E*)-2-Cyano-3-(6-fluoro-1-phenyl-1H-indol-3-yl)acrylic acid (JXL027)

¹H NMR (500 MHz, DMSO-d₆) δ 13.59 (br. s, 1H), 8.62 (s, 1H), 8.59 (s, 1H), 8.16 (m, 1H), 7.66 (m, 4H), 7.56 (m, 1H), 7.36 (d, *J* = 9.2 Hz, 1H), 7.25 (m, 1H).

¹³C NMR (126 MHz, DMSO-d₆) δ 164.6, 160.8 (d, *J*_{C-F} = 240.0 Hz), 145.7, 137.6, 136.6, 134.4, 130.8, 129.1, 125.0, 124.7, 121.6, 118.4, 111.8 (d, *J*_{C-F} = 24.2 Hz), 111.5, 98.7 (d, *J*_{C-F} = 26.2 Hz), 97.7.




JXL028

(E)-2-Cyano-3-(7-fluoro-1-phenyl-1H-indol-3-yl)acrylic acid (JXL028)

¹H NMR (500 MHz, DMSO-*d*₆) δ 13.62 (br. s, 1H), 8.56 (s, 1H), 8.47 (s, 1H), 7.89 (br. s, 1H), 7.61 (m, 5H), 7.30 (br. s, 1H), 7.17 (br. s, 1H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.4, 149.7 (d, J_{C-F} = 247.5 Hz), 145.5, 139.0, 135.5, 131.9, 129.9, 129.3, 126.2, 124.3, 124.0, 118.3, 115.9, 111.8, 110.9 (d, J_{C-F} = 17.4 Hz), 98.1.




JXL029

(E)-2-Cyano-3-(5-fluoro-1-phenyl-1H-indol-3-yl)acrylic acid (JXL029)

¹H NMR (500 MHz, DMSO-*d*₆) δ 13.57 (br. s, 1H), 8.67 (s, 1H), 8.60 (s, 1H), 8.01 (d, J = 9.0 Hz, 1H), 7.70 (m, 4H), 7.58 (m, 2H), 7.24 (m, 1H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.6, 159.4 (d, J_{C-F} = 237.8 Hz), 145.8, 137.7, 135.1, 133.1, 130.7, 129.2, 129.1, 125.2, 118.5, 113.7, 113.2 (d, J_{C-F} = 26.5 Hz), 111.4, 105.7 (d, J_{C-F} = 24.2 Hz), 97.2.




JXL035

(E)-3-(4-Chloro-1-phenyl-1H-indol-3-yl)-2-cyanoacrylic acid (JXL035)

¹H NMR (500 MHz, DMSO-*d*₆) δ 9.22 (s, 1H), 8.68 (s, 1H), 7.64 (m, 3H), 7.59 (m, 2H), 7.48 (d, J = 7.6 Hz, 1H), 7.38 (m, 1H), 7.32 (m, 1H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.5, 146.4, 138.0, 137.4, 134.5, 130.7, 130.5, 129.5, 125.7, 125.3, 124.9, 123.7, 118.4, 111.7, 110.9, 97.5.




JXL093

Ethyl (*E*)-3-(4-chloro-1-phenyl-1*H*-indol-3-yl)-2-cyanoacrylate (JXL093)

¹H NMR (500 MHz, CDCl₃) δ 9.51 (s, 1H), 8.78 (s, 1H), 7.59 (m, 2H), 7.51 (m, 3H), 7.40 (dd, *J* = 8.3, 0.8 Hz, 1H), 7.32 (dd, *J* = 7.7, 0.8 Hz, 1H), 7.22 (app. t, *J* = 8.0 Hz, 1H), 4.38 (q, *J* = 7.1 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H).

¹³C NMR (126 MHz, CDCl₃) δ 163.6, 147.6, 138.1, 137.4, 134.2, 130.1, 129.0, 126.9, 125.3, 124.6, 124.5, 124.3, 118.0, 111.8, 110.6, 96.0, 62.1, 14.3.

Experimental details for the synthesis of JXL020

To the solution of indole-3-carboxaldehyde (3 mmol, 435 mg) in dry DMF (6 mL) were added 3,5-bis(trifluoromethyl)benzyl bromide (1.2 equiv, 3.6 mmol, 660 μL) and KOH (1.2 equiv, 3.6 mmol, 200 mg) at 0 °C. The reaction mixture was stirred at 21 °C for 2 h. After the reaction completion shown by TLC, water (6 mL) was added to the reaction vial. The reaction mixture was extracted by dichloromethane (15 mL × 3). The combined organic layer was dried by sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes/EtOAc = 8:1) to provide the desired product. yield: 90%, 1001.7 mg.

To the solution of 1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indole-3-carbaldehyde (1 mmol, 371 mg) in ethanol (1 mL) were added ethyl 2-cyanoacetate (1.3 equiv, 1.3 mmol, 140 μL) and *L*-proline (40 mol%, 0.4 mmol, 58 mg). The reaction was stirred at 21 °C for 12 h and yellow solid precipitated gradually. After completion of the reaction, ice-cold water (2 mL) was added into the reaction vial. The solid was separated by Buchner funnel filtration and washed with water (2 mL × 3) and dried to afford the desired product. yield: 93%, 433 mg.


To the solution of (*E*)-ethyl 3-(1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylate (0.21 mmol, 100 mg) in THF (2 mL) was added 0.5N LiOH solution (3 equiv, 0.4 mmol, 0.8 mL). The reaction mixture was stirred at 21 °C for 1 h. After reaction completion shown by TLC, THF was evaporated. Concentrated HCl was added dropwise to acidify the reaction mixture until pH was lower than 1, meanwhile yellow solid precipitated. Ice-cold water (5 mL) was added to the reaction mixture and the solid was separated by Buchner funnel filtration and washed with water (5 mL × 3). After dried by vacuum, the solid was washed by 2 mL of solvent mixture (hexanes/EtOAc = 5:1) 5 to 10 times and monitored by TLC until non-polar impurities disappear (The non-polar compound was the retro-Aldol condensation product, which can be recovered from the filtrate). Finally, the purity of the product was checked by NMR. yield: 55%, 52 mg.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylic acid (JXL020)

¹H NMR (500 MHz, DMSO-*d*₆) δ 13.37 (br. s, 1H), 8.75 (s, 1H), 8.48 (s, 1H), 7.99 (m, 4H), 7.65 (s, 1H), 7.28 (m, 2H), 5.83 (s, 2H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.7, 145.7, 140.3, 136.3, 134.8, 131.1, 130.8 (q, *J* = 31.1 Hz), 128.9, 128.7, 127.9, 124.8, 124.3, 122.9 (q, *J* = 273.4 Hz), 122.2, 119.3, 118.3, 95.6, 49.2.


The following compounds were synthesized by a route similar to that described for JXL020: JXL008, JXL009, JXL010, JXL011, JXL015, JXL016, JXL017, JXL018, JXL019, JXL036, JXL037, JXL038, JXL039, JXL040, JXL041, JXL050, JXL051, JXL052, JXL053, JXL054, JXL055, JXL56, JXL057, JXL058, JXL059, JXL060, JXL061, JXL062, JXL063, JXL064, JXL065, JXL066, JXL068, JXL069, JXL072, JXL073, JXL076, JXL077, JXL078, JXL081, JXL082, JXL087, JXL088, JXL089, JXL090, JXL091.



Ethyl (*E*)-2-cyano-3-(1-(4-fluorobenzyl)-1*H*-indol-3-yl)acrylate (JXL008)

¹H NMR (500 MHz, CDCl₃) δ 8.60 (app. s, 2H), 7.85 (d, *J* = 6.8 Hz, 1H), 7.32 (m, 3H), 7.15 (m, 2H), 7.03 (app. t, 2H), 5.39 (s, 2H), 4.37 (q, *J* = 7.1 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H).


¹³C NMR (126 MHz, CDCl₃) δ 163.7, 162.5 (d, *J*_{C-F} = 247.7 Hz), 145.7, 136.1, 133.8, 130.9, 128.6, 128.5, 124.0, 122.7, 118.6, 118.0, 116.0 (d, *J*_{C-F} = 21.9 Hz), 110.9, 110.4, 94.6, 61.9, 50.7, 14.2.



Ethyl (*E*)-2-cyano-3-(1-(3,4-difluorobenzyl)-1*H*-indol-3-yl)acrylate (JXL009)

¹H NMR (500 MHz, CDCl₃) δ 8.60 (s, 1H), 8.59 (s, 1H), 7.86 (d, *J* = 7.8 Hz, 1H), 7.33 (m, 2H), 7.28 (s, 1H), 7.13 (m, 1H), 6.95 (m, 1H), 6.89 (m, 1H), 5.39 (s, 2H), 4.38 (q, *J* = 7.1 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H).


¹³C NMR (126 MHz, CDCl₃) δ 163.7, 150.7 (dd, *J* = 251.2, 13.2 Hz), 150.2 (dd, *J* = 250.4, 12.6 Hz), 145.7, 136.1, 133.7, 132.3, 128.6, 124.3, 122.9, 122.7, 120.0, 118.8, 118.0 (d, *J* = 17.5 Hz), 115.9 (d, *J* = 18.0 Hz), 110.8, 110.6, 95.2, 62.1, 50.4, 14.4.



Ethyl (*E*)-2-cyano-3-(1-(3,5-difluorobenzyl)-1*H*-indol-3-yl)acrylate (JXL010)

¹H NMR (500 MHz, CDCl₃) δ 8.61 (s, 1H), 8.59 (s, 1H), 7.87 (d, *J* = 7.1 Hz, 1H), 7.33 (m, 2H), 7.26 (d, *J* = 7.2 Hz, 1H), 6.75 (app. t, *J* = 8.7 Hz, 1H), 6.64 (app. d, *J* = 5.7 Hz, 2H), 5.41 (s, 2H), 4.38 (q, *J* = 7.1 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H).


¹³C NMR (126 MHz, CDCl₃) δ 163.6, 163.3 (dd, *J*_{c-f} = 251.0, 12.5 Hz), 145.5, 139.2, 136.0, 133.6, 128.4, 124.3, 122.9, 118.7, 118.0, 110.7, 110.6, 109.5 (dd, *J*_{c-f} = 19.9, 6.4 Hz), 103.8 (t, *J*_{c-f} = 25.2 Hz), 95.3, 62.0, 50.4, 14.2.



Ethyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylate (JXL011)

¹H NMR (500 MHz, CDCl₃) δ 8.62 (s, 1H), 8.61 (s, 1H), 7.90 (d, *J* = 7.7 Hz, 1H), 7.85 (s, 1H), 7.57 (s, 2H), 7.35 (m, 2H), 7.23 (d, *J* = 7.8 Hz, 1H), 5.56 (s, 2H), 4.38 (q, *J* = 7.1 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H).


¹³C NMR (126 MHz, CDCl₃) δ 163.6, 145.6, 138.1, 136.0, 133.4, 132.7 (q, *J* = 33.8 Hz), 128.6, 126.8, 124.7, 123.2, 122.9 (q, *J* = 273.4 Hz), 122.6, 119.0, 118.0, 111.2, 110.4, 95.9, 62.1, 50.5, 14.3.



(E)-2-Cyano-3-(1-(4-fluorobenzyl)-1H-indol-3-yl)acrylic acid (JXL015)

¹H NMR (500 MHz, DMSO-d₆) δ 8.64 (s, 1H), 8.46 (s, 1H), 7.93 (d, *J* = 7.1 Hz, 1H), 7.61 (d, *J* = 7.3 Hz, 1H), 7.33 (m, 2H), 7.26 (m, 2H), 7.16 (m, 2H), 5.60 (s, 2H).


¹³C NMR (126 MHz, DMSO-d₆) δ 164.8, 162.0 (d, *J*_{C-F} = 244.3 Hz), 145.6, 136.4, 134.6, 133.1, 130.0, 128.0, 124.1, 122.8, 119.2, 118.5, 116.0 (d, *J*_{C-F} = 21.7 Hz), 112.0, 109.8, 95.0, 49.6.



(E)-2-Cyano-3-(1-(3,4-difluorobenzyl)-1H-indol-3-yl)acrylic acid (JXL016)

¹H NMR (500 MHz, DMSO-d₆) δ 13.34 (br. s, 1H), 8.62 (s, 1H), 8.47 (s, 1H), 7.94 (d, *J* = 7.3 Hz, 1H), 7.61 (d, *J* = 7.6 Hz, 1H), 7.40 (m, 2H), 7.27 (m, 2H), 7.10 (br. s, 1H), 5.61 (s, 2H).


¹³C NMR (126 MHz, DMSO-d₆) δ 164.8, 149.7 (dd, *J*_{C-F} = 253.3, 13.6 Hz), 149.4 (dd, *J*_{C-F} = 246.3, 11.6 Hz), 145.7, 136.4, 134.7, 134.5, 128.0, 124.7 (dd, *J*_{C-F} = 5.9, 3.0 Hz), 124.2, 122.8, 119.2, 118.4, 118.3 (d, *J*_{C-F} = 17.0 Hz), 117.1 (d, *J*_{C-F} = 17.6 Hz), 112.0, 109.9, 95.0, 49.3.



(E)-2-Cyano-3-(1-(3,5-difluorobenzyl)-1H-indol-3-yl)acrylic acid (JXL017)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.67 (s, 1H), 8.48 (s, 1H), 7.95 (d, *J* = 4.8 Hz, 1H), 7.59 (d, *J* = 4.4 Hz, 1H), 7.27 (m, 2H), 7.15 (s, 1H), 6.98 (br. s, 2H), 5.65 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.8, 162.9 (dd, *J*_{c-f} = 247.0, 12.8 Hz), 145.6, 141.4, 136.4, 134.8, 128.0, 124.2, 122.9, 119.3, 118.4, 111.9, 111.0 (d, *J*_{c-f} = 26.1 Hz), 110.8, 103.8 (t, *J*_{c-f} = 26.5 Hz), 95.5, 49.5.



Ethyl (E)-3-(1-benzyl-1H-indol-3-yl)-2-cyanoacrylate (JXL018)

¹H NMR (500 MHz, CDCl₃) δ 8.61 (s, 1H), 8.60 (s, 1H), 7.85 (d, *J* = 7.9 Hz, 1H), 7.33 (m, 6H), 7.17 (m, 2H), 5.42 (s, 2H), 4.37 (q, *J* = 7.1 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H).

¹³C NMR (126 MHz, CDCl₃) δ 163.8, 145.7, 136.2, 135.1, 134.0, 129.0, 128.5, 128.2, 126.8, 124.0, 122.6, 118.5, 118.1, 111.0, 110.3, 94.3, 61.8, 51.4, 14.2.




JXL019

(E)-3-(1-Benzyl-1H-indol-3-yl)-2-cyanoacrylic acid (JXL019)

$^1\text{H NMR}$ (500 MHz, DMSO- d_6) δ 13.34 (br. s, 1H), 8.65 (s, 1H), 8.48 (s, 1H), 7.93 (d, $J = 6.9$ Hz, 1H), 7.60 (d, $J = 6.8$ Hz, 1H), 7.25 (m, 7H), 5.62 (s, 2H).

$^{13}\text{C NMR}$ (126 MHz, DMSO- d_6) δ 164.8, 145.7, 136.8, 136.5, 134.7, 129.1, 128.2, 127.6, 124.0, 122.7, 120.0, 119.2, 118.5, 112.1, 109.7, 94.7, 50.4.




JXL036

(E)-2-Cyano-3-(1-(3,4-difluorobenzyl)-5-fluoro-1H-indol-3-yl)acrylic acid (JXL036)

$^1\text{H NMR}$ (500 MHz, DMSO- d_6) δ 8.69 (s, 1H), 8.47 (s, 1H), 7.84 (d, $J = 9.6$ Hz, 1H), 7.63 (dd, $J = 8.9, 4.3$ Hz, 1H), 7.41 (m, 2H), 7.14 (m, 2H), 5.61 (s, 2H).


$^{13}\text{C NMR}$ (126 MHz, DMSO- d_6) δ 164.9, 159.4 (d, $J_{\text{C-F}} = 237.3$ Hz), 149.8 (dd, $J_{\text{C-F}} = 247.1, 12.7$ Hz), 149.5 (dd, $J_{\text{C-F}} = 246.6, 12.3$ Hz), 146.0, 136.1, 134.5, 133.1, 129.1, 125.0, 118.5, 118.4, 117.4, 113.6, 112.5 (d, $J_{\text{C-F}} = 26.2$ Hz), 110.1, 105.2 (d, $J_{\text{C-F}} = 25.1$ Hz), 95.5, 49.7.



(E)-2-Cyano-3-(1-(3,5-difluorobenzyl)-5-fluoro-1H-indol-3-yl)acrylic acid (JXL037)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.71 (s, 1H), 8.48 (s, 1H), 7.85 (dd, *J* = 9.6, 2.0 Hz, 1H), 7.62 (dd, *J* = 8.9, 4.3 Hz, 1H), 7.16 (m, 2H), 7.00 (app. d, *J* = 6.2 Hz, 2H), 5.65 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.9, 163.0 (d, *J*_{C-F} = 247.7 Hz), 159.3 (d, *J*_{C-F} = 237.6 Hz), 145.9, 141.4, 136.3, 133.2, 129.1, 118.5, 113.6, 112.6 (d, *J*_{C-F} = 26.3 Hz), 111.3, 110.2, 105.3 (d, *J*_{C-F} = 24.9 Hz), 104.0 (t, *J*_{C-F} = 25.2 Hz), 95.9, 49.8.



(E)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-5-fluoro-1H-indol-3-yl)-2-cyanoacrylic acid (JXL038)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.80 (s, 1H), 8.49 (s, 1H), 8.06 (s, 1H), 8.04 (s, 2H), 7.86 (dd, *J* = 9.6, 2.1 Hz, 1H), 7.69 (dd, *J* = 8.9, 4.3 Hz, 1H), 7.17 (dt, *J* = 9.0, 2.2 Hz, 1H), 5.83 (s, 2H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.8, 159.3 (d, *J*_{C-F} = 237.5 Hz), 145.9, 140.3, 136.2, 133.1, 131.1 (q, *J*_{C-F} = 33.1 Hz), 129.1, 123.6 (q, *J*_{C-F} = 272.2 Hz), 118.4, 113.5, 112.7, 112.5, 110.4, 105.5, 105.3, 96.1, 49.6.




JXL039

(E)-2-Cyano-3-(1-(3,4-difluorobenzyl)-6-fluoro-1H-indol-3-yl)acrylic acid (JXL039)

¹H NMR (500 MHz, DMSO-d₆) δ 8.64 (s, 1H), 8.46 (s, 1H), 7.99 (dd, *J* = 8.7, 5.1 Hz, 1H), 7.58 (dd, *J* = 9.8, 1.8 Hz, 1H), 7.43 (m, 2H), 7.12 (m, 2H), 5.57 (s, 2H).

¹³C NMR (126 MHz, DMSO-d₆) δ 164.8, 160.3 (d, *J*_{C-F} = 239.3 Hz), 149.8 (dd, *J*_{C-F} = 239.3, 25.2 Hz), 149.6 (dd, *J*_{C-F} = 246.3, 25.2 Hz), 145.8, 136.8, 135.3, 134.5, 125.1, 125.0, 124.6, 121.1, 118.5, 117.5, 111.3 (d, *J*_{C-F} = 23.9 Hz), 110.2, 98.7 (d, *J*_{C-F} = 26.5 Hz), 96.2, 49.4.




JXL040

(E)-2-Cyano-3-(1-(3,5-difluorobenzyl)-6-fluoro-1H-indol-3-yl)acrylic acid (JXL040)

¹H NMR (500 MHz, DMSO-d₆) δ 8.65 (s, 1H), 8.46 (s, 1H), 8.00 (dd, *J* = 8.6, 5.2 Hz, 1H), 7.57 (d, *J* = 9.7 Hz, 1H), 7.15 (m, 2H), 7.03 (s, 1H), 7.02 (s, 1H), 5.62 (s, 2H).


¹³C NMR (126 MHz, DMSO-d₆) δ 164.8, 163.0 (d, *J*_{C-F} = 239.4 Hz), 162.9 (d, *J*_{C-F} = 248.6 Hz), 160.4 (d, *J*_{C-F} = 239.4 Hz), 145.5, 141.3, 136.8, 135.2, 124.6, 121.1, 118.4, 111.3, 111.1, 110.3, 104.0 (t, *J*_{C-F} = 25.2 Hz), 98.7 (d, *J*_{C-F} = 26.5 Hz), 96.9, 49.6.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-6-fluoro-1H-indol-3-yl)-2-cyanoacrylic acid (JXL041)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.74 (s, 1H), 8.48 (s, 1H), 8.06 (app. s, 3H), 8.01 (dd, *J* = 8.7, 5.1 Hz, 1H), 7.66 (dd, *J* = 9.8, 2.0 Hz, 1H), 7.13 (dt, *J* = 9.3, 2.1 Hz, 1H), 5.78 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.7, 159.9 (d, *J*_{C-F} = 264.6 Hz), 145.8, 140.2, 136.8, 135.3, 131.1 (q, *J*_{C-F} = 33.3 Hz), 129.1, 124.5, 123.6 (q, *J*_{C-F} = 273.7 Hz), 122.5, 121.2, 118.4, 111.4 (d, *J*_{C-F} = 25.2 Hz), 110.4, 98.6 (d, *J*_{C-F} = 27.2 Hz), 96.7, 49.4.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-chloro-1H-indol-3-yl)-2-cyanoacrylic acid (JXL050)

¹H NMR (500 MHz, DMSO-*d*₆) δ 9.20 (s, 1H), 8.90 (s, 1H), 8.06 (m, 3H), 7.72 (d, *J* = 7.5 Hz, 1H), 7.32 (m, 2H), 5.86 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.7, 146.7, 140.0, 138.0, 135.6, 131.1 (q, *J*_{C-F} = 33.1 Hz), 129.1, 125.6, 125.1, 124.7, 124.5, 123.7, 123.6 (q, *J*_{C-F} = 273.7 Hz), 122.5, 111.6, 109.9, 96.7, 49.6.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-bromo-1H-indol-3-yl)-2-cyanoacrylic acid (JXL051)

¹H NMR (500 MHz, DMSO-*d*₆) δ 9.37 (s, 1H), 8.91 (s, 1H), 8.07 (app. s, 3H), 7.77 (d, *J* = 8.2 Hz, 1H), 7.51 (d, *J* = 7.6 Hz, 1H), 7.23 (t, *J* = 8.0 Hz, 1H), 5.85 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.7, 146.3, 140.0, 138.0, 135.8, 131.0 (q, *J*_{c-f} = 33.1 Hz), 129.1, 127.9, 125.4, 124.9, 123.7 (q, *J*_{c-f} = 273.2 Hz), 122.5, 118.2, 113.6, 122.1, 110.2, 96.6, 49.5.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-fluoro-1H-indol-3-yl)-2-cyanoacrylic acid (JXL052)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.80 (s, 1H), 8.58 (s, 1H), 8.05 (app. s, 3H), 7.54 (d, *J* = 8.2 Hz, 1H), 7.29 (d, *J* = 12.9 Hz, 1H), 7.09 (dd, *J* = 11.1, 8.2 Hz, 1H), 5.85 (s, 2H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.5, 155.7 (d, *J*_{c-f} = 239.4 Hz), 146.5, 140.1, 138.8, 134.5, 131.1 (q, *J*_{c-f} = 33.0 Hz), 129.0, 125.1 (d, *J*_{c-f} = 7.6 Hz), 124.7, 123.6 (q, *J*_{c-f} = 273.5 Hz), 122.5, 118.1, 116.1 (d, *J*_{c-f} = 18.5 Hz), 108.8, 108.5, 97.2, 49.7.




JXL053

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-7-fluoro-1H-indol-3-yl)-2-cyanoacrylic acid (JXL053)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.73 (br. s, 1H), 8.48 (br. s, 1H), 8.06 (br. s, 1H), 7.89 (br. s, 2H), 7.78 (br. s, *J* = 7.4 Hz, 1H), 7.22 (br. s, 1H), 7.10 (br. s, 1H), 5.89 (s, 2H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.6, 159.7 (d, *J*_{C-F} = 245.7 Hz), 145.6, 141.0, 136.2, 132.1, 130.0 (q, *J*_{C-F} = 33.0 Hz), 128.4, 123.9, 123.8, 123.6 (q, *J*_{C-F} = 273.3 Hz), 122.4 (d, *J*_{C-F} = 18.9 Hz), 118.1, 115.8, 110.6, 110.2 (d, *J*_{C-F} = 18.9 Hz), 97.3, 52.0.




JXL054

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-5-chloro-1H-indol-3-yl)-2-cyanoacrylic acid (JXL054)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.78 (s, 1H), 8.51 (s, 1H), 8.12 (s, 1H), 8.06 (s, 1H), 8.03 (s, 2H), 7.70 (d, *J* = 7.4 Hz, 1H), 7.33 (d, *J* = 7.0 Hz, 1H), 5.83 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.7, 145.7, 140.2, 135.9, 135.1, 131.1 (q, *J*_{C-F} = 33.0 Hz), 129.3, 129.0, 127.9, 124.4, 123.6 (q, *J*_{C-F} = 274.0 Hz), 122.5, 119.5, 118.4, 113.6, 109.9, 96.9, 49.5.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-cyano-1H-indol-3-yl)-2-cyanoacrylic acid (JXL055)

¹H NMR (500 MHz, DMSO-*d*₆) δ 9.03 (s, 1H), 9.00 (s, 1H), 8.09 (m, 4H), 7.80 (d, *J* = 7.3 Hz, 1H), 7.47 (t, *J* = 7.8 Hz, 1H), 5.90 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.4, 144.4, 139.9, 136.7, 131.1 (q, *J*_{C-F} = 33.0 Hz), 129.7, 129.2, 126.6, 124.3, 123.6 (q, *J*_{C-F} = 273.7 Hz), 122.7, 118.6, 117.9, 117.8, 109.3, 101.7, 98.1, 49.5.



(E)-1-(3,5-Bis(trifluoromethyl)benzyl)-3-(2-carboxy-2-cyanovinyl)-1H-indole-4-carboxylic acid (JXL056)

¹H NMR (500 MHz, DMSO-*d*₆) δ 9.26 (br. s, 1H), 8.88 (br. s, 1H), 8.05 (app. s, 3H), 7.94 (br. s, 1H), 7.76 (br. s, 1H), 7.37 (br. s, 1H), 5.85 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 169.3, 165.0, 150.6, 140.2, 137.7, 136.0, 131.1 (q, *J*_{C-F} = 33.0 Hz), 129.0, 125.9, 125.2, 124.7, 123.6 (q, *J*_{C-F} = 273.7 Hz), 123.5, 122.5, 118.3, 116.1, 110.1, 96.4, 49.3.



(E)-3-(4-(Benzyloxy)-1-(3,5-bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylic acid (JXL057)

¹H NMR (500 MHz, DMSO-d₆) δ 9.15 (s, 1H), 8.72 (s, 1H), 8.04 (s, 1H), 7.99 (s, 2H), 7.54 (br. s, 3H), 7.37 (br. s, 2H), 7.28 (m, 2H), 6.95 (s, 1H), 5.81 (s, 2H), 5.28 (s, 2H).


¹³C NMR (126 MHz, DMSO-d₆) δ 165.0, 153.7, 148.6, 140.5, 137.9, 133.4, 131.2 (q, *J*_{c-f} = 32.8 Hz), 128.9, 128.8, 128.1, 127.5, 125.4, 124.7, 123.6 (q, *J*_{c-f} = 273.7 Hz), 122.3, 118.5, 117.0, 110.5, 105.6, 105.2, 95.5, 70.0, 49.5.



(E)-3-(6-(Benzyloxy)-1-(3,5-bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylic acid (JXL058)

¹H NMR (500 MHz, DMSO-d₆) δ 8.63 (s, 1H), 8.43 (s, 1H), 8.05 (s, 1H), 8.03 (s, 2H), 7.86 (d, *J* = 8.5 Hz, 1H), 7.32 (m, 6H), 6.97 (d, *J* = 8.3 Hz, 1H), 5.77 (s, 2H), 5.09 (s, 2H).

¹³C NMR (126 MHz, DMSO-d₆) δ 165.0, 156.7, 146.1, 140.5, 137.5, 137.3, 134.3, 131.0 (q, *J*_{c-f} = 32.8 Hz), 128.9, 128.8, 128.3, 128.2, 126.9, 123.6 (q, *J*_{c-f} = 273.4 Hz), 122.4, 121.9, 118.5, 113.3, 110.5, 96.8, 95.6, 70.2, 49.2.




JXL059

(E)-3-(7-(Benzyloxy)-1-(3,5-bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylic acid (JXL059)

¹H NMR (500 MHz, DMSO-d₆) δ 8.61 (br. s, 1H), 8.45 (br. s, 1H), 7.97 (br. s, 1H), 7.60 (br. s, 2H), 7.51 (br. s, 1H), 7.25 (br. s, 2H), 7.16 (br. s, 2H), 6.91 (br. s, 1H), 5.94 (s, 2H), 5.13 (s, 2H).

¹³C NMR (126 MHz, DMSO-d₆) δ 164.8, 146.6, 145.7, 142.3, 136.6, 135.4, 130.7 (q, *J*_{C-F} = 32.8 Hz), 130.5, 128.8, 128.4, 128.0, 127.5, 125.7, 124.1, 123.6 (q, *J*_{C-F} = 273.7 Hz), 121.8, 118.3, 111.8, 110.3, 107.1, 96.2, 70.3, 52.4.




JXL060

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-methoxy-1H-indol-3-yl)-2-cyanoacrylic acid (JXL060)

¹H NMR (500 MHz, DMSO-d₆) δ 8.99 (s, 1H), 8.71 (s, 1H), 8.05 (s, 1H), 8.00 (s, 2H), 7.24 (app. s, 2H), 6.82 (d, *J* = 6.0 Hz, 1H), 5.81 (s, 2H), 3.92 (s, 3H).


¹³C NMR (126 MHz, DMSO-d₆) δ 165.0, 154.8, 148.6, 140.5, 137.8, 133.3, 131.0 (q, *J*_{C-F} = 32.8 Hz), 128.9, 125.4, 123.6 (q, *J*_{C-F} = 273.7 Hz), 122.5, 118.5, 116.8, 110.5, 105.0, 104.3, 95.3, 56.2, 49.5.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-5-bromo-1H-indol-3-yl)-2-cyanoacrylic acid (JXL061)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.75 (s, 1H), 8.49 (s, 1H), 8.25 (s, 1H), 8.06 (s, 1H), 8.02 (m, 3H), 7.64 (app. s, 1H), 7.44 (app. s, 1H), 5.82 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.7, 145.3, 140.3, 135.5, 135.3, 131.1 (q, *J*_{c-f} = 32.8 Hz), 129.8, 128.9, 127.0, 124.7, 123.6 (q, *J*_{c-f} = 274.0 Hz), 122.5, 122.3, 118.5, 113.9, 109.9, 97.7, 49.5.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-6-bromo-1H-indol-3-yl)-2-cyanoacrylic acid (JXL062)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.72 (s, 1H), 8.47 (s, 1H), 8.06 (s, 2H), 8.03 (s, 1H), 7.94 (d, *J* = 8.2 Hz, 1H), 7.39 (d, *J* = 7.9 Hz, 1H), 5.81 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.7, 145.6, 140.2, 137.4, 135.3, 131.1 (q, *J*_{c-f} = 32.8 Hz), 129.1, 127.0, 125.9, 123.6 (q, *J*_{c-f} = 274.0 Hz), 122.5, 121.5, 118.2, 117.2, 114.8, 110.3, 96.9, 49.3.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-6-chloro-1H-indol-3-yl)-2-cyanoacrylic acid (JXL063)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.75 (s, 1H), 8.47 (s, 1H), 8.05 (m, 3H), 8.00 (d, *J* = 7.4 Hz, 1H), 7.89 (s, 1H), 7.28 (d, *J* = 6.5 Hz, 1H), 5.81 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.7, 145.6, 140.2, 137.0, 135.4, 131.1 (q, *J*_{c-f} = 32.8 Hz), 129.2, 129.0, 126.7, 123.6 (q, *J*_{c-f} = 274.0 Hz), 123.3, 122.5, 121.2, 118.2, 111.9, 110.3, 97.0, 49.3.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-7-chloro-1H-indol-3-yl)-2-cyanoacrylic acid (JXL064)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.69 (s, 1H), 8.50 (s, 1H), 8.03 (s, 1H), 7.97 (d, *J* = 7.5 Hz, 1H), 7.73 (s, 2H), 7.32 (d, *J* = 7.1 Hz, 1H), 7.25 (m, 1H), 6.14 (s, 2H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.5, 145.1, 142.1, 136.9, 131.5, 131.2 (q, *J*_{c-f} = 32.8 Hz), 127.6, 126.1, 124.1, 123.6 (q, *J*_{c-f} = 274.0 Hz), 122.5, 122.0, 118.8, 118.0, 116.9, 110.3, 97.9, 51.8.




JXL065

Ethyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-4-bromo-1*H*-indol-3-yl)-2-cyanoacrylate (JXL065)

¹H NMR (500 MHz, DMSO-*d*₆) δ 9.67 (s, 1H), 8.71 (s, 1H), 7.86 (s, 1H), 7.56 (s, 2H), 7.52 (d, *J* = 7.4 Hz, 1H), 7.17 (m, 2H), 5.55 (s, 2H), 4.37 (q, *J* = 7.1 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 163.4, 146.9, 137.6, 137.4, 134.3, 132.7 (q, *J*_{C-F} = 33.9 Hz), 128.2, 126.7, 125.6, 125.1, 122.8 (q, *J*_{C-F} = 273.4 Hz), 122.7, 118.0, 114.9, 111.8, 109.9, 96.3, 62.2, 50.6, 14.3.




JXL066

Ethyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-4-fluoro-1*H*-indol-3-yl)-2-cyanoacrylate (JXL066)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.86 (s, 1H), 8.61 (s, 1H), 7.86 (s, 1H), 7.56 (s, 2H), 7.23 (m, 1H), 7.01 (m, 2H), 5.54 (s, 2H), 4.37 (q, *J* = 7.1 Hz, 2H), 1.39 (t, *J* = 7.1 Hz, 3H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 163.1, 157.4 (d, *J* = 243.9 Hz), 147.4, 138.2 (d, *J* = 10.1 Hz), 137.8, 133.2, 132.7 (q, *J*_{C-F} = 33.9 Hz), 126.7, 125.3, 122.8 (q, *J*_{C-F} = 273.4 Hz), 122.7, 117.9, 116.9 (d, *J*_{C-F} = 17.6 Hz), 110.0, 109.0 (d, *J*_{C-F} = 19.5 Hz), 106.6, 97.1, 62.2, 50.7, 14.3.



***tert*-Butyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylate (JXL068)**

¹H NMR (500 MHz, CDCl₃) δ 8.57 (s, 1H), 8.54 (s, 1H), 7.86 (d, *J* = 7.6 Hz, 1H), 7.84 (s, 1H), 7.56 (s, 2H), 7.33 (m, 2H), 7.22 (d, *J* = 7.4 Hz, 1H), 5.55 (s, 2H), 1.59 (s, 9H).


¹³C NMR (126 MHz, CDCl₃) δ 162.4, 144.7, 138.2, 135.9, 133.0, 132.7 (q, *J*_{C-F} = 32.8 Hz), 128.6, 126.7, 124.5, 123.0, 122.9 (q, *J*_{C-F} = 277.2 Hz), 122.6, 119.0, 118.2, 111.1, 110.4, 97.7, 82.9, 50.4, 28.1.



(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-2-cyanoacrylic acid (JXL069)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.85 (s, 1H), 8.47 (m, 3H), 8.09 (s, 2H), 8.04 (s, 1H), 7.35 (dd, *J* = 7.1, 4.6 Hz, 1H), 5.84 (s, 2H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 185.9, 164.6, 147.8, 146.1, 145.5, 141.0, 135.2, 131.0 (q, *J*_{C-F} = 32.8 Hz), 129.4, 126.9, 123.6 (q, *J*_{C-F} = 274.0 Hz), 122.3, 120.1, 119.1, 118.2, 97.1, 47.8.




JXL072

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-pyrrolo[3,2-b]pyridin-3-yl)-2-cyanoacrylic acid (JXL072)

¹H NMR (500 MHz, DMSO-d₆) δ 8.94 (s, 1H), 8.68 (s, 1H), 8.54 (s, 1H), 8.11 (m, 4H), 7.35 (s, 1H), 5.87 (s, 2H).

¹³C NMR (126 MHz, DMSO-d₆) δ 164.5, 145.9, 144.2, 140.1, 135.9, 131.1 (q, *J*_{C-F} = 32.8 Hz), 129.6, 129.2, 124.7, 123.6 (q, *J*_{C-F} = 274.0 Hz), 122.5, 120.0, 119.5, 117.9, 110.1, 97.2, 49.8.




JXL073

(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-pyrrolo[2,3-c]pyridin-3-yl)-2-cyanoacrylic acid (JXL073)

¹H NMR (500 MHz, DMSO-d₆) δ 9.14 (s, 1H), 8.96 (s, 1H), 8.55 (s, 1H), 8.40 (d, *J* = 5.1 Hz, 1H), 8.19 (s, 2H), 8.11 (app. s, 2H), 5.95 (s, 2H).


¹³C NMR (126 MHz, DMSO-d₆) δ 164.5, 145.5, 140.7, 139.9, 138.1, 134.5, 133.6, 131.1 (q, *J*_{C-F} = 32.8 Hz), 129.4, 127.2, 123.6 (q, *J*_{C-F} = 274.0 Hz), 122.7, 118.0, 114.6, 109.7, 97.9, 49.8.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-chloro-1H-pyrrolo[2,3-b]pyridin-3-yl)-2-cyanoacrylic acid (JXL076)

¹H NMR (500 MHz, DMSO-*d*₆) δ 9.00 (s, 1H), 8.98 (s, 1H), 8.36 (s, 1H), 8.12 (s, 2H), 8.04 (br. s, 1H), 7.47 (br. s, 1H), 5.85 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.4, 148.7, 145.8, 145.7, 140.2, 136.1, 135.3, 131.0 (q, *J*_{C-F} = 32.8 Hz), 129.5, 123.6 (q, *J*_{C-F} = 274.0 Hz), 122.4, 120.4, 117.7, 116.8, 108.2, 98.1, 48.2.



(E)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-bromo-1H-pyrrolo[2,3-b]pyridin-3-yl)-2-cyanoacrylic acid (JXL077)

¹H NMR (500 MHz, DMSO-*d*₆) δ 9.17 (s, 1H), 9.02 (s, 1H), 8.27 (d, *J* = 5.0 Hz, 1H), 8.12 (s, 2H), 8.05 (s, 1H), 7.63 (d, *J* = 5.0 Hz, 1H), 5.85 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 164.4, 148.2, 145.5, 145.3, 140.2, 135.4, 131.0 (q, *J*_{C-F} = 32.8 Hz), 129.6, 124.9, 123.7, 123.6 (q, *J*_{C-F} = 274.0 Hz), 122.4, 118.4, 117.7, 108.6, 97.9, 48.2.



Methyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylate (JXL078)

¹H NMR (500 MHz, CDCl₃) δ 8.62 (s, 1H), 8.61 (s, 1H), 7.89 (d, *J* = 7.7 Hz, 1H), 7.84 (s, 1H), 7.57 (s, 2H), 7.35 (m, 2H), 7.24 (m, 1H), 5.55 (s, 2H), 3.92 (s, 3H).


¹³C NMR (126 MHz, CDCl₃) δ 164.5, 145.8, 138.1, 136.0, 133.5, 132.7 (q, *J*_{C-F} = 32.8 Hz), 128.6, 126.8, 124.7, 123.9, 123.6 (q, *J*_{C-F} = 274.0 Hz), 123.2, 119.1, 118.0, 111.2, 110.5, 95.4, 53.0, 50.5.



(*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-(morpholine-4-carbonyl)acrylonitrile (JXL081)

¹H NMR (500 MHz, CDCl₃) δ 8.48 (s, 1H), 8.34 (s, 1H), 7.84 (s, 2H), 7.56 (app. s, 2H), 7.32 (m, 2H), 7.22 (m, 1H), 5.54 (s, 2H), 3.77 (br. s, 8H).


¹³C NMR (126 MHz, CDCl₃) δ 164.0, 145.2, 138.4, 135.8, 132.7 (q, *J*_{C-F} = 34.0 Hz), 131.8, 128.4, 126.8, 125.0, 124.5, 123.1 (q, *J*_{C-F} = 273.3 Hz), 122.8, 121.8, 119.0, 118.6, 111.4, 110.3, 98.1, 66.7, 50.3, 50.0.



Ethyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-2-cyanoacrylate (JXL082)

¹H NMR (500 MHz, CDCl₃) δ 8.64 (s, 1H), 8.51 (s, 1H), 8.48 (dd, *J* = 4.6, 1.2 Hz, 1H), 8.21 (dd, *J* = 7.9, 1.2 Hz, 1H), 7.83 (s, 1H), 7.77 (s, 2H), 7.34 (dd, *J* = 7.9, 4.6 Hz, 1H), 5.69 (s, 2H), 4.38 (q, *J* = 7.1 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H).


¹³C NMR (126 MHz, CDCl₃) δ 163.3, 147.6, 145.7, 145.0, 138.5, 132.7, 132.3 (q, *J*_{C-F} = 33.6 Hz), 127.9, 127.7, 123.1 (q, *J*_{C-F} = 273.3 Hz), 122.5, 120.3, 119.0, 117.6, 109.4, 97.0, 62.3, 48.3, 14.3.



Ethyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-4-chloro-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-2-cyanoacrylate (JXL087)

¹H NMR (500 MHz, CDCl₃) δ 9.22 (s, 1H), 8.77 (s, 1H), 8.32 (d, *J* = 5.2 Hz, 1H), 7.83 (s, 1H), 7.77 (s, 2H), 7.31 (d, *J* = 5.2 Hz, 1H), 5.67 (s, 2H), 4.37 (q, *J* = 7.1 Hz, 2H), 1.39 (t, *J* = 7.1 Hz, 3H).


¹³C NMR (126 MHz, CDCl₃) δ 163.0, 148.5, 146.2, 145.4, 138.1, 137.4, 133.1, 132.4 (q, *J*_{C-F} = 33.7 Hz), 128.0, 122.9 (q, *J*_{C-F} = 273.4 Hz), 122.6, 121.8, 117.6, 117.2, 109.7, 97.8, 62.4, 48.6, 14.3.



Ethyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-4-bromo-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-2-cyanoacrylate (JXL088)

¹H NMR (500 MHz, CDCl₃) δ 9.40 (s, 1H), 8.78 (s, 1H), 8.22 (d, *J* = 5.1 Hz, 1H), 7.83 (s, 1H), 7.77 (s, 2H), 7.49 (d, *J* = 5.1 Hz, 1H), 5.67 (s, 2H), 4.37 (q, *J* = 7.1 Hz, 2H), 1.39 (t, *J* = 7.1 Hz, 3H).


¹³C NMR (126 MHz, CDCl₃) δ 163.1, 148.1, 145.7, 145.1, 138.1, 137.4, 133.4, 132.4 (q, *J*_{c-f} = 33.7 Hz), 123.7, 122.9 (q, *J*_{c-f} = 273.4 Hz), 122.6, 119.7, 117.6, 110.0, 101.4, 97.4, 62.4, 48.6, 14.3.



Ethyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-5-chloro-1*H*-indol-3-yl)-2-cyanoacrylate (JXL089)

¹H NMR (500 MHz, CDCl₃) δ 8.60 (s, 1H), 8.50 (s, 1H), 7.85 (s, 1H), 7.83 (s, 1H), 7.55 (s, 2H), 7.28 (dd, *J* = 8.7, 1.7 Hz, 1H), 7.14 (d, *J* = 8.7 Hz, 1H), 5.54 (s, 2H), 4.38 (q, *J* = 7.1 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H).


¹³C NMR (126 MHz, CDCl₃) δ 163.2, 144.8, 137.7, 134.3, 134.1, 132.8 (q, *J*_{c-f} = 33.7 Hz), 129.7, 129.3, 126.7, 125.1, 122.8 (q, *J*_{c-f} = 273.4 Hz), 122.7, 118.9, 117.7, 111.6, 110.6, 96.9, 62.3, 50.6, 14.3.



Ethyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-4-cyano-1*H*-indol-3-yl)-2-cyanoacrylate (JXL090)

¹H NMR (500 MHz, CDCl₃) δ 9.29 (s, 1H), 8.78 (s, 1H), 7.87 (s, 1H), 7.68 (dd, *J* = 7.4, 0.8 Hz, 1H), 7.56 (s, 2H), 7.49 (dd, *J* = 8.4, 0.8 Hz, 1H), 7.38 (dd, *J* = 8.4, 7.4 Hz, 1H), 5.62 (s, 2H), 4.38 (q, *J* = 7.1 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H).


¹³C NMR (126 MHz, CDCl₃) δ 162.6, 144.3, 137.3, 136.2, 135.1, 132.9 (q, *J*_{c-f} = 33.7 Hz), 129.4, 127.3, 126.7, 124.1, 123.0, 122.8 (q, *J*_{c-f} = 273.4 Hz), 117.9, 117.5, 115.4, 110.7, 103.5, 98.6, 62.4, 50.6, 14.3.



Methyl (*E*)-1-(3,5-bis(trifluoromethyl)benzyl)-3-(2-cyano-3-ethoxy-3-oxoprop-1-en-1-yl)-1*H*-indole-4-carboxylate (JXL091)

¹H NMR (500 MHz, CDCl₃) δ 9.36 (s, 1H), 8.71 (s, 1H), 7.93 (dd, 1H, *J* = 7.4, 1.1 Hz, 1H), 7.85 (s, 1H), 7.55 (s, 2H), 7.40 (dd, *J* = 8.3, 1.1 Hz, 1H), 7.34 (dd, *J* = 8.3, 7.4 Hz, 1H), 5.58 (s, 2H), 4.37 (q, *J* = 7.1 Hz, 2H), 4.04 (s, 3H), 1.40 (t, *J* = 7.1 Hz, 3H).

¹³C NMR (126 MHz, CDCl₃) δ 167.8, 163.5, 150.1, 137.8, 137.2, 134.9, 132.8 (q, *J*_{c-f} = 33.7 Hz), 126.7, 125.8, 125.1, 123.6, 122.8 (q, *J*_{c-f} = 273.4 Hz), 122.7, 121.7, 118.0, 114.6, 111.3, 96.4, 62.1, 52.6, 50.5, 14.3.

Experimental detail for the synthesis of JXL024


To the solution of 1-phenyl-1*H*-indole-3-carbaldehyde (0.4 mmol, 90 mg) in AcOH (3 mL) were added thiazolidine-2,4-dione (1 equiv, 0.4 mmol, 46.8 mg) and NaOAc (3 equiv, 98 mg). The reaction mixture was stirred at reflux for 24 hours. After it was cooled to 21 °C, the reaction mixture was filtered by vacuum filtration and washed by AcOH (3 mL \times 3) and water (5 mL \times 3). After drying by vacuum, the desired product was produced. yield: 34%, 44 mg.

(Z)-5-((1-Phenyl-1*H*-indol-3-yl)methylene)thiazolidine-2,4-dione (JXL024)

¹H NMR (500 MHz DMSO-*d*₆) δ 7.98 (m, 2H), 7.79 (s, 1H), 7.66 (app. d, *J* = 7.7 Hz, 2H), 7.62 (app. t, *J* = 7.7 Hz, 2H), 7.54 (d, *J* = 8.0 Hz, 1H), 7.49 (t, *J* = 7.1 Hz, 1H), 7.30 (m, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 172.5, 169.5, 138.4, 136.2, 130.5, 129.9, 128.3, 128.2, 125.0, 124.6, 122.4, 121.5, 121.4, 119.6, 113.0, 111.5.

The following compounds were synthesized by a route similar to that described for JXL024.: JXL067, JXL070, JXL072, JXL074, JXL075.

**(Z)-2-Imino-5-((1-phenyl-1*H*-indol-3-yl)methylene)thiazolidin-4-one (JXL023)**

¹H NMR (500 MHz, DMSO-*d*₆) δ 11.94 (br. s, 1H), 9.33 (br. s, 1H), 8.97 (s, 1H), 7.64 (m, 10H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 180.9, 174.8, 172.5, 138.6, 136.2, 130.5, 129.3, 128.1, 126.3, 124.8, 124.5, 122.2, 120.3, 119.7, 113.5, 111.5.



(Z)-5-((1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)methylene)-2-iminothiazolidin-4-one (JXL067)

¹H NMR (500 MHz, DMSO-*d*₆) δ 9.21 (s, 1H), 9.01 (s, 1H), 8.03 (s, 1H), 7.94 (br. s, 3H), 7.86 (d, *J* = 7.7 Hz, 1H), 7.81 (s, 1H), 7.58 (d, *J* = 7.9 Hz, 1H), 7.24 (m, 1H), 7.19 (m, 1H), 5.76 (s, 2H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 180.0, 174.9, 141.4, 136.5, 131.0 (q, *J*_{C-F} = 32.8 Hz), 130.4, 128.5, 127.9, 125.1, 123.8, 123.6 (q, *J*_{C-F} = 274.0 Hz), 122.1, 121.7, 120.8, 119.3, 111.8, 111.3, 48.9.



(Z)-5-((4-Fluoro-1-phenyl-1H-indol-3-yl)methylene)thiazolidine-2,4-dione (JXL070)

¹H NMR (500 MHz, DMSO-*d*₆) δ 12.45 (br. s, 1H), 8.11 (s, 1H), 7.83 (s, 1H), 7.65 (m, 4H), 7.52 (s, 1H), 7.32 (m, 2H), 7.12 (m, 1H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 168.0, 167.7, 156.9 (d, *J*_{C-F} = 245.7 Hz), 138.7 (d, *J*_{C-F} = 10.1 Hz), 137.9, 130.8, 130.5, 128.8, 125.5, 125.4, 124.3, 120.0, 116.2 (d, *J*_{C-F} = 18.9 Hz), 110.7, 108.5, 108.1 (d, *J*_{C-F} = 18.9 Hz).



(Z)-5-((6-Fluoro-1-phenyl-1H-indol-3-yl)methylene)thiazolidine-2,4-dione (JXL071)

¹H NMR (500 MHz, DMSO-*d*₆) δ 12.42 (s, 1H), 8.06 (app. s, 2H), 7.85 (s, 1H), 7.67 (m, 2H), 7.62 (m, 2H), 7.50 (m, 1H), 7.31 (d, *J* = 9.6 Hz, 1H), 7.16 (t, *J* = 8.5 Hz, 1H).


¹³C NMR (126 MHz, DMSO-*d*₆) δ 168.0, 167.7, 160.6 (d, *J*_{C-F} = 245.7 Hz), 138.0, 136.3 (d, *J*_{C-F} = 12.6 Hz), 131.0, 130.6, 128.5, 125.0, 124.8, 123.5, 121.3 (d, *J*_{C-F} = 10.0 Hz), 120.0, 112.6, 111.1 (d, *J*_{C-F} = 18.9 Hz), 98.2 (d, *J*_{C-F} = 18.9 Hz).



(Z)-5-Benzylidenethiazolidine-2,4-dione (JXL074)

¹H NMR (500 MHz, DMSO-*d*₆) δ 12.60 (br. s, 1H), 7.77 (s, 1H), 7.58 (app. d, *J* = 7.3 Hz, 2H), 7.51 (app. t, *J* = 7.4 Hz, 2H), 7.46 (m, 1H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 168.4, 167.8, 133.5, 132.3, 130.9, 130.5, 129.8, 124.0.




(Z)-5-((1H-Indol-3-yl)methylene)thiazolidine-2,4-dione (JXL075)

¹H NMR (500 MHz, DMSO-*d*₆) δ 12.28 (s, 1H), 12.11 (s, 1H), 8.03 (s, 1H), 7.87 (d, *J* = 7.3 Hz, 1H), 7.72 (s, 1H), 7.48 (d, *J* = 7.6 Hz, 1H), 7.23 (m, 1H), 7.18 (m, 1H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 168.2, 167.8, 136.7, 129.1, 127.3, 125.0, 123.5, 121.5, 118.8, 116.7, 112.9, 110.9.

Experimental detail for the synthesis of JXL022



To the solution of 4-pyridinecarboxaldehyde (1 mmol, 107 mg) in ethanol (1 mL) were added ethyl 2-cyanoacetate (1.3 equiv, 1.3 mmol, 140 μL) and L-proline (40 mol%, 0.4 mmol, 58 mg). The reaction was stirred at 21 °C for 12 h and yellow solid precipitated gradually. After completion of the reaction, ice-cold water (2 mL) was added into the reaction vial. The solid

was separated by Buchner funnel filtration and washed with water (2 mL \times 3) and dried to afford the desired product, ethyl (*E*)-2-cyano-3-(pyridin-4-yl)acrylate, which was used for the next step without further purification.


To the solution of (*E*)-2-cyano-3-(pyridin-4-yl)acrylate (0.21 mmol, 42.4 mg) in THF (2 mL) was added 0.5N LiOH solution (3 equiv, 0.4 mmol, 0.8 mL). The reaction mixture was stirred at 21 °C for 1 h. After reaction completion shown by TLC, THF was evaporated. Concentrated HCl was added dropwise to acidify the reaction mixture until pH was lower than 1, meanwhile yellow solid precipitated. Ice-cold water (5 mL) was added to the reaction mixture and the solid was separated by Buchner funnel filtration and washed with water (5 mL \times 3). After dried by vacuum, the solid was washed by 2 mL of solvent mixture (hexanes/EtOAc = 5:1) 5 to 10 times and monitored by TLC until non-polar impurities disappear. Finally, the purity of the product was checked by NMR. yield: 64%, 23.4 mg.

Ethyl (*E*)-2-cyano-3-(pyridin-4-yl)acrylate (JXL022)

¹H NMR (500 MHz, CDCl₃) δ 8.81 (d, J = 5.2 Hz, 2H), 8.18 (s, 1H), 7.74 (d, J = 5.2 Hz, 2H), 4.41 (q, J = 7.1 Hz, 2H), 1.41 (t, J = 7.1 Hz, 3H).

¹³C NMR (126 MHz, CDCl₃) δ 161.2, 152.0, 151.0, 137.9, 123.2, 114.2, 108.2, 63.2, 14.0.


The following compounds were synthesized by a route similar to that described for JXL022: JXL030, JXL031, JXL032, JXL033, JXL034, JXL042, JXL43, JXL044, JXL045, JXL046, JXL047, JXL048, JXL049.



(*E*)-2-Cyano-3-(2-fluorophenyl)acrylic acid (JXL030)

¹H NMR (500 MHz, DMSO-*d*₆) δ 8.49 (s, 1H), 8.31 (t, J = 7.4 Hz, 1H), 7.63 (m, 1H), 7.36 (t, J = 7.4 Hz, 1H), 7.29 (m, 1H).

¹³C NMR (126 MHz, DMSO-*d*₆) δ 162.9, 161.5 (d, J_{C-F} = 256.2 Hz), 145.4 (d, J_{C-F} = 7.8 Hz), 135.0 (d, J_{C-F} = 9.2 Hz), 128.7, 124.7, 119.8 (d, J_{C-F} = 10.9 Hz), 115.8 (d, J_{C-F} = 21.9 Hz), 114.9, 105.9.




JXL032

(E)-2-Cyano-3-(4-fluorophenyl)acrylic acid (JXL032)

¹H NMR (500 MHz, DMSO-d₆) δ 8.30 (s, 1H), 8.10 (m, 2H), 7.29 (m, 2H).


¹³C NMR (126 MHz, DMSO-d₆) δ 165.2 (d, *J*_{C-F} = 255.2 Hz), 163.5, 153.1, 133.3 (d, *J*_{C-F} = 9.3 Hz), 128.3, 116.0 (d, *J*_{C-F} = 22.4 Hz), 115.3, 103.2.



JXL033

(E)-2-cyano-3-(3-(trifluoromethyl)phenyl)acrylic acid (JXL033)

¹H NMR (500 MHz, CD₃OD) δ 8.24 (s, 1H), 8.17 (m, 2H), 7.80 (d, *J* = 7.8 Hz, 1H), 7.70 (app. t, *J* = 7.8 Hz, 1H).




JXL034

(E)-2-Cyano-3-(4-(trifluoromethyl)phenyl)acrylic acid (JXL034)

¹H NMR (500 MHz, DMSO-d₆) δ 8.39 (s, 1H), 8.17 (d, *J* = 7.7 Hz, 2H), 7.84 (d, *J* = 7.7 Hz, 2H).

¹³C NMR (126 MHz, DMSO-d₆) δ 162.8, 152.6, 135.4, 133.1 (q, *J*_{C-F} = 32.9 Hz), 131.0, 125.7, 123.7 (q, *J*_{C-F} = 272.2 Hz), 114.8, 106.7.




JXL042

(E)-2-Cyano-3-(3-fluoro-4-methylphenyl)acrylic acid (JXL042)

¹H NMR (500 MHz, DMSO-d₆) δ 8.28 (s, 1H), 7.78 (m, 2H), 7.49 (t, *J* = 8.0 Hz, 1H), 2.30 (s, 3H).


¹³C NMR (126 MHz, DMSO-d₆) δ 163.6, 160.9 (d, *J*_{C-F} = 244.6 Hz), 153.5, 133.0, 131.6 (d, *J*_{C-F} = 7.5 Hz), 130.8 (d, *J*_{C-F} = 17.6 Hz), 127.2, 117.9 (d, *J*_{C-F} = 23.9 Hz), 116.4, 104.5, 15.0.



(E)-2-Cyano-3-(3,4-difluorophenyl)acrylic acid (JXL043)

¹H NMR (500 MHz, DMSO-d₆) δ 8.32 (s, 1H), 8.09 (d, *J* = 8.0 Hz, 1H), 7.94 (br. s, 1H), 7.67 (d, *J* = 8.8 Hz, 1H).


¹³C NMR (126 MHz, DMSO-d₆) δ 163.3, 152.3, 152.2 (dd, *J*_{c-f} = 255.4, 12.6 Hz), 149.9 (dd, *J*_{c-f} = 248.2, 12.6 Hz), 129.8, 128.8, 120.1 (d, *J*_{c-f} = 17.6 Hz), 119.1 (d, *J*_{c-f} = 17.6 Hz), 116.3, 105.8.



(E)-2-Cyano-3-(2,4-difluorophenyl)acrylic acid (JXL044)


¹H NMR (500 MHz, DMSO-d₆) δ 14.2 (br. s, 1H), 8.27 (s, 1H), 8.24 (m, 1H), 7.52 (m, 1H), 7.35 (dt, *J* = 8.6, 2.2 Hz, 1H).

¹³C NMR (126 MHz, DMSO-d₆) δ 165.5 (dd, *J*_{c-f} = 255.8, 12.6 Hz), 163.1, 161.9 (dd, *J*_{c-f} = 270.5, 12.6 Hz), 145.1, 131.1, 117.0, 115.9, 113.5 (d, *J*_{c-f} = 22.7 Hz), 106.8, 105.6 (t, *J*_{c-f} = 26.5 Hz).



(E)-3-(3,5-bis(trifluoromethyl)phenyl)-2-cyanoacrylic acid (JXL045)


¹H NMR (500 MHz, CD₃OD) δ 8.59 (s, 2H), 8.48 (s, 1H), 8.19 (s, 1H).



(E)-3-(2-Chloro-3-(trifluoromethyl)phenyl)-2-cyanoacrylic acid (JXL046)

¹H NMR (500 MHz, DMSO-d₆) δ 8.50 (s, 1H), 8.23 (d, *J* = 7.7 Hz, 1H), 8.05 (d, *J* = 7.4 Hz, 1H), 7.75 (app. t, *J* = 7.9 Hz, 1H).


^{13}C NMR (126 MHz, DMSO- d_6) δ 162.6, 150.5, 134.4, 133.5, 132.1, 131.2, 128.9, 128.3 (q, $J_{\text{c-f}}$ = 30.2 Hz), 123.1 (q, $J_{\text{c-f}}$ = 274.1 Hz) 115.1, 111.1.



(E)-2-Cyano-3-(4-fluoro-3-(trifluoromethyl)phenyl)acrylic acid (JXL047)

^1H NMR (500 MHz, DMSO- d_6) δ 8.46 (dd, J = 7.1, 1.8 Hz, 1H), 8.44 (s, 1H), 8.40 (m, 1H), 7.75 (m, 1H).


^{13}C NMR (126 MHz, DMSO- d_6) δ 163.3, 161.1 (d, $J_{\text{c-f}}$ = 262.1 Hz), 152.2, 137.6 (d, $J_{\text{c-f}}$ = 10.1 Hz), 130.5, 129.3, 122.6 (q, $J_{\text{c-f}}$ = 272.8 Hz), 118.9 (d, $J_{\text{c-f}}$ = 21.2 Hz), 118.2 (qd, $J_{\text{c-f}}$ = 32.9, 12.6 Hz), 116.2, 106.3.



(E)-2-Cyano-3-phenylacrylic acid (JXL048)

^1H NMR (500 MHz, DMSO- d_6) δ 8.33 (s, 1H), 8.02 (m, 2H), 7.59 (m, 3H).

^{13}C NMR (126 MHz, DMSO- d_6) δ 163.7, 154.9, 133.6, 132.0, 131.1, 129.8, 116.5, 104.3.




(E)-2-cyano-3-(4-hydroxyphenyl)acrylic acid (JXL049)

^1H NMR (500 MHz, DMSO- d_6) δ 8.16 (s, 1H), 7.95 (d, J = 8.8 Hz, 2H), 6.92 (d, J = 8.8 Hz, 2H).

^{13}C NMR (126 MHz, DMSO- d_6) δ 164.2, 163.0, 153.7, 133.6, 122.8, 117.2, 116.3, 99.2.

Experimental detail for the synthesis of JXL079



A flask containing Mg powder (10 mmol, 240 mg) and a stir bar was sealed and vacuumed and refilled with argon three times. Anhydrous diethyl ether (32 mL) and bis(trifluoromethyl)benzyl bromide (8 mmol, 1.46 mL) was added to the reaction flask. The reaction mixture was stirred and refluxed for 30 min and then ground dry ice powder (5 g) was added into the reaction flask. After 1 h, the reaction was complete as shown by TLC. The extra Mg powder was filtered off and the solvent was evaporated under vacuum. 1N HCl (20 mL) was added to the residue and the precipitate was filtered and dried to provide the desired carboxylic acid.

A flask containing a stir bar was sealed, vacuumed and refilled with argon three times. Anhydrous dichloromethane (20 mL) and DIBAL (1 M in hexanes, 6 mmol, 6 mL) were added to the flask. The crude carboxylic acid (2 mmol, 544 mg) dissolved in dry dichloromethane (10 mL) was added to the reaction flask at -78 °C. After 2 h, the reaction was complete as shown by TLC and it was then quenched by adding sat. ammonium chloride (10 mL). The resulting mixture was extracted with dichloromethane (20 mL×3) and the organic phases were combined and evaporated on the rotavap. The residue was purified by flash column chromatography (hexanes:ethyl acetate = 10:1) to afford the desired product **2-(3,5-bis(trifluoromethyl)phenyl)ethan-1-ol** (yield: 90%, 464 mg).

To a solution of 2-(3,5-bis(trifluoromethyl)phenyl)ethan-1-ol (0.2 mmol, 51.6 mg) in dichloro-methane (2 mL) was added triethylamine (0.22 mmol, 31 µL) and mesyl chloride (MsCl, 0.2 mmol, 17 µL) at 0 °C. After stirring for 1 h, the reaction was complete as shown by TLC. The solvent was removed by flowing air over the open flask. The residue was purified

by flash column chromatography (hexanes:ethyl acetate = 10:1) to afford the desired product **3,5-bis(trifluoromethyl)phenethyl methanesulfonate** (yield: 82%, 55.1 mg).

A flask containing NaH (60%, 0.22 mmol, 8.8 mg) and a stir bar was sealed, vacuumed and refilled with argon three times. Anhydrous THF (3 mL) and a solution of *tert*-butyl (*E*)-2-cyano-3-(1*H*-indol-3-yl)acrylate (0.2 mmol, 53.6 mg) in THF (2 mL) were added into the reaction flask. The reaction mixture was stirred for 30 min and then 3,5-bis(trifluoromethyl)phenethyl methanesulfonate (0.164 mmol, 55.1 mg) in 2 mL THF was added. The reaction was stirred for 24 h and quenched by sat. NH₄Cl solution. The resulting mixture was extracted with dichloromethane (4 mL×3) and the organic phases were combined and evaporated on the rotavap. The residue was purified by flash column chromatography (hexanes:ethyl acetate = 10:1) to afford the desired product ***tert*-butyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)phenethyl)-1*H*-indol-3-yl)-2-cyano-acrylate** (yield: 68%, 69 mg).


To a solution of methyl *tert*-butyl (*E*)-3-(1-(3,5-bis(trifluoromethyl)phenethyl)-1*H*-indol-3-yl)-2-cyanoacrylate (0.1 mmol, 50.8 mg) in dichloromethane (2 mL) was added trifluoroacetic acid (3 equiv, 0.3 mmol, 34 μ L). The reaction mixture was stirred at 21 °C for 30 min and a yellow solid precipitated. After the reaction was complete as shown by TLC, the reaction solvent was evaporated by flowing air over the open flask. The solid was washed by 2 mL of solvent mixture (hexanes/EtOAc = 5:1) 5 to 10 times and monitored by TLC until all the non-polar impurities disappeared. Finally, the purity of the product was checked by NMR. yield: 87%, 39 mg.

(*E*)-3-(1-(3,5-Bis(trifluoromethyl)phenethyl)-1*H*-indol-3-yl)-2-cyanoacrylic acid (JXL079)

¹H NMR (500 MHz, CD₃OD) δ 8.51 (s, 1H), 8.14 (s, 1H), 7.83 (d, *J* = 7.3 Hz, 1H), 7.74 (s, 1H), 7.52 (m, 3H), 7.31 (m, 2H), 4.64 (t, *J* = 6.4 Hz, 2H), 3.35 (t, *J* = 6.3 Hz, 2H).

¹³C NMR (126 MHz, CD₃OD) δ 165.3, 145.3, 141.1, 136.1, 133.6, 131.3 (q, *J*_{C-F} = 32.8 Hz), 129.3, 128.3, 123.7, 123.3 (q, *J*_{C-F} = 272.5 Hz), 122.3, 120.3, 118.1, 117.6, 110.6, 109.6, 94.1, 47.8, 35.0.

Experimental detail for the synthesis of JXL080




To a solution of ethyl (*E*)-2-cyano-3-(1*H*-indol-3-yl)acrylate (0.5 mmol, 112 mg) in dichloromethane (5 mL) were added 2-(3,5-bis(trifluoromethyl)phenyl)acetic acid (0.55 mmol, 150 mg), DMAP (catalytic amount, 6 mg), and DCC (0.5 mmol, 103 mg) at 0 °C. The mixture was allowed to reach 21 °C and stirred overnight. The white precipitate was filtered, and the resulting solution was concentrated in vacuum. The solid was purified by flash column chromatography (hexanes:ethyl acetate = 10:1) to afford the desired product (yield: 78%, 192.6 mg).

Ethyl (*E*)-3-(1-(2-(3,5-bis(trifluoromethyl)phenyl)acetyl)-1*H*-indol-3-yl)-2-cyanoacrylate (JXL080)

¹H NMR (500 MHz, CDCl₃) δ 8.90 (s, 1H), 8.52 (s, 1H), 8.49 (d, *J* = 8.1 Hz, 1H), 7.88 (s, 1H), 7.86 (s, 2H), 7.78 (d, *J* = 7.5 Hz, 1H), 7.47 (m, 2H), 4.50 (s, 2H), 4.41 (q, *J* = 7.1 Hz, 2H), 1.43 (t, *J* = 7.1 Hz, 3H).

¹³C NMR (126 MHz, CDCl₃) δ 168.0, 162.2, 144.2, 135.6, 134.4, 132.3 (q, *J*_{C-F} = 33.6 Hz), 130.1, 129.8, 128.8, 127.8, 127.3, 125.5, 123.1 (q, *J*_{C-F} = 273.3 Hz), 118.3, 117.1, 117.0, 115.9, 101.8, 62.8, 41.9, 14.3.

Experimental detail for the synthesis of JXL083



To a solution of 1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indole-3-carbaldehyde (10 mmol, 3.71 g) in acetone (60 mL) was added 2-methyl-2-butene (9 mL), NaH₂PO₄ (3 equiv, 4.4 g) and NaClO₂ (6.6 mmol, 6 g) in 6 mL water. The reaction mixture was stirred at 21 °C for 24 h. After the reaction was complete as shown by TLC, the reaction solvent was evaporated on the rotavap. The crude material was dissolved in ethyl acetate (30 mL) and water (30 mL) and extracted with ethyl acetate (30 mL × 3). The organic phase was combined, dried with sodium sulfate and evaporated on the rotavap. The solid was purified by flash column chromatography (hexanes:ethyl acetate = 2:1) to afford the desired product **1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indole-3-carboxylic acid** (yield: 89%, 3.44 g).


A 100 mL round bottom flask with a stir bar containing the carboxylic acid (5 mmol, 1935 mg) from the previous step was sealed, vacuumed and refilled with argon three times. To the flask was added 50 mL dichloromethane and oxalyl chloride (25 mmol, 2.1 mL) dropwise. The reaction mixture was stirred at 21 °C for 1.5 h. The reaction solvent was evaporated by vacuum and the resulting compound was used for the next step.

A 100 mL round bottom flask with a stir bar was sealed, vacuumed and refilled with argon three times. Diisopropylamine (5.5 mmol, 765 µL) and THF (10 mL) were added into the flask and it was cooled to -78 °C. nBuLi (2.5 M in hexanes, 5 mmol, 2 mL) was added slowly to the flask. After the mixture had stirred for 30 min, a solution of ethyl 2-cyanoacetate (5 mmol, 590 µL) in THF (10 mL) was added slowly to the flask. After the mixture had stirred for 1 h, a solution of the acyl chloride (5 mmol from previous step) in THF (5 mL) was added slowly to the reaction mixture. After 1 h, the reaction was quenched by adding aqueous 1M HCl solution (10 mL) and extracted with ethyl acetate (10 mL × 3). The organic phase was combined, dried with sodium sulfate and evaporated by rotavap. The solid was purified by flash column chromatography (hexanes:ethyl acetate = 10:1) to afford the desired product **Ethyl (Z)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyano-3-hydroxyacrylate (JXL083)** (yield: 80%, 1.93 g).

¹H NMR (500 MHz, CDCl₃) δ 14.61 (s, 1H), 8.67 (s, 1H), 8.31 (dd, *J* = 7.0 1.3 Hz, 1H), 7.84 (s, 1H), 7.57 (s, 2H), 7.33 (m, 2H), 7.21 (d, *J* = 7.4 Hz, 1H), 5.53 (s, 2H), 4.40 (q, *J* = 7.1 Hz, 2H), 1.41 (t, *J* = 7.1 Hz, 3H).

¹³C NMR (126 MHz, CDCl₃) δ 179.1, 172.2, 138.1, 136.0, 135.7, 132.7 (q, *J*_{C-F} = 33.6 Hz), 126.9, 124.5, 123.7, 123.6, 122.6, 122.9 (q, *J*_{C-F} = 273.4 Hz), 121.8, 118.3, 110.1, 109.6, 73.1, 62.3, 50.4, 14.3.

Experimental detail for the synthesis of JXL084




To a solution of **JXL083** (0.5 mmol, 240 mg) in dichloromethane (10 mL) was added pyridine (0.5 mmol, 40 μ L) and acetyl chloride (1.0 mmol, 84 μ L). The reaction mixture was stirred for 1h and TLC indicated that the reaction was complete. The reaction solvent was evaporated by flowing air over the open flask. The residue was purified by flash column chromatography (hexanes:ethyl acetate = 10:1) to afford the desired product **Ethyl (Z)-3-(1-(3,5-bis(trifluoro-methyl)benzyl)-1H-indol-3-yl)-2-cyanoacrylate (JXL084)** (yield: 86%, 225 mg).

^1H NMR (500 MHz, CDCl_3) δ 8.56 (s, 1H), 7.90 (m, 1H), 7.85 (s, 1H), 7.61 (s, 2H), 7.32 (m, 2H), 7.24 (m, 1H), 5.51 (s, 2H), 4.29 (q, $J = 7.1$ Hz, 2H), 2.48 (s, 3H), 1.36 (t, $J = 7.1$ Hz, 3H).

^{13}C NMR (126 MHz, CDCl_3) δ 166.9, 161.4, 137.7, 136.2, 135.5, 132.8 (q, $J_{\text{C-F}} = 33.9$ Hz), 127.0, 126.7, 124.6, 123.6, 122.8 (q, $J_{\text{C-F}} = 273.4$ Hz), 122.7, 121.8, 117.7, 110.7, 109.8, 89.3, 61.8, 50.5, 29.7, 21.3, 14.2.

Experimental detail for the synthesis of JXL085




To a solution of **JXL083** (0.5 mmol, 240 mg) in dichloromethane (10 mL) was added triethylamine (1.0 mmol, 139.5 μ L) and phosphoryl chloride (0.55 mmol, 520 μ L). The reaction mixture was stirred for 1h at reflux and TLC indicated that the reaction was complete. The reaction solvent was evaporated by flowing air over the open flask. The residue was

purified by flash column chromatography (hexanes:ethyl acetate = 10:1) to afford the desired product **Ethyl (Z)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-3-chloro-2-cyanoacrylate (JXL085)** (yield: 84%, 210 mg).

¹H NMR (500 MHz, CDCl₃) δ 7.91 (s, 1H), 7.85 (s, 1H), 7.77 (m, 1H), 7.65 (s, 2H), 7.63 (s, 1H), 7.30 (m, 2H), 5.48 (s, 2H), 4.18 (q, *J* = 7.1 Hz, 2H), 1.18 (t, *J* = 7.1 Hz, 3H).

¹³C NMR (126 MHz, CDCl₃) δ 161.0, 155.6, 138.1, 136.2, 135.4, 133.4, 132.7 (q, *J*_{C-F} = 33.7 Hz), 127.1, 126.6, 124.5, 124.3, 122.8 (q, *J*_{C-F} = 273.4 Hz), 121.5, 115.8, 122.1, 110.4, 102.5, 62.4, 50.0, 13.9.

Experimental detail for the synthesis of JXL086




Diethyl (E)-2-(1-(3,5-bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-1-cyanovinylphosphonate (JXL086)

To the solution of 1-(3,5-bis(trifluoromethyl)benzyl)-1H-indole-3-carboxaldehyde (1 mmol, 371 mg) in ethanol (3 mL) were added diethyl cyanomethylphosphate (1.3 equiv, 1.3 mmol, 204 μL) and L-proline (40 mol%, 0.4 mmol, 58 mg). The reaction was stirred at 50 °C for 24 h. After completion of the reaction as indicated by TLC, the reaction solvent was evaporated by flowing air over the open flask. The solid was purified by flash column chromatography (hexanes:ethyl acetate = 2:1) to afford the desired product **JXL086** (yield: 90%, 477 mg).

¹H NMR (500 MHz, CDCl₃) δ 8.55 (s, 1H), 8.33 (d, *J* = 19.7 Hz, 1H), 7.86 (d, *J* = 7.9 Hz, 1H), 7.83 (s, 1H), 7.58 (s, 2H), 7.31 (m, 2H), 7.22 (m, 1H), 5.54 (s, 2H), 4.21 (m, 4H), 1.39 (t, *J* = 7.0 Hz, 6H).

¹³C NMR (126 MHz, CDCl₃) δ 149.8, 138.3, 135.8, 132.7 (q, *J*_{C-F} = 33.7 Hz), 132.6, 128.2, 126.8, 124.5, 123.0, 122.9 (q, *J*_{C-F} = 273.4 Hz), 122.5, 119.0, 117.8 (d, *J*_{C-P} = 11.3 Hz), 112.2 (d, *J*_{C-P} = 18.9 Hz), 110.3, 91.6 (d, *J*_{C-P} = 207.9 Hz), 63.2, 50.4, 16.3.

JXL095 was synthesized by the similar route as **JXL086**.




Diethyl (E)-2-(1-(3,5-bis(trifluoromethyl)benzyl)-1H-pyrrolo[2,3-b]pyridin-3-yl)-1-cyanovinylphosphonate (JXL095)

¹H NMR (500 MHz, CDCl₃) δ 8.58 (s, 1H), 8.46 (dd, *J* = 4.7, 1.4 Hz, 1H), 8.24 (d, *J* = 19.5 Hz, 1H), 8.20 (dd, *J* = 8.0, 1.4 Hz, 1H), 7.82 (s, 1H), 7.77 (s, 2H), 7.32 (dd, *J* = 8.0, 4.7 Hz, 1H), 5.67 (s, 2H), 4.22 (m, 4H), 1.40 (t, *J* = 7.1 Hz, 6H).

¹³C NMR (126 MHz, CDCl₃) δ 149.2 (d, *J*_{C-P} = 8.2 Hz), 147.4, 145.6, 138.6, 132.4 (q, *J*_{C-F} = 33.7 Hz), 132.0, 127.9, 127.8, 123.0 (q, *J*_{C-F} = 273.4 Hz), 122.4, 119.9, 118.9, 117.4 (d, *J*_{C-P} = 11.3 Hz), 110.5 (d, *J*_{C-P} = 19.5 Hz), 92.8 (d, *J*_{C-P} = 205.1 Hz), 63.4, 48.3, 16.3.

Experimental detail for the synthesis of JXL096




(E)-2-(1-(3,5-Bis(trifluoromethyl)benzyl)-1H-indol-3-yl)-1-cyanovinylphosphonic acid (JXL096)

A solution of **JXL086** (30 mg, 0.057 mmol) in dichloromethane (2 mL) was cooled to 0 °C and bromotrimethylsilane (40 μL, 0.3 mmol) was added dropwise under argon. The mixture was warmed to 21 °C and stirred for 12 h. The solvent was evaporated under vacuum and the resulting residue was then dissolved in methanol (2 mL). The mixture was stirred at 21 °C for 2 h. Evaporation of all volatiles under vacuum gave the phosphoric acid **JXL096** (yield: 92%, 25 mg).

¹H NMR (500 MHz, CD₃OD) δ 8.58 (s, 1H), 8.25 (d, *J* = 19.6 Hz, 1H), 7.90 (s, 1H), 7.87 (m, 1H), 7.76 (s, 2H), 7.45 (m, 1H), 7.31 (m, 2H), 5.75 (s, 2H).

^{13}C NMR (126 MHz, CD_3OD) δ 146.7 ($J_{\text{c-p}} = 7.2$ Hz), 140.2, 136.1, 132.1, 131.9 (q, $J_{\text{c-f}} = 33.7$ Hz), 128.0, 127.2, 123.8, 122.3, 123.2 (q, $J_{\text{c-f}} = 273.4$ Hz), 121.4, 118.2, 117.4 (d, $J_{\text{c-p}} = 11.3$ Hz), 111.5 ($J_{\text{c-p}} = 18.4$ Hz), 110.5, 94.6 (d, $J_{\text{c-p}} = 201.2$ Hz), 49.1.

Experimental detail for the synthesis of JXL092



To a solution of methyl 1-(3,5-bis(trifluoromethyl)benzyl)-3-formyl-1H-indole-4-carboxylate (1 mmol, 429 mg) in ethanol (3 mL) were added *tert*-butyl 2-cyanoacetate (1.3 equiv, 1.3 mmol, 183 μL) and L-proline (40 mol%, 0.4 mmol, 58 mg). The reaction was stirred at 21 $^{\circ}\text{C}$ for 12 h and a yellow solid precipitated gradually. After completion of the reaction, ice-cold water (2 mL) was added into the reaction. The solid was separated by Buchner funnel filtration and washed with water (2 mL \times 3) and dried to afford the desired product. yield: 95%, 524 mg.


(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-4-(methoxycarbonyl)-1H-indol-3-yl)-2-cyanoacrylic acid (JXL092)

To the solution of methyl (*E*)-1-(3,5-bis(trifluoromethyl)benzyl)-3-(3-(*tert*-butoxy)-2-cyano-3-oxoprop-1-en-1-yl)-1H-indole-4-carboxylate (0.5 mmol, 276 mg) in dichloromethane (2 mL) was added trifluoroacetic acid (3 equiv, 1.5 mmol, 0.2 mL). The reaction mixture was stirred at 21 $^{\circ}\text{C}$ for 30 min and a yellow solid precipitated. After the reaction was complete as shown by TLC, the reaction solvent was evaporated by flowing air over the open flask. The solid was washed by 2 mL of solvent mixture (hexanes/EtOAc = 5:1) 5 to 10 times and monitored by TLC until all the non-polar impurities disappeared. Finally, the purity of the product was checked by NMR. yield: 90%, 223 mg.

^1H NMR (500 MHz, CDCl_3) δ 9.09 (s, 1H), 8.58 (s, 1H), 7.75 (d, $J = 7.5$ Hz, 1H), 7.68 (s, 1H), 7.47 (s, 2H), 7.34 (d, $J = 8.2$ Hz, 1H), 7.20 (app. t, $J = 7.9$ Hz, 1H), 5.51 (s, 2H), 3.87 (s, 3H).

^{13}C NMR (126 MHz, CDCl_3) δ 167.8, 165.1, 149.5, 138.2, 137.2, 134.8, 132.3 (q, $J_{\text{c-f}} = 33.7$ Hz), 126.8, 126.2, 125.5, 124.8, 123.3, 122.8 (q, $J_{\text{c-f}} = 273.4$ Hz), 122.3, 118.2, 114.7, 110.9, 97.2, 52.4, 50.2.

Experimental detail for the synthesis of JXL094



(*E*)-3-(1-(3,5-Bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylamide (JXL094)

The mixture of (*E*)-3-(1-(3,5-bis(trifluoromethyl)benzyl)-1*H*-indol-3-yl)-2-cyanoacrylic acid **JXL001** (0.1 mmol, 43.8 mg) and thionyl chloride (0.5 ml) was refluxed for 1 h. After concentration under vacuum, the resulting acyl chloride was dissolved in 1 ml benzene and 30% ammonia in water (1 ml) was added. The reaction mixture was stirred at 21 °C for 24 h. After completion of the reaction as indicated by TLC, the reaction solvent was evaporated under vacuum. The solid was purified by flash column chromatography (hexanes:ethyl acetate = 2:1) to afford the desired product **JXL094** (yield: 85%, 37 mg).

¹H NMR (500 MHz, CDCl₃) δ 8.70 (s, 1H), 8.47 (s, 1H), 7.92 (d, *J* = 7.1 Hz, 1H), 7.85 (s, 1H), 7.58 (s, 2H), 7.34 (m, 2H), 7.23 (m, 1H), 5.55 (s, 2H).

¹³C NMR (126 MHz, CDCl₃) δ 163.0, 144.8, 138.1, 136.0, 132.7 (q, *J*_{C-F} = 33.7 Hz), 132.5, 128.6, 126.8, 126.1, 124.6, 123.1, 122.9 (q, *J*_{C-F} = 273.4 Hz), 122.6, 119.3, 111.3, 110.3, 96.1, 50.4.

Additional exemplary compounds of the present invention can be prepared by methods analogous to those described above.

Example 2: Treatment of Epithelial Cells with Exemplary Compounds

To determine whether these compounds could promote cellular lactate production, we treated cultured epithelial cells with the compounds and measured lactate levels in the culture media using a Nova Biomedical BioProfile Basic Analyzer. Briefly, cultured epithelial cells were treated with DMSO, UK-5099 (also called JXL001), or certain of the exemplary compounds disclosed herein for 24-30 hours, and media lactate levels were measured and normalized to cell number and duration of the experiment to acquire a cellular lactate production rate (nmol lactate, million cells, hour).

Lactate production rates of treated cells are shown in FIGs. 8, 9, and 12. As expected based on the present disclosure, since they are UK-5099 analogues, most of the novel

compounds assayed increased lactate production. Furthermore, the total cell count following treatment with the UK-5099 analogues is shown in Figure 13. Most of the compounds were tolerated by the cells. A separate assay was performed to calculate the EC50 of some of the compounds as shown in FIG. 10.

Example 3: *In vivo* test of Exemplary Compounds

To determine the efficacy of the compounds on the hair cycle, mice were shaved at postnatal day 50, and topically treated every other day with a compound disclosed herein suspended in lotion in every other day for 2 weeks, and pictures were taken. As seen in FIG. 11, all the analogues that showed the ability to promote lactate production in the in vitro assay were also able to stimulate hair growth over the course of 2 weeks.

INCORPORATION BY REFERENCE

All publications and patents mentioned herein are hereby incorporated by reference in their entirety as if each individual publication or patent was specifically and individually indicated to be incorporated by reference. In case of conflict, the present application, including any definitions herein, will control.


EQUIVALENTS

While specific embodiments of the subject invention have been discussed, the above specification is illustrative and not restrictive. Many variations of the invention will become apparent to those skilled in the art upon review of this specification and the claims below. The full scope of the invention should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

CLAIMS


We claim:


1. A compound of formula I or II,



wherein:

each A is independently CH, CR⁴, or N;



Y is carboxyl, ester, amide, or  ;

Z is CH, CR⁴, or N.

R² is CN or carboxyl;

R³ is H, aryl, aralkyl, or aralkylacyl, and is optionally substituted by one or more R⁵, wherein

each R⁵ is independently selected from alkyl, alkoxy, or halo;

each instance of R⁴ is independently alkyl, carboxyl, halo, hydroxy, ester, or CN;

R⁷ is hydrogen, alkyl, halo, hydroxyl, alkoxy, or acyloxy;


R¹⁰ is hydrogen or alkyl; and

n is 0-4;


or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1, wherein Z is CH or N.

3. The compound of claim 1, wherein the compound is a compound of formula III,



wherein,



Y is carboxyl, ester, amide, or ;

R² is CN or carboxyl;

R³ is H, aryl, aralkyl, or aralkylacyl, and is optionally substituted by one or more R⁵, wherein

each R⁵ is independently selected from alkyl, alkoxy, or halo;

each instance of R⁴ is independently alkyl, carboxyl, halo, hydroxy, ester, or CN;

R⁶ is from H, alkyl, or cycloalkyl; and


R⁷ is hydrogen, alkyl, halo, hydroxyl, alkoxy, or acyloxy;

R¹⁰ is hydrogen or alkyl


R¹¹ is H or alkyl; and

n is 0-4;

or a pharmaceutically acceptable salt thereof.



4. The compound of any one of claims 1-3, wherein Y is
5. The compound of claim 4, wherein R¹⁰ is H.
6. The compound of claim 4, wherein R¹⁰ is alkyl (e.g., ethyl).
7. The compound of any one of claims 1-3, wherein Y is ester or amide.
8. The compound of any one of claims 3-7, wherein R¹¹ is alkyl (e.g., methyl).
9. The compound of claim 1, wherein the compound is a compound of formula V, VI, or VII,



wherein:

each A is independently CH, CR⁴, or N;

X is NR⁶ or O;

R¹ is H or lower alkyl; or either R¹ and R⁶ or R¹ and R², together with the atoms that separate them, complete a heterocycle;

R² is CN or carboxyl;

R³ is H, aryl, aralkyl, or aralkylacyl, and is optionally substituted by one or more R⁵, wherein each R⁵ is independently selected from alkyl, alkoxy, or halo;

each instance of R⁴ is independently alkyl, carboxyl, halo, hydroxy, or CN;

R⁶ is from H, alkyl, or cycloalkyl;

R⁷ is hydrogen, alkyl, halo, hydroxyl, alkoxy, or acyloxy; and


n is 0-4;

or a pharmaceutically acceptable salt thereof.

10. The compound of any one of claims 1-9, wherein at least one A is N.

11. The compound of any one of claims 1-10, wherein exactly one A is N.

12. The compound of any one of claims 1-11, wherein the compound is of formula Va, VIa, or VIIa:



wherein:

X is NR⁶ or O;

R¹ is H or lower alkyl;

R² is CN or carboxyl; or R¹ and R², together with the atoms that separate them, complete a heterocycle;

R³ is H, phenyl, or benzyl, and is optionally substituted by one or more R⁵, wherein each R⁵ is independently selected from alkyl, alkoxy, or halo;

each instance of R⁴ is independently selected from alkyl, carboxyl, halo, hydroxy, or CN; and

R⁶ is selected from H, alkyl, or cycloalkyl;


n is 0-4;

or a pharmaceutically acceptable salt thereof.

13. The compound of any one of claims 8-12, wherein X is NH.
14. The compound of any one of claims 8-12, wherein X is O.
15. The compound of any one of claims 9-12, wherein R¹ is H.
16. The compound of any one of claims 9-12, wherein R¹ is lower alkyl.
17. The compound of any one of claims 9-12, wherein R¹ and R⁶, together with the atoms that separate them, complete a heterocycle (e.g., morpholinyl).
18. The compound of any one of claims 9-13, wherein R⁶ is hydrogen.
19. The compound of any one of claims 9-18, wherein R² is CN.
20. The compound of any one of claims 9-18, wherein R² is carboxyl.
21. The compound of any one of claims 9-13, wherein R¹ and R², together with the atoms that separate them, complete a heterocyclyl selected from thiazolidine-2,4-dione-5-ylidene or 2-iminothiazolidin-4-one-5-ylidene.
22. The compound of any one of claims 12-21, wherein the compound is of formula Va.
23. The compound of any one of claims 12-21, wherein the compound is of formula VIa.
24. The compound of any one of claims 1-23, wherein R³ is H.
25. The compound of any one of claims 1-23, wherein R³ is phenyl.


26. The compound of any one of claims 1-23, wherein R^3 is phenyl and is substituted by one or more R^5 .
27. The compound of claim 26, wherein R^3 is substituted by one R^5 , and wherein R^5 is an alkoxy.
28. The compound of claim 26 or 27, wherein R^3 is aralkyl (e.g., benzyl or phenethyl).
29. The compound of claim 26 or 27, wherein R^3 is aralkyl.
30. The compound of claim 26 or 27, wherein R^3 is aralkylacyl (e.g., phenylacetyl).
31. The compound of claim 26 or 27, wherein R^3 is benzyl.
32. The compound of claim 26 or 27, wherein R^3 is benzyl and is substituted by one or more R^5 .
33. The compound of claim 26 or 27, wherein R^3 is aralkyl (e.g., benzyl or phenethyl) and is substituted by one or more R^5 (preferably on the phenyl ring).
34. The compound of claim 26 or 27, wherein R^3 is aralkylacyl (e.g., phenylacetyl), and is substituted by one or more R^5 (preferably on the phenyl ring).
35. The compound of any one of claims 32-33, wherein R^3 is substituted by one or two R^5 , and wherein each R^5 is independently selected from fluoroalkyl or fluoro.
36. The compound of claim 35, wherein R^3 is substituted by two R^5 , and wherein each R^5 is trifluoromethyl.

37. The compound of any one of claims 9-36, wherein the compound is represented by formula Vb:




38. The compound of any one of claims 12-37, wherein n is 0.

39. The compound of claim 38, wherein the compound is represented by formula Vc:




40. The compound of any one of claims 12-37, wherein n is 1.

41. The compound of claim 40, wherein the compound is represented by formula Vd:




42. The compound of claim 40, wherein the compound is represented by formula Ve:





43. The compound of any one of claims 40-42, wherein R⁴ is selected from halo or haloalkyl.


44. The compound of claim 43, wherein R^4 is halo (e.g., chloro or bromo).
45. The compound of any one of claims 9-21, wherein the compound is of formula VI.
46. The compound of any one of claims 9-21, wherein the compound is of formula VIa.
47. The compound of claim 45 or 46, wherein n is 0.
48. The compound of claim 45 or 46, wherein n is 2, and R^4 is selected from halo or haloalkyl.
49. The compound of any one of claims 9-21, wherein the compound is of formula VII.
50. The compound of any one of claims 9-21, wherein the compound is of formula VIIa.
51. The compound of any one of the preceding claims, wherein R^7 is hydrogen, hydroxyl, halo (e.g., chloro), or acyloxy (e.g., acetyloxy).
52. The compound of claim 51, wherein R^7 is hydroxyl, halo (e.g., chloro), or acyloxy (e.g., acetyloxy).

53. A compound selected from












,




,




,




,




,




,




,




,




,




,




,



,





54. A pharmaceutical composition comprising a compound of any one of the preceding claims and a pharmaceutically acceptable excipient.
55. The pharmaceutical composition of claim 54, wherein the pharmaceutical composition is formulated for topical administration.
56. A method of enhancing lactate production in a cell, comprising contacting the cell with an MPO inhibitor, such as an MPC inhibitor.
57. A method of enhancing lactate production in a cell, comprising contacting the cell with a compound or composition of any one of claims 1-55.
58. The method of claim 56 or 57, wherein the cell is a hair follicle stem cell.
59. A method of promoting hair growth, comprising administering to a patient a compound or composition of any one of claims 1-55.
60. A method of treating a condition or disorder affecting hair growth, comprising administering to a patient a compound or composition of any one of claims 1-55.
61. The method of claim 60, wherein the condition or disorder is baldness or alopecia.
62. A method of promoting hair growth, comprising administering to a patient (e.g., topically) an MPC inhibitor.
63. A method of treating a condition or disorder affecting hair growth, comprising administering to a patient (e.g., topically) an MPC inhibitor.
64. The method of claim 63, wherein the condition or disorder is baldness or alopecia.
65. The method of any one of claims 62-65, wherein the MPC inhibitor is a compound or composition of any one of claims 1-55.

66. A method of promoting hair growth, comprising administering to a patient (e.g., topically) an MPO inhibitor.
67. A method of treating a condition or disorder affecting hair growth, comprising administering to a patient (e.g., topically) an MPO inhibitor.
68. The method of claim 67, wherein the condition or disorder is baldness or alopecia.
69. The method of any one of claims 66-68, wherein the MPC inhibitor is a compound or composition of any one of claims 1-55.




FIG. 1A.
HFSC HFSCs




FIG. 1B.




FIG. 1C.




FIG. 1D.




FIG. 1E.




FIG. 2A.




FIG. 2B.





FIG. 2C.


HFSCs in telogen vs telo-ana (RNA-seq)



GENE SYMBOL	RANK IN GENE LIST	RANK METRIC SCORE
Ldha	341	3.39
Aldoa	501	2.90
Aldh1a3	869	2.30
Aldh3a2	1046	2.09
Hkdc1	1097	2.05
Pdhb	1191	1.95
Tpi1	1472	1.71
Fbp1	1511	1.68
Adh5	1735	1.54
Aldh7a1	2020	1.36
Gm5506	2210	1.28

FIG. 3A.

HFSCs in telogen vs telo-ana (microarray)



GENE SYMBOL	RANK IN GENE LIST	RANK METRIC SCORE
Pgam1	58	21.9
Pkm2	92	16.2
Pfkf	234	9.6
Gapdh	262	8.9
Ldha	342	7.4
Eno1	993	3.9
Hkdc1	1676	2.8
Pgm2	1735	2.7
Fbp1	1919	2.5
Adpgk	2265	2.2
Adh5	2265	2.1
Acss2	2418	2.0
Hk3	2734	1.9
Pgk1	2791	1.9
Pklr	3234	1.7

FIG. 3B.




FIG. 3C.







FIG. 3D.






FIG. 5A.




FIG. 5B.




FIG. 5C.




FIG. 5D.




FIG. 5E.




FIG. 5F.




FIG. 5G.




FIG. 6A.




FIG. 6B.




FIG. 6C.







FIG. 6D.






FIG. 7D.





FIG. 8






FIG. 10




FIG. 11




FIG. 12





FIG. 13


5/31/18 t=30 hr 10 uM treatment



t-test: *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001

FIG. 14

5/31/18 t=30 hr 10 uM treatment



t-test: *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001

FIG. 15

Relative LPR, JXL001-093




FIG. 16




FIG. 17

Relative LPR, JXL001, 067-093




FIG. 18

Relative cell count, JXL001, 067-093




FIG. 19




FIG. 20

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2018/040385

A. CLASSIFICATION OF SUBJECT MATTER

See extra sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC (2018.01) A61K 31/405, A61K 8/49, A61P 17/14, A61K 31/404, C07D 209/18, C07D 417/06, C07C 255/41, C07D 471/04, C07D 275/03, C07D 413/12, C07F 9/38

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Databases consulted: CAPLUS, MARPAT, REGISTRY, Google Scholar, DWPI

Search terms used: hair loss, baldness, alopecia, lactate, hair follicle stem cells, mitochondrial pyruvate carrier, MPC, MPO, UK5099.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2015049365 A2 INSERM [FR] 09 Apr 2015 (2015/04/09) claim 4, page 5 lines 7-10, pages 22-25, page 22 lines 21 and 26, page 22 line 17, page 23 line 16.	1,7,9,12,14,15,19, 22,25,54,55
Y	whole document	1-69
X	WO 2013169956 A2 UNIVERSITY OF UTAH RESEARCH FOUNDATION [US] 14 Nov 2013 (2013/11/14) [00285],[00293], [00294], [00300] and [00302]	1,7,9,12,14,15,19, 22,25,54-57
Y	whole document	1-69
X	Kyle S. et al. Mitochondrial pyruvate transport: a historical perspective and future research directions. Biochem J. 2015 Mar 15; 466(3): 443-454. Kyle S. et al. 15 Mar 2015 (2015/03/15) whole document	1,2,7,9,12,14-16,19, 22,23,25,45-47,51, 54-57
Y	whole document	1-69

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

“A” document defining the general state of the art which is not considered to be of particular relevance

“E” earlier application or patent but published on or after the international filing date

“L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

“O” document referring to an oral disclosure, use, exhibition or other means

“P” document published prior to the international filing date but later than the priority date claimed

“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

“&” document member of the same patent family

Date of the actual completion of the international search

18 Oct 2018

Date of mailing of the international search report

21 Oct 2018

Name and mailing address of the ISA:

Israel Patent Office

Technology Park, Bldg.5, Malcha, Jerusalem, 9695101, Israel

Facsimile No. 972-2-5651616

Authorized officer

VOLKOV Karina

Telephone No. 972-2-5651777

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2018/040385

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>Pyrrolo[2,3-b]pyridine derivatives (7-azaindoles) viii. synthesis and some reactions of 4-methyl-1-phenyl-1h-4-methyl-1-phenyl-1 h-pyrrolo[2,3-b]pyridine-3- carboxaldehydel. All-Union Chemical and Pharmaceutical Research Institute Translated from Zhurnal Obshchei Khimii, Vol. 34, No. 8, pp. 2603-2610.</p> <p>N. Yakhontov et al. 01 Aug 1964 (1964/08/01) page 2627 compound IX</p>	1,3,7,9-11,14,16,20,25,37,39-41,51
X	CAS Registry Number: 1246086-21-4, Entered STN: 12 Oct 2010. 12 Oct 2010 (2010/10/12)	1,3,7,24,40,42-44
X	CAS Registry Number: 1232821-01-0, Entered STN: 19 Jul 2010 19 Jul 2010 (2010/07/19)	1,3,7,24,40,42-44
X	WO 2013128465 A1 (CONNEXIOS LIFE SCIENCES PVT LTD?[IN]) 06 Sep 2013 (2013/09/06) page 106 compound 41	1,7,9,12,14,15,19,22,24,53
X	<p>AMIN, Sk Abdul, et al. Exploring structural requirements of unconventional Knoevenagel-type indole derivatives as anticancer agents through comparative QSAR modeling approaches. Canadian Journal of Chemistry, 2016, 94.7: 637-644.</p> <p>01 Jan 2016 (2016/01/01) page 32 Table 1 compound 2b</p>	1,4,6,24,54
X	CAS Registry Number: 2094959-90-5, Entered STN: 05 May 2017 05 May 2017 (2017/05/05)	1-3,7,9,12-14,16,18,19,49-51
X	CAS Registry Number: 2022941-63-3, Entered STN: 02 Nov 2016 02 Nov 2016 (2016/11/02)	1-3,7,9,12-14,16,18,19,49-51
X	CAS Registry Number: 1417368-01-4, Entered STN: 23 Jan 2013 23 Jan 2013 (2013/01/23)	1,2,7,9-12,17,24,37-39
X	CAS Registry Number: 677327-34-3, Entered STN: 28 Apr 2004. 28 Apr 2004 (2004/04/28)	1,2,7,9-12,17,24,37-39
X	CAS Registry Number: 1993795-68-8, Entered STN: 16 Sep 2016 16 Sep 2016 (2016/09/16)	1,7,9,12,14,28,31
X	CAS Registry Number: 1993738-37-6, Entered STN: 16 Sep 2016 16 Sep 2016 (2016/09/16)	1,7,9,12,14,28,31
X	CAS Registry Number: 1360583-78-3, Entered STN: 09 Mar 2012 09 Mar 2012 (2012/03/09)	1,7,9,12,20,21,28,31-33

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2018/040385

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CAS Registry Number: 1360583-75-0, Entered STN: 09 Mar 2012 09 Mar 2012 (2012/03/09)	1,7,9,12,20,21,28, 31-33
X	WO 2004080481 A1 (NOVO NORDISK AS?[DK]) 23 Sep 2004 (2004/09/23) compounds 446,449,464,465	1,12,21,24
X	CAS Registry Number: 895304-22-0, Entered STN: 23 Jul 2006 23 Jul 2006 (2006/07/23)	1,9,12,28,29,31-33, 35
X	CAS Registry Number: 895303-82-9, Entered STN: 23 Jul 2006 23 Jul 2006 (2006/07/23)	1,9,12,28,29,31-33, 35
X	CAS Registry Number: 1025594-30-2, Entered STN: 05 Jun 2008. 05 Jun 2008 (2008/06/05)	1,9,12,30,34
X	CAS Registry Number: 904141-89-5, Entered STN: 24 Aug 2006 24 Aug 2006 (2006/08/24)	1,9,12,30,34
Y	Vishnyakova, K. S. et al. Possible role of autophagy activation in stimulation of regeneration. Molecular Biology (01 September 2013), 47(5), pages: 692-700. Retrieved from URL: < http://dx.booksc.org/22800000/libgen.scimag22819000-22819999.zip/browse/10.1134/S002689331305021X.pdf >. 01 Sep 2013 (2013/09/01) page 693	1-69
P,X	Flores, A. et al. Lactate dehydrogenase activity drives hair follicle stem cell activation. Nature cell biology (published online 14 August 2017), volume 19, issue 9, pages: 1017-1026. 14 Aug 2017 (2017/08/14) whole document	1-69
P,X	WO 2018039612 A1 (UNIV CALIFORNIA [US]) 01 Mar 2018 (2018/03/01) whole document	1-69

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/US2018/040385

Patent document cited search report	Publication date	Patent family member(s)	Publication Date
WO 2015049365 A2	09 Apr 2015	WO 2015049365 A2	09 Apr 2015
		WO 2015049365 A3	01 Oct 2015
		EP 3052133 A2	10 Aug 2016
		US 2016250249 A1	01 Sep 2016
<hr/>			
WO 2013169956 A2	14 Nov 2013	WO 2013169956 A2	14 Nov 2013
		WO 2013169956 A3	13 Mar 2014
		EP 2847356 A2	18 Mar 2015
		EP 2847356 A4	05 Aug 2015
		US 2013339749 A1	19 Dec 2013
		US 2015099790 A1	09 Apr 2015
		US 2016193364 A1	07 Jul 2016
<hr/>			
WO 2013128465 A1	06 Sep 2013	WO 2013128465 A1	06 Sep 2013
		AU 2012372019 A1	24 Jul 2014
		AU 2012372019 B2	15 Jun 2017
		BR 112014015301 A2	13 Jun 2017
		BR 112014015301 A8	04 Jul 2017
		CA 2860187 A1	06 Sep 2013
		CL 2014001704 A1	13 Mar 2015
		CN 104125956 A	29 Oct 2014
		CO 7160013 A2	15 Jan 2015
		DK 2800748 T3	17 Jul 2017
		EP 2800748 A1	12 Nov 2014
		EP 2800748 B1	29 Mar 2017
		ES 2630309 T3	21 Aug 2017
		IL 233270 D0	31 Aug 2014
		JP 2015521154 A	27 Jul 2015
		JP 6118340 B2	19 Apr 2017
		KR 20140127226 A	03 Nov 2014
		MA 35874 B1	01 Dec 2014

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2018/040385

Patent document cited search report	Publication date	Patent family member(s)	Publication Date
		MX 2014007719 A	19 Mar 2015
		MX 354361 B	28 Feb 2018
		NZ 626745 A	24 Jun 2016
		PH 12014501442 A1	08 Oct 2014
		PH 12014501442 B1	08 Oct 2014
		RU 2014126646 A	10 Feb 2016
		SG 11201403268Q A	30 Jul 2014
		US 2015158860 A1	11 Jun 2015
		US 9453014 B2	27 Sep 2016
		ZA 201405263 B	25 May 2016
<hr/>			
WO 2004080481 A1	23 Sep 2004	WO 2004080481 A1	23 Sep 2004
		EP 1605967 A1	21 Dec 2005
		JP 2006519791 A	31 Aug 2006
		US 2006258561 A1	16 Nov 2006
<hr/>			
WO 2018039612 A1	01 Mar 2018	WO 2018039612 A1	01 Mar 2018
		WO 2018039615 A1	01 Mar 2018
<hr/>			

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2018/040385

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (2018.01) A61K 31/405, A61K 8/49, A61P 17/14, A61K 31/404, C07D 209/18, C07D 417/06, C07C 255/41, C07D 471/04, C07D 275/03, C07D 413/12, C07F 9/38